

Exercises for Stochastic Processes

Tutorial exercises:

T1. Let $S = \{0, 1\}$. Consider the general Q -matrix

$$\begin{pmatrix} -\beta & \beta \\ \delta & -\delta \end{pmatrix}, \quad (1)$$

for some $\beta, \delta \geq 0$. Show that the corresponding transition probabilities are $p_t(x, y) = \mathbb{1}_{\{x=y\}}$ if $\beta + \delta = 0$, and otherwise they are given by

$$\begin{aligned} p_t(0, 0) &= \frac{\delta}{\beta + \delta} + \frac{\beta}{\beta + \delta} e^{-t(\beta + \delta)}, & p_t(0, 1) &= \frac{\beta}{\beta + \delta} (1 - e^{-t(\beta + \delta)}), \\ p_t(1, 1) &= \frac{\beta}{\beta + \delta} + \frac{\delta}{\beta + \delta} e^{-t(\beta + \delta)}, & p_t(1, 0) &= \frac{\delta}{\beta + \delta} (1 - e^{-t(\beta + \delta)}). \end{aligned} \quad (2)$$

T2. Consider the following stochastic process $X(t)$ on $\{0, 1\}$. If the process is in 0 it stays in this state for an exponential distributed time with parameter β and then jumps to state 1. If the process is in state 1 it stays in this state for an exponential distributed time with parameter δ and then goes to 0. Let $p_t(i, j)$ be the probability that $X(t) = j$ if $X(0) = i$.

(a) Show that

$$p_t(0, 1) = \int_0^t \beta e^{-\beta s} p_{t-s}(1, 1) ds,$$

and

$$p_t(1, 0) = \int_0^t \delta e^{-\delta s} p_{t-s}(0, 0) ds,$$

(b) Show that the Q -matrix for this process is the same as in (1), so that the transition probabilities for this process are given in (2).

T3. With the notations used in the lecture in the probabilistic construction of a Markov chain with a given Q -matrix, show that the following statements are equivalent:

- (a) $\mathbb{P}(N(t) < \infty) = 1$ for all $t \geq 0$.
- (b) $\sum \tau_n = \infty$ a.s.
- (c) $\sum \frac{1}{c(Z_n)} = \infty$ a.s.

Homework exercises:

H1. Let Q be a Q-matrix on a finite state space. Show that $p_t(x, y)$ defined by

$$P_t := \sum_{k=0}^{\infty} \frac{t^k Q^k}{k!}$$

is a transition function and show that

$$q(x, y) = \left. \frac{d}{dt} p_t(x, y) \right|_{t=0}.$$

H2. Show that, for any continuous time Markov chain with deterministic starting point, the time of the first jump has an exponential distribution (possibly with parameter 0 or ∞). Does this generalize to arbitrary initial distributions?

H3. Let (X_n) be a sequence of independent continuous time Markov chains on $\{0, 1\}$ with Q-matrices $\begin{pmatrix} -\beta & \beta \\ \delta & -\delta \end{pmatrix}$. Assume that $\sum \frac{\beta_n}{\beta_n + \delta_n} < \infty$. Define

$$X(t) := (X_1(t), X_2(t), \dots)$$

and

$$S := \left\{ x \in \{0, 1\}^{\mathbb{N}} \mid \sum x_n < \infty \right\}.$$

- (a) Show that S is countable and $\mathbb{P}(X(t) \in S \mid X(0) \in S) = 1$!
- (b) Show that $p_t(x, y) := \mathbb{P}(X(t) = y \mid X(0) = x)$ is a transition function on S !
- (c) Assume that, moreover, $\sum \beta_n = \infty$. Show that $c(x) = \infty$ for all $x \in S$!
- (d) Show that, for any $x \in S$ and $\epsilon > 0$,

$$\mathbb{P}^x(X(t) = x \text{ for all } t < \epsilon) = 0.$$

Conclude that there is no Markov chain with transition function p !

Deadline: Monday, 5.12.16