Percolation and number of phases in the two-dimensional lsing model

Hans-Otto Georgii^{a)}

Mathematisches Institut der Universität München, Theresienstr. 39, D-80333 München, Germany

Yasunari Higuchi^{b)}

Department of Mathematics, Faculty of Science, Kobe University, Rokko, Kobe 657-8501, Japan

(Received 18 November 1999; accepted for publication 30 November 1999)

We reconsider the percolation approach of Russo, Aizenman, and Higuchi for showing that there exist only two phases in the Ising model on the square lattice. We give a fairly short alternative proof which is only based on stochastic monotonicity and avoids the use of symmetry inequalities originally needed for some background results. Our proof extends to the Ising model on other planar lattices such as the triangular and honeycomb lattice. We can also treat the Ising antiferromagnet in a homogeneous field and the hard-core lattice gas model on \mathbb{Z}^2 . © 2000 American Institute of Physics. [S0022-2488(00)00103-1]

I. INTRODUCTION

One of the fundamental results on the two-dimensional ferromagnetic Ising model is the following theorem obtained independently in the late 1970s by Aizenman¹ and Higuchi² on the basis of the seminal work of Russo.³

Theorem: For the ferromagnetic Ising model on \mathbb{Z}^2 with no external field and inverse temperature $\beta > \beta_c$, there exist only two distinct extremal Gibbs measures μ^+ and μ^- .

The basic technique initiated by Russo consists of an interplay of three features of the Ising model:

- (1) the strong Markov property for random sets defined by geometric conditions involving clusters of constant spin,
- (2) the symmetry of the interaction under spin-flip and lattice automorphisms, and
- (3) the ferromagnetic character of the interaction which manifests itself in FKG order and positive correlations.

These ingredients led to a detailed understanding of the geometric features of typical configurations as described by the concepts of percolation theory. In addition to these tools, the authors of Refs. 1–3 also needed the result that the limiting Gibbs measure with \pm boundary condition is a mixture of the two pure phases. This result of Messager and Miracle-Solé⁴ had been had been obtained by quite different means, namely some correlation inequalities of symmetry type in the spirit of Griffiths–Kelly–Sherman (GKS) and Lebowitz inequalities. While such symmetry inequalities are a beautiful and powerful tool, they are quite different in character from the Fortvín– Kasteleyn–Ginibre (FKG) inequality and have their own restrictions. It is therefore natural to ask whether Russo's random cluster method is flexible enough to prove the theorem without recourse to symmetry inequalities. On the one hand, this would allow one to extend the theorem to models with less symmetries, while on the other hand one might gain a deeper understanding of possible geometric features of typical configurations.

In this paper we propose such a purely geometric reasoning which is only based on the three above-mentioned features and avoids the use of the symmetry inequalities of Messager and

0022-2488/2000/41(3)/1153/17/\$17.00

1153

© 2000 American Institute of Physics

^{a)}Electronic mail: georgii@rz.mathematik.uni-muenchen.de

^{b)}Electric mail: higuchi@math.kobe-u.ac.jp

Miracle-Solé.⁴ Despite this reduction of tools we could simplify the proof by an efficient combination of known geometric arguments. These include:

- (1) the Burton-Keane uniqueness theorem for infinite clusters,⁵
- (2) a version of Zhang's argument for the impossibility of simultaneous plus- and minuspercolation in \mathbb{Z}^2 (cf. Theorem 5.18 of Ref. 6),
- (3) Russo's symmetry trick for simultaneous flipping of spins and reflection of the lattice, 3 and
- (4) Aizenman's idea of looking at contour intersections in a duplicated system.¹

We have tried to keep the paper reasonably self-contained, so that the reader will find a complete proof of the theorem. As a payoff of the method we also obtain some generalizations. On the one hand, the arguments carry over to the Ising model on other planar lattices such as the triangular or the hexagonal lattice. On the other hand, in the case of the square lattice they also cover the antiferromagnetic Ising model in an external field as well as the hard-core lattice gas model.

II. SETUP AND BASIC FACTS

Although we assume that the reader is familiar with the definition of the Ising model, let us start by recalling a number of fundamental facts and introducing some notations. We assume throughout that the inverse temperature β exceeds the Onsager threshold β_c , and that there is no external field, h=0. The main ingredients we need are as follows.

(1) The configuration space $\Omega = \{-1,1\}^{\mathbb{Z}^2}$, which is equipped with the Borel σ -algebra \mathcal{F} and the local σ -algebras \mathcal{F}_{Λ} of events depending only on the spins in $\Lambda \subset \mathbb{Z}^2$.

(2) The *Gibbs distributions* μ_{Λ}^{ω} in finite regions $\Lambda \subset \mathbb{Z}^2$ with boundary condition $\omega \in \Omega$. These enjoy the *Markov property*, which says that $\mu_{\Lambda}^{\omega}(A)$ for $A \in \mathcal{F}_{\Lambda}$ depends only on the restriction of ω to the boundary $\partial \Lambda = \{x \notin \Lambda : |x-y| = 1 \text{ for some } y \in \Lambda\}$ of Λ , and the *finite energy property*, which states that $\mu_{\Lambda}^{\omega}(A) > 0$ when $\emptyset \neq A \in \mathcal{F}_{\Lambda}$.

(3) The *Gibbs measures* μ on (Ω, \mathcal{F}) which, by definition, satisfy $\mu(\cdot | \mathcal{F}_{\Lambda^c})(\omega) = \mu_{\Lambda}^{\omega}$ for μ -almost all ω and any finite Λ ; we write \mathcal{G} for the convex set of all Gibbs measures and ex \mathcal{G} for the set of all extremal Gibbs measures.

(4) The *strong Markov property* of Gibbs measures, stating that $\mu(\cdot | \mathcal{F}_{\Gamma^c})(\omega) = \mu_{\Gamma(\omega)}^{\omega}$ for μ -almost all ω when Γ is any finite *random* subset of \mathbb{Z}^2 which is *determined from outside*, in that $\{\Gamma = \Lambda\} \in \mathcal{F}_{\Lambda^c}$ for all finite Λ , and \mathcal{F}_{Γ^c} is the σ -algebra of all events Λ outside Γ in the sense that $\Lambda \cap \{\Gamma = \Lambda\} \in \mathcal{F}_{\Lambda^c}$ for all finite Λ . (Using the conventions $\mu_{\emptyset}^{\omega} = \delta_{\omega}$ and $\mathcal{F}_{\emptyset^c} = \mathcal{F}$ we can in fact allow that Γ takes the value \emptyset .) For a proof one simply splits Ω into the disjoint sets $\{\Gamma = \Lambda\}$ for finite Λ .

(5) The *stochastic monotonicity* (or FKG order) of Gibbs distributions. Writing $\mu \leq \nu$ when $\mu(f) \leq \nu(f)$ for all increasing local (or, equivalently, all increasing bounded measurable) real functions f on Ω , we have $\mu_{\Lambda}^{\omega} \leq \mu_{\Lambda}^{\omega'}$ when $\omega \leq \omega'$, and $\mu_{\Lambda}^{\omega} \leq \mu_{\Delta}^{\omega}$ when $\Delta \subset \Lambda$ and $\omega \equiv +1$ on $\Lambda \setminus \Delta$ (the opposite relation holds when $\omega \equiv -1$ on $\Lambda \setminus \Delta$).

(6) The *pure phases* $\mu^+, \mu^- \in \mathcal{G}$ obtained as limits for $\Lambda \uparrow \mathbb{Z}^2$ of μ_{Λ}^{ω} with $\omega \equiv +1$ and -1, respectively, their invariance under all graph automorphisms of \mathbb{Z}^2 , the sandwich relation $\mu^- \leq \mu \leq \mu^+$ for any other $\mu \in \mathcal{G}$, and the resulting extremality of μ^+ and μ^- .

(7) The characterization of extremal Gibbs measures by their *triviality on the tail* σ -algebra $\mathcal{T}=\cap{\{\mathcal{F}_{\Lambda c}:\Lambda\subset \mathbb{Z}^2 \text{ finite}\}};$ the fact that extremal Gibbs measures have *positive correlations*; and the *extremal decomposition* representing any Gibbs measure as the barycenter of a mass distribution on ex \mathcal{G} .

A general account of Gibbs measures can be found in Ref. 7, and Ref. 6 contains an exposition of the Ising model and its properties related to stochastic monotonicity.

We will also use a class of transformations of Ω which preserve the Ising Hamiltonian, and thereby the class \mathcal{G} of Gibbs measures. These transformations are as follows.

(1) The spin-flip transformation $T: \omega = (\omega(x))_{x \in \mathbb{Z}^2} \rightarrow (-\omega(x))_{x \in \mathbb{Z}^2}$.

(2) The *translations* $\vartheta_x, x \in \mathbb{Z}^2$, which are defined by $\vartheta_x \omega(y) = \omega(y-x)$ for $y \in \mathbb{Z}^2$, and in particular the horizontal and vertical shifts $\vartheta_{\text{hor}} = \vartheta_{(1,0)}$ and $\vartheta_{\text{vert}} = \vartheta_{(0,1)}$, respectively.

(3) The *reflections* in lines *l* through lattice sites: For any $k \in \mathbb{Z}$ we write

$$R_{k,\text{hor}}: \mathbb{Z}^2 \ni x = (x_1, x_2) \longrightarrow (x_1, 2k - x_2) \in \mathbb{Z}^2$$

for the reflection in the horizontal line $\{x_2=k\}$, and similarly $R_{k,vert}$ for the reflection in the vertical line $\{x_1=k\}$. For k=0 we simply write $R_{hor}=R_{0,hor}$ and $R_{vert}=R_{0,vert}$. All these reflections act canonically on Ω .

We will investigate the geometric behavior of typical configurations in *half-planes* of \mathbb{Z}^2 . These are sets of the form

$$\pi = \{x = (x_1, x_2) \in \mathbb{Z}^2 : \mathbf{z}^2 : x_i \ge k\}$$

with $k \in \mathbb{Z}, i \in \{1,2\}$, or with " \geq " replaced by " \leq ." The line $l = \{x \in \mathbb{Z}^2 : x_i = k\}$ is called the associated *boundary line*. In particular, we will consider the following

(1) The upper half-plane $\pi_{up} = \{x = (x_1, x_2) \in \mathbb{Z}^2 : x_2 \ge 0\}.$

(2) The downwards half-plane $\pi_{\text{down}} = \{x = (x_1, x_2) \in \mathbb{Z}^2 : x_2 \leq 0\}$, and the analogously defined right half-plane π_{right} and left half-plane π_{left} . We will also work with

(3) The left horizontal semiaxis $l_{\text{left}} = \{x = (x_1, x_2) \in \mathbb{Z}^2 : x_1 \leq 0, x_2 = 0\}$, and

(4) the right semiaxis $l_{\text{right}} = \{x = (x_1, x_2) \in \mathbb{Z}^2 : x_1 \ge 0, x_2 = 0\}.$

In the rest of this section we state three fundamental results on percolation in the Ising model. By the symmetry between the spin values +1 and -1, these results also hold when the minus and plus signs are interchanged. Similarly, all notations introduced with one sign will be used accordingly for the opposite sign.

We first recall some basic concepts of percolation theory. A finite *path* is a sequence $p = (x_1, x_2, ..., x_k)$ of pairwise distinct lattice points such that, for any $i \in \{2,...,k\}$, x_{i-1} and x_i are nearest neighbors (i.e., have Euclidean distance 1). The number k is called the *length* of p, and x_1 and x_k are its starting and final point, respectively. A path p is called a *path in a subset* $S \subset \mathbb{Z}^2$ if all x_i belong to S. We say that p meets or touches S if some x_i is contained in S or a nearest neighbor of a point in S. We will also speak of infinite paths $(x_1, x_2, ...)$ and doubly infinite paths $(..., x_{-1}, x_0, x_1, ...)$ in the obvious sense. A path p is called a *circuit* if x_1 and x_k are nearest neighbors, and a *semicircuit* in a half-plane π if it is contained in π and x_1 and x_k belong to the boundary line of π . A region $C \subset \mathbb{Z}^2$ is called *connected* if for any $x, y \in C$ there exists a path in C from x to y. A *cluster* in a region $S \subseteq \mathbb{Z}^2$ is a maximal connected subset C of S. It is called infinite if its cardinality is infinite. Infinite clusters will be denoted by the letter I, with suitable sub- and superscripts.

Given any configuration $\omega \in \Omega$, we consider the set $S^+(\omega) = \{x \in \mathbb{Z}^2 : \omega(x) = +1\}$ of + spins. A path (respectively, circuit, semicircuit, cluster) in $S^+(\omega)$ is called a +*path* (respectively, +*circuit*, +*semicircuit*, +*cluster*) for ω , and two points x, y are said to be +*connected* if there exists a +path from x to y.

We also need to work with the conjugate graph structure on \mathbb{Z}^2 , for which two points are considered as neighbors if their Euclidean distance is either 1 or $\sqrt{2}$, i.e., if they are either nearest neighbors or diagonal neighbors. This graph structure is indicated by a star and leads to the concepts of *paths, *circuits, *semicircuits, *connectedness, *clusters, +*paths, +*semicircuits, and so on. Note that each path is *a fortiori* a *path, and each cluster is contained in some *cluster.

The starting point of the random cluster method is the following result of Refs. 8 and 3. Let E^+ denote the event that there exists an infinite +cluster I^+ in \mathbb{Z}^2 , and define E^-, E^{+*}, E^{-*} analogously. Note that $E^+ \subset E^{+*}$ and $E^- \subset E^{-*}$. (Throughout this paper we will use the letter E to denote events concerning the existence of infinite clusters.)

Lemma 2.1 (Existence of infinite clusters): If $\mu \in \mathcal{G}$ is different from μ^- , there exists with positive probability an infinite + cluster. That is, $\mu(E^+) > 0$ when $\mu \neq \mu^-$.

Proof: Suppose that $\mu(E^+)=0$. Then any given square Δ is almost surely surrounded by a -*circuit, and with probability close to 1 such a circuit can already be found within a square

 $\Lambda \supset \Delta$ provided Λ is large enough. If this occurs, we let Γ be the largest random subset of Λ which is the interior of such a -*circuit. (The largest such set exists because the union of such sets is again the interior of a -*circuit.) In the alternative case we set $\Gamma = \emptyset$. By maximality, Γ is determined from outside. The strong Markov property together with the stochastic monotonicity $\mu_{\Gamma}^{-} \leq \mu^{-}$ therefore implies (in the limit $\Lambda \uparrow \mathbb{Z}^{2}$) that $\mu \leq \mu^{-}$ on \mathcal{F}_{Δ} . Since Δ was arbitrary and μ^{-} is minimal we find that $\mu = \mu^{-}$, and the lemma is proved.

The next lemma is a variant of another result of Russo.³

Lemma 2.2 (Flip-reflection domination): Let $\mu \in \mathcal{G}$ and R be any reflection, and suppose that for μ -almost all ω each finite $\Delta \subset \mathbb{Z}^2$ is surrounded by an R-invariant *circuit c such that $\omega \ge R \circ T(\omega)$ on c. Then $\mu \ge \mu \circ R \circ T$.

Proof: Another way of stating the assumption is that for any finite *R*-invariant Δ and μ -almost all ω there exists a finite *R*-invariant random set $\Gamma(\omega) \supset \Delta$ such that $\omega \ge R \circ T(\omega)$ on $\partial \Gamma(\omega)$. Given any $\epsilon > 0$, we can thus find an *R*-invariant Λ so large that with probability at least $1 - \epsilon$ such an *R*-invariant $\Gamma(\omega)$ exists within Λ . Since the union of any two such $\Gamma(\omega)$'s enjoys the same properties, we can assume that $\Gamma(\omega)$ is chosen maximal in Λ ; in the case when no such $\Gamma(\omega)$ exists we set $\Gamma(\omega) = \emptyset$. The maximality of Γ implies that the events { $\Gamma = G$ } are measurable with respect to $\mathcal{F}_{\Lambda \setminus G}$. For any increasing \mathcal{F}_{Δ} -measurable function $f \ge 0$ we thus get from the strong Markov property

$$\mu(f) \ge \mu(\mu_{\Gamma}(f) \mathbf{1}_{\{\Gamma \neq \emptyset\}}).$$

However, if $\Gamma(\omega) \neq \emptyset$ then $\omega \ge R \circ T(\omega)$ on $\partial \Gamma(\omega)$. By stochastic monotonicity, for such ω we have

$$\mu^{\omega}_{\Gamma(\omega)}(f) \geq \mu^{R \circ T(\omega)}_{\Gamma(\omega)}(f) = \mu^{\omega}_{\Gamma(\omega)}(f \circ R \circ T),$$

where the identity follows from the *R*-invariance of Γ and the *R* \circ *T*-invariance of the interaction. Hence

$$\mu(f) \ge \mu(f \circ R \circ T \mathbf{1}_{\{\Gamma \neq \emptyset\}}) \ge \mu(f \circ R \circ T) - \boldsymbol{\epsilon} \|f\|_{\infty}.$$

The lemma thus follows by letting $\epsilon \rightarrow 0$ and $\Delta \uparrow \mathbb{Z}^2$.

A third useful result of Russo³ is the following. To state it we need to introduce two notations. First, let

$$\theta = \mu^+ (0 \in I^{+*})$$

be the μ^+ -probability that the origin belongs to an infinite +*cluster. Lemma 2.1 implies that $\theta > 0$. Second, for a half-plane π with boundary line *l* and a *semicircuit σ in π we write Int σ for the unique subset of \mathbb{Z}^2 which is invariant under the reflection *R* in *l* and satisfies $\pi \cap \partial(\operatorname{Int} \sigma) = \sigma$; we call Int σ the interior of σ .

Lemma 2.3 (Point-to-semicircuit lemma): Let π be some half-plane with boundary line $l, x \in l$, and σa *semicircuit in π with interior $\Lambda = \operatorname{Int} \sigma \Im x$. Let $\omega \in \Omega$ be such that $\omega \equiv +1$ on σ . Then

 $\mu^{\omega}_{\Lambda}(x \text{ is in } \Lambda + * \text{connected to } \sigma) \ge \theta/2.$

Proof: By stochastic monotonicity we can assume that $\omega \equiv -1$ on $\partial \Lambda \setminus \sigma$. We then have $\omega \ge R \circ T(\omega)$ on $\partial \Lambda$, and therefore $\mu_{\Lambda}^{\omega} \ge \mu_{\Lambda}^{\omega} \circ R \circ T$. To exploit this relation we let $B_{x,\sigma}$ be the event that there exists a +*path in Λ from x to σ , $C_{x,\sigma}$ the event that x is surrounded by a +*circuit in $\Lambda \cup \sigma$ which is +*connected to σ , and $D_{x,\sigma} = B_{x,\sigma} \cup C_{x,\sigma}$. Then

$$\mu^{\omega}_{\Lambda}(D_{x,\sigma} \cup RT(D_{x,\sigma})) = 1. \tag{1}$$

Indeed, suppose that $\omega(x) = +1$, but $B_{x,\sigma}$ does not occur. Then the +*cluster containing x does not meet σ . Its outer boundary belongs to a -*cluster, which either touches $R(\sigma)$ so that

 $R \circ T(C_{x,\sigma})$ occurs, or not—in which case we consider the +*cluster containing its outer boundary, and so on. After finitely many steps we see that either $C_{x,\sigma}$ or $R \circ T(C_{x,\sigma})$ must occur. (1) is an immediate consequence. It follows that $\mu_{\Lambda}^{\omega}(D_{x,\sigma}) \ge 1/2$. Hence

$$\mu_{\Lambda}^{\omega}(B_{x,\sigma}) \geq \mu_{\Lambda}^{\omega}(B_{x,\sigma}|C_{x,\sigma}) \mu(D_{x,\sigma}) \geq \mu_{\Lambda}^{\omega}(B_{x,\sigma}|C_{x,\sigma})/2.$$

But if $C_{x,\sigma}$ occurs then there exists a largest random set $\Gamma \subset \Lambda$ containing x such that $\partial \Gamma$ forms a +*circuit and is +*connected to σ . Writing $B_{x,\partial\Gamma}$ for the event that x is +*connected to $\partial\Gamma$ and using the strong Markov property we thus find that

$$\mu_{\Lambda}^{\omega}(B_{x,\sigma}|C_{x,\sigma}) = \mu_{\Lambda}^{\omega}(\mu_{\Gamma}^{+}(B_{x,\partial\Gamma})|C_{x,\sigma}) \ge \theta$$

because $\mu_{\Gamma}^+(B_{x,\partial\Gamma}) \ge \theta$ by stochastic monotonicity. Together with the previous inequality this gives the result.

III. PERCOLATION IN HALF-PLANES

In this section we will prove that there exist plenty of infinite clusters of constant spin in the half-planes of \mathbb{Z}^2 . In particular, this will show that all translation invariant $\mu \in \mathcal{G}$ are mixtures of μ^+ and μ^- . We will use two pearls of percolation theory, the Burton–Keane uniqueness theorem for infinite clusters⁵ and Zhang's argument for the nonexistence of two infinite clusters of opposite sign in \mathbb{Z}^2 . (In the present context, these two results were obtained first in Ref. 9).

For a given half-plane π we let E_{π}^+ denote the event that there exists an infinite +cluster in π . When this occurs, we will write I_{π}^+ for such an infinite +cluster in π . (As we will see, such clusters are unique, so that this notation does not lead to conflicts.) In the case of the standard half-planes, we will only keep the directional index and omit the π ; for example, we write E_{up}^+ for $E_{\pi_{up}}^+$. Similar notations will be used for +*clusters and for the minus sign instead of the plus sign.

Let us say that (π, π') is a pair of *conjugate half-planes* if π and π' share only a common boundary line. An associated pair $(I_{\pi}^+, I_{\pi'}^+)$ or $(I_{\pi}^-, I_{\pi'}^-)$ of infinite clusters of the same sign in π and π' will be called an *infinite butterfly*. (This name alludes to the assumption that the two infinite "wings" have the same "color," but is not meant to suggest that they are symmetric and connected to each other, although the latter will turn out to be true.) We will say that a statement holds \mathcal{G} -almost surely if it holds μ -almost surely for all $\mu \in \mathcal{G}$.

Lemma 3.1 (Butterfly lemma): G-almost surely there exists at least one infinite butterfly.

Proof: Suppose the contrary. By the extremal decomposition theorem and the fact that the existence of infinite butterflies is a tail measurable event, there is then some $\mu \in \operatorname{ex} \mathcal{G}$ for which there exists no infinite butterfly μ -almost surely. We will show that this is impossible.

Step 1. First we observe that μ is $R \circ T$ -invariant for all reflections $R = R_{k,hor}$ or $R_{k,vert}$, and in particular is periodic under translations. Indeed, let (π, π') be conjugate half-planes with common boundary line l and R the reflection in l mapping π onto π' . By the absence of infinite butterflies, at least one of the half-planes π and π' contains no infinite –cluster, and this or the other half-plane contains no infinite +cluster. In view of the tail triviality of μ , we can assume that $\mu(E_{\pi}^{-})=0$. This means that for μ -almost all ω every finite $\Delta \subset \pi$ is surrounded by some +*semicircuit γ in π . For such a $\gamma, c = \gamma \cup R(\gamma)$ is an R-invariant *circuit that surrounds $\Delta \cup R(\Delta)$ and satisfies $\omega \ge R \circ T(\omega)$ on c. By Lemma 2.2, this gives the flip-reflection domination $\mu \ge \mu \circ R \circ T$. Since also $\mu(E_{\pi}^{+})=0$ or $\mu(E_{\pi'}^{+})=0$, we conclude in the same way that $\mu \le \mu \circ R \circ T$, so that $\mu = \mu \circ R \circ T$. Since both ϑ_{hor}^2 and ϑ_{vert}^2 are compositions of two reflections, the invariance under the translation group $(\vartheta_{x})_{x \in \mathbb{Z}^2}$ follows.

Step 2. We now take advantage of the Burton-Keane uniqueness theorem,⁵ stating that for every periodic μ with finite energy there exists at most one infinite + (respectively, -) cluster, and Zhang's symmetry argument (cf. Ref. 6, Theorem 5.18) deducing from this uniqueness the absence of simultaneous + and -percolation. (In Ref. 5, the uniqueness of the infinite cluster is only stated for translation invariant μ , but the argument works in the same way by applying the ergodic theorem to the subgroup $(\vartheta_x)_{x \in 2\mathbf{Z}^2}$. It is also not shown there that the finite energy property remains valid under ergodic decomposition. Although this follows from Theorem (14.17) of Ref. 7 in the present setting, and a similar argument in general, we do not need this here because our μ is extremal, and therefore $(\vartheta_x)_{x \in 2\mathbb{Z}^2}$ -ergodic by Proposition (14.9) of Ref. 7).

We start noting that, by the flip-reflection symmetry of μ , μ is different from μ^+ and μ^- , so that by Lemma 2.1, the tail triviality of μ , and the Burton–Keane uniqueness theorem there exist both a unique infinite +cluster I^+ and a unique infinite -cluster I^- in the whole plane \mathbb{Z}^2 μ -almost surely. We now choose a square $\Lambda = [-n,n]^2 \subset \mathbb{Z}^2$ so large that $\mu(\Lambda \cap I^+ \neq \emptyset) > 1 - 2^{-12}$. Let $\partial_k \Lambda$ be the intersection of $\partial \Lambda$ with the *k*th quadrant, and let A_k^+ be the increasing event that there exists an infinite +path in Λ^c starting from some site in $\partial_k \Lambda$. Define A_k^- analogously. Since

$$\{\Lambda \cap I^+ \neq \emptyset\} \subset \bigcup_{k=1}^4 A_k^+$$

and μ (as an extremal Gibbs measure) has positive correlations, it follows that

$$\prod_{k=1}^{4} \mu(\Omega \backslash A_{k}^{+}) \leq \mu \left(\bigcap_{k=1}^{4} \Omega \backslash A_{k}^{+} \right) \leq \mu(\Lambda \cap I^{+} = \emptyset) < 2^{-12},$$

whence there exists some $k \in \{1,...,4\}$ such that $\mu(\Omega \setminus A_k^+) < 2^{-3}$. For notational convenience we assume that k = 1. By the above-shown flip-reflection symmetry, we find that

$$\mu(A_1^+ \cap A_2^- \cap A_3^+ \cap A_4^-) > 1 - 4 \cdot 2^{-3} = 1/2,$$

which is impossible because on this intersection the infinite clusters I^+ and I^- cannot be both unique. This contradiction concludes the proof of the lemma.

The butterfly lemma leads immediately to the following result first obtained by Messager and Miracle-Solé⁴ by means of correlation inequalities of symmetry type; the following proof appeared first in Ref. 6.

Corollary 3.2 (Periodic Gibbs measures): Any periodic $\mu \in \mathcal{G}$ is a mixture of μ^+ and μ^- .

Proof: Suppose $\mu \in \mathcal{G}$ is invariant under $(\vartheta_x)_{x \in p\mathbb{Z}^2}$ for some period $p \ge 1$. Conditioning μ on any periodic tail event E we obtain again a periodic Gibbs measure. It is therefore sufficient to show that $\mu(E^+ \cap E^-) = 0$. Indeed, the butterfly lemma then shows that $\mu(E^+) + \mu(E^-) = 1$, and Lemma 2.1 implies that $\mu(\cdot|E^+) = \mu^+$ and $\mu(\cdot|E^-) = \mu^-$ whenever these conditional probabilities are defined. Hence $\mu = \mu(E^+)\mu^+ + \mu(E^-)\mu^-$.

Suppose by contraposition that $\mu(E^+ \cap E^-) > 0$. Since $E^+ \cap E^-$ is invariant and tail measurable, we can in fact assume that $\mu(E^+ \cap E^-) = 1$; otherwise we replace μ by $\mu(\cdot | E^+ \cap E^-)$. By the butterfly lemma, there exists a pair (π, π') of conjugate half-planes, say π_{up} and π_{down} , and a sign, say +, such that both half-planes contain infinite clusters of this sign with positive probability. Since $\mu(E^-) = 1$ by assumption, we can find a large square Δ such that with positive probability Δ meets infinite +clusters in π_{up} and π_{down} and also an infinite -cluster. This -cluster leaves Δ either on the left or on the right between the two infinite +clusters. We can assume that the latter occurs with positive probability. By the finite energy property, it then follows that also $\mu(A_0) > 0$, where for $k \in p\mathbf{Z}$ we write A_k for the event that the point (k, 0) belongs to a two-sided infinite +path with its two halves staying in π_{up} and π_{down} , respectively, and (k+1, 0) belongs to an infinite -cluster.

Let *A* be the event that A_k occurs for infinitely many k < 0 and infinitely many k > 0. The horizontal periodicity and Poincaré's recurrence theorem [cf. Lemma (18.15) of Ref. 7] then show that $\mu(A_0 \setminus A) = 0$, and therefore $\mu(A) > 0$. But on *A* there exist infinitely many –clusters which are separated from each other by the infinitely many "vertical" +paths. This contradicts the Burton–Keane theorem.

The preceding argument actually shows that $\mu(E^{-*}\cap E^{+*})=0$ whenever $\mu \in \mathcal{G}$ is periodic. Since $\mu^+(E^+)=1$ by Lemma 2.1 and tail triviality, this shows that in the +phase the +spins form an infinite sea with only finite islands.

Corollary 3.3 (Plus-sea in the plus-phase): $\mu^+(E^{-*})=0$. Hence, μ^+ -almost surely there exists a unique infinite +cluster I^+ in \mathbb{Z}^2 which surrounds each finite set.

We note that in contrast to Zhang's argument (cf. Theorem 5.18 of Ref. 6) our proof of the preceding corollary does not rely on the reflection invariance of μ^+ but only on its periodicity, and thus can be extended to the setting of Sec. VI.

We conclude this section with the observation that percolation in half-planes is not affected by spatial shifts.

Lemma 3.4 (Shift lemma): Let π and $\tilde{\pi}$ be two half-planes such that $\pi \supset \tilde{\pi}$, i.e., π and $\tilde{\pi}$ are translates of each other. Then $E_{\pi}^+ = E_{\tilde{\pi}}^+$ G-almost surely, and similarly with - instead of +.

Proof: Since trivially $E_{\pi}^+ \supset E_{\tilde{\pi}}^+$, we only need to show that $E_{\pi}^+ \subset E_{\tilde{\pi}}^+ \mathcal{G}$ -almost surely. For definiteness we consider the case when $\pi = \pi_{up} = \{x_2 \ge 0\}$ and $\tilde{\pi} = \{x_2 \ge 1\}$. Take any $\mu \in \operatorname{ex} \mathcal{G}$, and suppose that $\mu(E_{\tilde{\pi}}^+) = 0$. Then for almost all ω and any $n \ge 1$ there exists a smallest $-\operatorname{*semicircuit} \sigma_n(\omega)$ in $\tilde{\pi}$ containing $\Delta_n \cup \sigma_{n-1}(\omega)$ in its interior; here $\Delta_n = [-n,n] \times [1,n]$ and $\sigma_0 = \emptyset$. Let $x_n(\omega) \in l_{\text{left}}$ and $y_n(\omega) \in l_{\text{right}}$ be the two points facing the two end points of $\sigma_n(\omega)$; these are $\mathcal{F}_{\tilde{\pi}}$ -measurable functions of ω , and the random sets $\{x_n, y_n\}$ are pairwise disjoint. Let A_n be the event that the spins at x_n and y_n take value -1.

We claim that A_n occurs for infinitely many n with probability 1. Indeed, fix any $N \ge 1$, $x \in l_{\text{left}}$, $y \in l_{\text{right}}$ and let $B_{N,x,y} = \{x_N = x, y_N = y\} \cap \bigcap_{n \ge N} A_n^c$. Then we can write

$$\mu(A_N \cap B_{N,x,y}) = \mu(\mu_{\{x,y\}}(\omega(x) = \omega(y) = -1) \mathbf{1}_{B_{N,x,y}}) \ge \delta^2 \mu(B_{N,x,y})$$

because $B_{N,x,y}$ only depends on the configuration outside $\{x,y\}$, and the one-point conditional probabilities of μ are bounded from below by $\delta = [1 + e^{8\beta}]^{-1}$. Summing over x, y we obtain $\mu(\bigcap_{n \ge N} A_n^c) \le (1 - \delta^2) \mu(\bigcap_{n \ge N} A_n^c)$, and iteration gives $\mu(\bigcap_{n \ge N} A_n^c) = 0$. Letting $N \to \infty$ we get the claim.

We now can conclude that with probability 1 each box $[-n,n] \times [0,n]$ is surrounded by a -*semicircuit in π_{up} , which means that $\mu(E_{up}^+)=0$. As $\mu(E_{\tilde{\pi}}^+)$ is either 0 or 1, the lemma follows.

IV. UNIQUENESS OF SEMI-INFINITE CLUSTERS

Our next subject is the uniqueness of infinite clusters in half-planes, together with the stronger property that such clusters touch the boundary line infinitely often. This was already a key result of Russo.³

Lemma 4.1 (Line touching lemma): For any half-plane π , there exists *G*-almost surely at most one infinite + (respectively +*) cluster I_{π}^+ (respectively, I_{π}^{+*}) in π . When it exists, this infinite cluster *G*-almost surely intersects the boundary line l of π infinitely often, in the sense that outside any finite Δ one can find an infinite path in this cluster starting from l.

Just as Russo did, we derive this lemma from the absence of percolation for the +phase in the upper half-plane π_{up} with -boundary condition in π_{up}^c (which implies the uniqueness of the semi-infinite Gibbs measure, by the argument of Lemma 2.1). But for the latter we will give here a different argument using stochastic domination by a translation invariant Gibbs measure and Corollary 3.2. To state the result we write \pm for the configuration which is +1 on π_{up} and -1 on π_{up}^c , and consider the semi-infinite limit

$$\mu_{\rm up}^{\pm} = \lim_{\Delta \uparrow \pi_{\rm up}} \mu_{\Delta}^{\pm} \tag{2}$$

which exists by stochastic monotonicity.

Lemma 4.2 (No percolation on a bordered half-plane) $\mu_{up}^{\pm}(E_{up}^{+*})=0$.

Proof: To begin we note that μ_{up}^{\pm} is invariant under horizontal translations and stochastically maximal in the set of all Gibbs measures on π_{up} with -boundary condition in π_{up}^c . This follows just as in the case of the plus-phase μ^+ on the whole lattice. In particular, μ_{up}^{\pm} is trivial on the π_{up} -tail $\mathcal{T}_{up} = \bigcap \{\mathcal{F}_{\pi_{up} \setminus \Lambda} : \Lambda \subset \pi_{up} \text{ finite}\}$. We think of μ_{up}^{\pm} as a probability measure on Ω for which almost all configurations are identically equal to -1 on π_{up}^c .

almost all configurations are identically equal to -1 on $\pi_{\underline{\mu}}^c$. Next we consider the downwards translates $\mu_{n,-}^+ = \mu_{\underline{\mu}p}^- \vartheta_{\text{vert}}^{-n}$, $n \ge 0$. Evidently, $\mu_{n,-}^+$ is obtained by an analogous infinite-volume limit in the half-plane $\{x_2 \ge -n\}$. This shows that $\mu_{n,+}^+ \le \mu_{n+1,-}^+$ by stochastic monotonicity, so that the stochastically increasing limit $\mu_{-}^+ = \lim_{n \to \infty} \mu_{n,-}^+$ exists. Clearly $\mu_{-}^+ \in \mathcal{G}$. Also, μ_{-}^+ inherits the horizontal invariance of the $\mu_{n,-}^+$ and is in addition vertically invariant. Corollary 3.2 therefore implies that $\mu_{-}^+ = a\mu^- + (1-a)\mu^+$ for some coefficient $a \in [0,1]$.

We claim that a>0. For $n\ge 1$ let B_n denote the event that the origin is -*connected to the horizontal line $\{x_2=-n\}$. By the finite energy property and the horizontal ergodicity of $\mu_{n,-}^+$, there exist for $\mu_{n,-}^+$ -almost all ω some random integers $m_{\text{left}}(\omega) < 0 < m_{\text{right}}(\omega)$ such that $\omega \equiv -1$ on

$$\sigma(\omega) = \{ x \in \mathbb{Z}^2 : x_1 \in \{ m_{\text{left}}(\omega), m_{\text{right}}(\omega) \}, -n \leq x_z \leq 0 \}.$$

Together with a segment of the line $\{x_2 = -n-1\}$ on which $\omega = -1\mu_{n,-}^+$ -almost surely, $\sigma(\omega)$ forms a -semicircuit in π_{down} surrounding the origin. An immediate application of the strong Markov property (applied to the largest such σ in a large box) and the point-to-semicircuit lemma thus implies that $\mu_{n,-}^+(B_n) \ge \theta/2$. Therefore, writing $E_{0,m}^{-*}$ for the event that the origin belongs to some -* cluster of size at least m we find $\mu_{n,-}^+(E_{0,m}^{-*}) \ge \theta/2$ when $n \ge m$. Letting first $n \to \infty$ and then $m \to \infty$ we see that $\mu_{-}^+(E^{-*}) \ge \theta/2$. Since $\mu^+(E^{-*}) = 0$ by Corollary 3.3, it follows that $a \ge \theta/2$, and the claim is proved.

To conclude the proof we observe that

$$\mu_{\rm up}^{\pm}(E_{\rm up}^{+*}) \leq \mu_{-}^{+}(E^{+*}) = 1 - a < 1,$$

again by Corollary 3.3. Since μ_{up}^{\pm} is trivial on \mathcal{T}_{up} , the lemma follows.

We are now able to prove Lemma 4.1 along the lines of Russo.³

Proof of Lemma 4.1: For definiteness we assume that $\pi = \pi_{up}$; other half-planes merely correspond to a change of coordinates. We consider only infinite +clusters in π_{up} ; the case of +*clusters is similar. It is also clear that any result proved for the +sign is also valid with the -sign.

Uniqueness: The uniqueness of infinite +clusters in π_{up} is a consequence of the second statement, the line-touching property for infinite -*clusters. Indeed, suppose there exists no infinite -*cluster in π_{up} ; then each finite set in π_{up} is surrounded by a +semicircuit, so that any two infinite +paths are necessarily +connected to each other. In the alternative case when an infinite -*cluster I_{up}^{-*} in π_{up} exists, this I_{up}^{-*} meets l_{left} or l_{right} infinitely often, so that each infinite +cluster must meet the other half-line infinitely often. Hence, two such +clusters must cross each other, and are thus identical.

Line touching: Let $\mu \in ex \mathcal{G}$ and $x \in \pi_{up}$ and consider the event A_x^+ that x belongs to an infinite + cluster in π_{up} which does not touch the horizontal axis l_{hor} . We will show that $\mu(A_x^+)=0$. Once this is established, we can take the union over all x and use the finite energy property to see that for each finite Δ the event "an infinite + cluster in π_{up} is not connected to l_{hor} outside Δ " also has probability zero, which means that almost surely any infinite + cluster in π_{up} must meet l_{hor} infinitely often.

Intuitively, if A_x^+ occurs then the infinite +cluster containing x is separated from l_{hor} by an infinite -*path; but the spins "above" this path feel only the -boundary condition and thus believe to be in the -phase μ^- , so that they will not form an infinite +cluster.

To make this intuition precise we fix some integer $k \ge 1$ and consider the event $A_{x,k}^+$ that x belongs to a +cluster of size at least k which does not meet l_{hor} . Take a box $\Delta \subset \pi_{\text{up}}$ containing

x and so large that there exists no path of length *k* from *x* to Δ^c . For $\omega \in A_{x,k}^+$ we consider the largest set $\Gamma(\omega) \subset \Delta$ containing *x* such that $\omega = -1$ on $\partial \Gamma(\omega) \setminus \partial_{up} \Delta$, where $\partial_{up} \Delta = \partial \Delta \cap \pi_{up}$. We also consider the event $E_{x,k}^+$ that *x* belongs to a +cluster in π_{up} of size at least *k*. Using the fact that $A_{x,k}^+$ is contained in the \mathcal{F}_{Δ} -measurable event { Γ exists} $\cap E_{x,k}^+$, we obtain by the strong Markov property and the stochastic monotonicity of Gibbs distributions that

$$\mu(A_{x,k}^+) \leq \mu(\mu_{\Gamma}(E_{x,k}^+) \leq \mu_{\Delta}^{\pm}(E_{x,k}^+),$$

where the \pm boundary condition is defined as in (2). Now, taking first the limit $\Delta \uparrow \pi_{up}$ as in (2) and then letting $k \to \infty$ we find that $\mu(A_x^+) \leq \mu_{up}^{\pm}(E_{up}^+)$. But the last expression vanishes by Lemma 4.2.

The butterfly lemma and shift lemma together still leave the possibility that all infinite butterflies have the same orientation, either horizontal or vertical. As a consequence of the line touching lemma, we can now show that both orientations must occur.

Lemma 4.3 (Orthogonal butterflies): G-almost surely there exist both a horizontal infinite butterfly in π_{up} and π_{down} as well as a vertical infinite butterfly in π_{left} and π_{right} .

Proof: Suppose there exists some $\mu \in \text{ex } \mathcal{G}$ having almost surely no vertical infinite butterfly. By the first step in the proof of the butterfly lemma, it then follows that $\mu = \mu \circ R_{k,\text{vert}} \circ T$ for all $k \in \mathbb{Z}$, and thus $\mu = \mu \circ \vartheta_{\text{hor}}^{-2}$. By the tail triviality, μ is in fact ergodic under ϑ_{hor}^2 ; cf. Proposition (14.9) of Ref. 7. By the butterfly lemma, horizontal infinite butterflies do exist, say of color +.

We now use an argument similar to that in Corollary 3.2, with the line touching lemma in place of the Burton-Keane theorem. Fix any $n \ge 1$. For $k \in \mathbb{Z}$ let A_k denote the event that all spins along the straight path $p_{k,n} = ((k,l):l = -n,...,n)$ are +1, (k, n) belongs to an infinite +cluster in $\pi_{n,\text{down}} = \{x \in \mathbb{Z}^2: x_2 \ge n\}$, and (k, -n) belongs to an infinite +cluster in $\pi_{n,\text{down}} = \{x \in \mathbb{Z}^2: x_2 \ge n\}$, and (k, -n) belongs to an infinite +cluster in $\pi_{n,\text{down}} = \{x_2 \le -n\}$. Let A be the event that A_k occurs for infinitely many k < 0 and infinitely many k > 0. The finite energy property then shows that $\mu(A_0) > 0$, and the horizontal ergodicity and Poincaré's recurrence theorem (or the ergodic theorem) imply that $\mu(A) = 1$. But the line touching lemma guarantees that the infinitely many doubly infinite ''vertical'' +paths passing through the horizontal axis are connected to each other in $\pi_{n,\text{up}}$ and $\pi_{n,\text{down}}$. As n was arbitrary, it follows that almost surely each finite set is surrounded by a +circuit, and an infinite -cluster cannot exist. In view of Lemma 2.1, this implies that $\mu = \mu^+$. But μ^+ is not invariant under $R_{\text{vert}} \circ T$, in contradiction to what we derived for μ .

The preceding argument can be used to derive the result of Russo³ that μ^+ and μ^- are the only phases which are periodic in one direction. We will not need this intermediate result.

V. NONCOEXISTENCE OF PHASES

In this section we will prove the following proposition.

Proposition 5.1 (Absence of nonperiodic phases): Any Gibbs measure $\mu \in \mathcal{G}$ is invariant under translations, i.e., $\mu = \mu \circ \vartheta_{hor}^{-1}$ and $\mu = \mu \circ \vartheta_{vert}^{-1}$.

Together with Corollary 3.2 this will immediately imply the main theorem that each Gibbs measure is a mixture of the two phases μ^+ and μ^- . Our starting point is the following lemma estimating the probability that a semi-infinite cluster can be pinned at a prescribed point.

Lemma 5.2 (Pinning lemma): Let $\mu \in \mathcal{G}$, and suppose that μ -almost surely there exists an infinite +*cluster I_{up}^{+*} in π_{up} which meets the right semiaxis l_{right} infinitely often. Then for each finite square $\Delta = [-n,n]^2$ and $x \in l_{right}$ we have

 $\mu(x \text{ is } + \text{*connected in } (\Delta \cup l_{\text{left}})^c \text{ to } I_{\text{up}}^{+*}) \geq \theta/4$

provided x lies sufficiently far to the right. The same holds when ''left'' and ''right'' or ''up'' and ''down'' are interchanged.

Proof: By hypothesis, the infinite component of $I_{up}^{+*} \setminus \Delta$ almost surely contains infinitely many points of l_{right} . Thus, if $x \in l_{right}$ is located far enough to the right then, with probability exceed-

ing 1/2, at least one such point can be found left from x, and another such point can be found right from x. This means that x is surrounded by a +*semicircuit σ in π_{up} which belongs to I_{up}^{+*} and satisfies $\Delta \cap \operatorname{Int} \sigma = \emptyset$.

Let Λ be a large square box containing x. If Λ is large enough, a semicircuit σ as above can be found within Λ with probability still larger than 1/2. We then can assume that σ has the largest interior among all such semicircuits in Λ . Using the strong Markov property and the point-to-semicircuit lemma we get the result.

Our main task in the following is to analyze the situation when a half-plane contains both an infinite +cluster and an infinite -cluster. (The line-touching lemma still allows this possibility.) In this situation it is useful to consider contours.

As is usually done in the Ising model, we draw lines of unit length between adjacent spins of opposite sign. We then obtain a system of polygonal curves running through the sites of the dual lattice $\mathbf{Z}^2 + (\frac{1}{2}, \frac{1}{2})$. A *contour* γ in the upper half-plane π_{up} is a part of these polygonal curves which separates a -cluster in π_{up} from a +*cluster in π_{up} . This corresponds to the convention that at crossing points the contours are supposed to bend around the -spins. (The artificial asymmetry between + and - does not matter, and we could clearly make the opposite convention.) On its two sides, γ is accompanied by a +*quasipath f_{γ}^+ and a -quasipath f_{γ}^- which will be called, respectively, the + and -face of γ ; the prefix "quasi" indicates that the faces are not necessarily self-avoiding but may contain loops.

Lemma 5.3 (Semi-infinite contours): G-almost surely on $E_{up}^{+*} \cap E_{up}^{-}$ there exists a unique semi-infinite contour γ_{up} in π_{up} . γ_{up} starts between two points of the horizontal axis l_{hor} and intersects each horizontal line in π_{up} only finitely often.

Proof: Let I_{up}^{+*} be the unique infinite +*cluster in π_{up} , and I_{up}^{-} the unique infinite -cluster in π_{up} . For definiteness we assume that I_{up}^{+*} meets l_{right} infinitely often, and I_{up}^{-} meets l_{left} infinitely often. Let x be the rightmost point of $I_{up}^{-} \cap l_{hor}$ and γ_{up} the contour in π_{up} starting from the line segment separating x and y = x + (1,0). Since I_{up}^{-} contains an infinite -path starting from x which cannot be traversed by γ_{up} , γ_{up} cannot return to l_{hor} on the left-hand side of x. But γ_{up} can also not return to l_{hor} on the right-hand side of y, since otherwise the -face of γ_{up} would establish a -connection in I_{up}^{-} from x to a point of l_{hor} to the right of y, in contradiction to the choice of x. Hence γ_{up} can never end and must therefore be infinite.

Let γ be any infinite contour in π_{up} . Then the infinite $-face f_{\gamma}^{-}$ must belong to I_{up}^{-} , by the uniqueness of the infinite -cluster. This implies that f_{γ}^{-} must lie on the "left-hand side" of γ_{up} . Likewise, the $+*face f_{\gamma}^{+*}$ must belong to the "side on the right" of γ_{up} . Hence $\gamma = \gamma_{up}$, proving the uniqueness of γ_{up} .

Finally, let $l = \{x_2 = n\}$, $n \ge 1$, be a horizontal line in π_{up} and $\pi = \{x_2 \ge n\}$ the half-plane above l. By the shift lemma and the above, π contains a unique semi-infinite contour γ starting from the line segment between two adjacent points u and v of l. u and v belong to the infinite faces of γ and therefore to I_{up}^{+*} and I_{up}^{-} , respectively. By the line touching lemma, this means that u and v are +*connected and -connected, respectively, to the axis l_{hor} . The unique continuation of γ can therefore visit only finitely many sites of π_{up} , and thus must reach l_{hor} after finitely many steps; this continuation is then equal to γ_{up} , by the uniqueness of the latter. This shows that γ_{up} visits the line l only finitely often.

From now on we consider a fixed external Gibbs measure $\mu \in \operatorname{ex} \mathcal{G}$. We want to prove that μ is horizontally invariant. (The proof of vertical invariance is similar.) To this end we consider its horizontal translate $\hat{\mu} = \mu \circ \vartheta_{hor}^{-1}$, as well as the product measure $\hat{\nu} = \mu \otimes \hat{\mu}$ on $\Omega \times \Omega$. It is convenient to think of the latter as a duplicated system consisting of two independent layers. The following lemma is a slight variation of a result of Aizenman¹ in his proof of the main theorem; our proof differs in part.

Lemma 5.4 (Fluctuations of the semi-infinite contour): Suppose π_{up} contains a semi-infinite contour $\gamma_{up} \mu$ -almost surely. Then for $\hat{\nu}$ -almost all $(\omega, \hat{\omega}) \in \Omega^2$, $\gamma_{up}(\omega)$ and $\gamma_{up}(\hat{\omega})$ intersect each other infinitely often.

Proof: By tail triviality, we can assume that γ_{up} has its +face on the left-hand side almost surely; the alternative case is analogous. For any $n \ge 1$ we let

$$a_n = \max\{k \in \mathbf{Z}: (k,n) \in I_{\pi_n, m}^{+*}\}$$

be the abscissa of the point at which γ_{up} enters definitely into the half-plane $\pi_{n,up} = \vartheta_{vert}^n \pi_{up}$ above the height *n*. Consider the product measure $\nu = \mu \otimes \mu$ and the event

 $F = \{(\omega, \omega') \in \Omega^2: \gamma_{up}(\omega) \text{ and } \gamma_{up}(\vartheta_{hor}\omega') \text{ meet each other only finitely often}\}.$

We need to show that $\nu(F) = 0$.

Suppose that *F* occurs. Then $\gamma_{up}(\omega)$ lies strictly on one side of $\gamma_{up}(\vartheta_{hor}\omega')$ above some random level *n*. Hence we have either $a_n(\omega) > a_n(\vartheta_{hor}\omega')$ eventually, or $a_n(\omega) < a_n(\vartheta_{hor}\omega')$ eventually. Using the abbreviation $d_n(\omega, \omega') = a_n(\omega) - a_n(\omega') = a_n(\omega) - a_n(\vartheta_{hor}\omega') + 1$, we thus see that

$$F \subset A \cup B \equiv \{d_n \ge 0 \text{ eventually}\} \cup \{d_n \le 0 \text{ eventually}\}.$$

Suppose now that $\nu(F) > 0$. Then, by symmetry, $\nu(A) = \nu(B) > 0$. By the tail triviality of μ , it follows that $\nu(A) = \nu(B) = 1$. This is because A, B are measurable with respect to the "product-tail" $\mathcal{T}^{(2)} = \bigcap \{ \mathcal{F}_{\Lambda^c} \otimes \mathcal{F}_{\Lambda^c} : \Lambda \subset \mathbb{Z}^2 \text{ finite} \}$ in Ω^2 , which is trivial by Fubini's theorem. (One should not be mistaken to believe that A was measurable with respect to the smaller "tail-product" $\mathcal{T} \otimes \mathcal{T}$. It is only the case that the ω -section A_{ω} of A belongs to \mathcal{T} for any ω , and the function $\omega \rightarrow \mu(A_{\omega})$ is \mathcal{T} -measurable.) We thus conclude that $\nu(A \cap B) = 1$, meaning that $d_n = 0$ eventually, almost surely. The lemma will therefore be proved once we have shown that this is impossible.

To this end we claim first that $\nu(d_n \ge 1) \ge \delta \nu(d_{n+1}=0)$ for all *n* and some constant $\delta \ge 0$. To see this let $A_{k,n} = \{(\omega, \omega'): a_{n+1}(\omega) = a_{n+1}(\omega') = k\}$, $\Delta_{k,n}$ the two-point set consisting of the points (k,n) and (k+1,n), and $B_{k,n}$ the event that $\omega = (+1,+1)$ on $\Delta_{k,n}$ and $\omega' = (+1,-1)$ on $\Delta_{k,n}$; see the following diagram:

First layer			Second layer	
n+1	+	_	+	_
n	+	+	+	-
	k	k+1	\boldsymbol{k}	k+1

We then have

$$\nu(B_{k,n}|\mathcal{F}_{\Delta_{k,n}^c}\otimes\mathcal{F}_{\Delta_{k,n}^c})(\omega,\omega')=\mu_{\Delta_{k,n}}^{\omega}\otimes\mu_{\Delta_{k,n}}^{\omega'}(B_{k,n})\ge[1+e^{8\beta}]^{-4}\equiv\delta$$

and thus

$$\nu(\{d_n \ge 1\} \cap A_{k,n}) \ge \nu(\nu(B_{k,n} | \mathcal{F}_{\Delta_{k,n}^c} \otimes \mathcal{F}_{\Delta_{k,n}^c}) \mathbf{1}_{A_{k,n}}) \ge \delta \nu(A_{k,n})$$

because $A_{k,n}$ is an event in $\Delta_{k,n}^c$. Summing over k we get the claim.

Now, if $d_n = 0$ eventually almost surely then

$$\inf_{n \to \infty} \lim \nu(d_n \ge 1) \ge \delta \liminf_{n \to \infty} \nu(d_{n+1} = 0) = \delta,$$

so that with positive probability we have simultaneously $d_n \ge 1$ infinitely often and $d_n = 0$ eventually. Since this is impossible, we conclude that $\nu(F) = 0$.

The following percolation result for the duplicated system with distribution $\hat{\nu}$ was already a cornerstone of Aizenman's argument.¹ We prove it here differently, avoiding his use of the fact that the limiting Gibbs measure for the \pm boundary condition is translation invariant. We will say that a path in \mathbb{Z}^2 is a \leq path for a pair $(\omega, \hat{\omega}) \in \Omega^2$ if $\omega(x) \leq \hat{\omega}(x)$ for all its sites *x*. In the same way we define \leq *paths, and we can speak of \leq *circuits and \leq *clusters.

Lemma 5.5 (No (+, -)percolation in the duplicated system): $\hat{\nu}$ -almost surely each finite square $\Delta = [-n,n]^2$ is surrounded by $a \leq *$ circuit in \mathbb{Z}^2 .

Proof: Consider any two points $x \in l_{\text{left}}$ and $y \in l_{\text{right}}$. We claim that with $\hat{\nu}$ -probability at least $(\theta/4)^2$ there exists a $\leq *$ path from x to y "above" Δ , provided x and y are located sufficiently far to the left and to the right, respectively. We distinguish three cases.

Case 1: $\mu(E_{up}^+)=0$. By Lemma 4.3, π_{up} then almost surely contains an infinite-cluster I_{up}^- , and each finite subset of π_{up} is surrounded by a -*semicircuit in π_{up} . In other words, an infinite -*cluster I_{up}^{-*} in π_{up} exists and touches both l_{left} and l_{right} infinitely often. By the pinning lemma and the positive correlations of μ , with μ -probability at least $(\theta/4)^2$ both x and y are -*connected to I_{up}^{-*} outside Δ , and therefore also -*connected to each other by a -*path p above Δ . However, this -*path p on the first layer is certainly also a \leq *path for the duplicated system, and the claim follows.

Case 2: $\mu(E_{up})=0$. In this case we also have $\hat{\mu}(E_{up})=0$. Interchanging + and - and replacing μ by $\hat{\mu}$ in Case 1, we find that with $\hat{\mu}$ -probability at least $(\theta/4)^2$, there exists a +* path \hat{p} in the second layer above Δ from x to y. Since \hat{p} is again a $\leq *$ path for the duplicated system, the claim follows as in the first case.

Case 3: $\mu(E_{up}^+) = \mu(E_{up}^-) = 1$. Then μ -almost surely there exists a unique semi-infinite contour γ_{up} , and by tail triviality we can assume (for definiteness) that γ_{up} has its +face on the left-hand side μ -almost surely, and thus also $\hat{\mu}$ -almost surely. By the pinning lemma and the independence of the two layers, the following event has $\hat{\nu}$ -probability of at least $(\theta/4)^2$:

- (1) In the first layer, y is -*connected off Δ to $I_{up}^{-}(\omega)$, and thus to the -face $f_{up}^{-}(\omega)$ of $\gamma_{up}(\omega)$; that is, there exists an infinite -*path $p_{y}^{-}(\omega)$ from y outside Δ eventually running along $\gamma_{up}(\omega)$.
- (2) In the second layer, x is +*connected off Δ to $I_{up}^{+*}(\hat{\omega})$, and thus to the +face $f_{up}^{+}(\hat{\omega})$ of $\gamma_{up}(\hat{\omega})$; that is, there exists an infinite +*path $p_x^{+}(\hat{\omega})$ from x outside Δ eventually running along $\gamma_{up}(\hat{\omega})$.

Since $\gamma_{up}(\omega)$ and $\gamma_{up}(\hat{\omega})$ intersect each other infinitely often by Lemma 5.4, the union of $p_y^-(\omega)$ and $p_x^+(\hat{\omega})$ contains a *path from *x* to *y* which by construction is a \leq *path for the duplicated system. This proves the claim in the final case.

To conclude the proof of the lemma, we let $A_{x,y}$ denote the event that there exists a $\leq *$ path from x to y above Δ , and $B_{x,y}$ the event that such a path exists below Δ . The indicator functions of these events can be written as increasing functions f and g respectively of the difference configuration $\hat{\omega} - \omega$. Using the positive correlations of $\hat{\omega}$ and μ we thus obtain

$$\hat{\nu}(A_{x,y} \cap B_{x,y}) = \int \mu(d\omega) \int \hat{\mu}(d\hat{\omega}) f(\hat{\omega} - \omega) g(\hat{\omega} - \omega)$$
$$\geq \int \mu(d\omega) \hat{\mu}(f(\cdot - \omega)) \hat{\mu}(g(\cdot - \omega))$$
$$\geq \hat{\nu}(A_{x,y}) \hat{\nu}(B_{x,y}) \geq (\theta/4)^4.$$

The last inequality follows from the claim and its analog for the lower half-plane. However, if $A_{x,y} \cap B_{x,y}$ occurs then Δ is surrounded by a \leq *circuit for the duplicated system. Letting $\Delta \uparrow \mathbb{Z}^2$ we see that with probability at least $(\theta/4)^4$ each finite set is surrounded by a \leq *circuit. Since this event is measurable with respect to the product-tail $\mathcal{T}^{(2)}$ on which $\hat{\nu}$ is trivial, the lemma follows.

It is now easy to complete the proof of Proposition 5.1 as in Ref. 1.

Proof of Proposition 5.1: Consider any square $\Delta = [-n,n]^2$, and let $\epsilon > 0$. By Lemma 5.5, Δ is $\hat{\nu}$ -almost surely surrounded by a \leq *circuit, and with probability at least $1 - \epsilon$ such a \leq *circuit can be found in a sufficiently large square Λ . Let Γ be the interior of the largest such \leq *circuit;

if no such \leq *circuit exists let $\Gamma = \emptyset$. Then we find for any increasing \mathcal{F}_{Δ} -measurable function $0 \leq f \leq 1$, using the strong Markov property of $\hat{\nu}$ and the fact that $\mu_{\Gamma}^{\omega} \leq \mu_{\Gamma}^{\hat{\omega}}$ when $\Gamma(\omega, \omega') \neq \emptyset$,

$$\begin{split} \iota(f) &= \hat{\nu}(f \otimes 1) \\ &\leq \int_{\{\Gamma \neq \emptyset\}} d\hat{\nu}(\omega, \omega') \mu^{\omega}_{\Gamma(\omega, \omega')}(f) + \epsilon \\ &\leq \int d\hat{\nu}(\omega, \omega') \mu^{\omega'}_{\Gamma(\omega, \omega')}(f) + \epsilon \\ &= \hat{\nu}(1 \otimes f) + \epsilon \\ &= \hat{\mu}(f) + \epsilon. \end{split}$$

Letting $\epsilon \to 0$ and $\Delta \uparrow \mathbb{Z}^2$ we find that $\mu \leq \hat{\mu}$. Interchanging μ and $\hat{\mu}$ (i.e., the roles of the layers) we get the reverse relation. Hence $\mu = \hat{\mu}$, so that μ is horizontally invariant. The vertical invariance follows similarly by an interchange of coordinates.

VI. EXTENSIONS

Which properties of the square lattice \mathbf{Z}^2 entered into the preceding arguments? The only essential feature was its invariance under the reflections in all horizontal and vertical lines with integer coordinates. We claim that the theorem remains true for the Ising model on any connected graph \mathcal{L} with these properties. (The Ising model on the triangular and hexagonal lattices has already been treated in Ref. 10.)

To be more precise, let $\mathcal{R} = \{R_{k,hor}, R_{k,vert} : k \in \mathbb{Z}\}$ denote the set of all reflections of the Euclidean plane \mathbb{R}^2 in horizontal or vertical lines with integer coordinates, and suppose \mathcal{L} is a locally finite subset of \mathbb{R}^2 which (after suitable scaling and rotation) is *R*-invariant for all $R \in \mathcal{R}$. Such an \mathcal{L} is uniquely determined by its finite intersection with the unit cube $[0,1]^2$, and it is periodic with period 2. Suppose further that \mathcal{L} is equipped with a symmetric neighbor relation "~" satisfying

(L1) each $x \in \mathcal{L}$ has only finitely many "neighbors" $y \in \mathcal{L}$ satisfying $x \sim y$;

(L2) $x \sim y$ if and only if $Rx \sim Ry$ for all $R \in \mathcal{R}$;

(L3) (\mathcal{L},\sim) is a connected graph.

If $x \sim y$ we say that x and y are connected by an edge, which is visualized by the straight line segment between x and y. The preceding assumptions simply mean that (\mathcal{L}, \sim) is a locally finite connected graph admitting the reflections $R \in \mathcal{R}$, and thereby the translations $\vartheta_x, x \in 2\mathbb{Z}^2$, as graph automorphisms. The fundamental further assumption is

(L4) (\mathcal{L},\sim) is planar, i.e., the edges in \mathbb{R}^2 between different pairs of neighboring points have only end points in common.

The complement (in \mathbf{R}^2) of the union of all edges then splits into connected components called the faces of (\mathcal{L}, \sim) .

As will be explained in more detail in the Appendix, the properties (L1) to (L4) are sufficient for all geometric arguments above. Some particular examples are as follows.

(1) The *triangular lattice* **T**. This is the \mathcal{R} -invariant lattice satisfying $\mathbf{T} \cap [0,1]^2 = \{(1,0),(0,1)\}$ and $(-1,0) \sim (1,0) \sim (0,1) \sim (2,1)$; the remaining edges result from (L2).

(2) The hexagonal or *honeycomb lattice* **H**. Here, for example, $\mathbf{H} \cap [0,1]^2 = \{(\frac{1}{3},1), (\frac{2}{3},0)\}$ and $(-\frac{1}{3},1) \sim (\frac{1}{3},1) \sim (\frac{2}{3},0) \sim (\frac{4}{3},0);$ all other edges are again determined by (L2).

(3) The *diced lattice*. This is obtained from the honeycomb lattice by placing points in the centers of the hexagonal faces and connecting them to the three points in the west, northeast and southeast of these faces; to obtain reflection symmetry an additional shift by $(-\frac{1}{3},0)$ is necessary. See p. 16 of Ref. 11 for more details.

(4) The covering lattice of the honeycomb lattice, the Kagomé lattice, cf. p. 37 of Ref. 11.

As for the interaction, it is neither necessary that all adjacent spins interact in the same way, nor that the interaction is invariant under the spin flip. Except for attractivity, we need only the invariance under *simultaneous* flip-reflections (which in particular implies periodicity with period 2). As a result, we can consider any system of spins $\omega(x) = \pm 1$ with formal Hamiltonian of the form

$$H(\omega) = \sum_{x \sim y} U_{x,y}(\omega(x), \omega(y)) + \sum_{x \in \mathcal{L}} V_x(\omega(x)),$$
(3)

where for all $a, b \in \{-1, 1\}$ we have $U_{x,y}(a, b) = U_{y,x}(b, a)$ and

(H1) $U_{x,y}(1,\cdot) - U_{x,y}(-1,\cdot)$ is decreasing on $\{-1, 1\}$;

(H2) $U_{x,y}(a,b) = U_{Rx,Ry}(-a,-b)$ and $V_x(a) = V_{Rx}(-a)$ for all $R \in \mathcal{R}$.

Assumption (H1) implies that the FKG inequality is applicable, and (H2) expresses the invariance under simultaneous spatial reflection and spin flip. We thus obtain the following general result.

Theorem 6.1: Consider a planar graph (\mathcal{L}, \sim) as above and an interaction of the form (3) satisfying (H1) and (H2). Then there exist no more than two extremal Gibbs measures.

The standard case, of course, is the ferromagnetic Ising model without external field; this corresponds to the choice $U_{x,y}(a,b) = -\beta ab$ and $V_x \equiv 0$.

But there is also another case of particular interest. Consider $\mathcal{L} = \mathbb{Z}^2 + (\frac{1}{2}, \frac{1}{2})$, the shifted square lattice with its usual graph structure. \mathcal{L} is bipartite, in the sense that \mathcal{L} splits into two disjoint sublattices, \mathcal{L}_{even} and \mathcal{L}_{odd} , such that all edges run from one sublattice to the other. If we set $U_{x,v}(a,b) = -\beta ab$ and define a staggered external field

$$V_x(a) = \begin{cases} -ha & \text{if } x \in \mathcal{L}_{\text{even}} \\ ha & \text{if } x \in \mathcal{L}_{\text{odd}} \end{cases}$$

with $h \in \mathbf{R}$ then the conditions (H1) and (H2) hold; here we take advantage of the fact that the reflections $R \in \mathcal{R}$ map \mathcal{L}_{even} into \mathcal{L}_{odd} and vice versa. But it is well-known that this model is isomorphic to the *anti*ferromagnetic Ising model on \mathbf{Z}^2 with homogeneous external field *h*; the isomorphism consists in flipping all spins in \mathcal{L}_{odd} . This gives us the following result.

Corollary 6.2: For the Ising antiferromagnet on \mathbb{Z}^2 for any inverse temperature and arbitrary external field there exist at most two extremal Gibbs measures.

This corollary does not extend to nonbipartite lattice such as the triangular lattice. In fact, for the Ising antiferromagnet on \mathbf{T} one expects the existence of three different phases for suitable h.

Another repulsive model to which our arguments can be applied is the hard-core lattice gas on \mathbb{Z}^2 , which is also known as the hard square model. In this model, the values -1 and 1 are interpreted as the absence and presence of a particle, respectively, and no particles are allowed to sit on adjacent sites. Its Hamiltonian is of the form (3) with

$$U_{x,y}(a,b) = \begin{cases} \infty & \text{if } a = b = 1 \\ 0 & \text{otherwise,} \end{cases}$$
$$V_x(a) = \begin{cases} -\log \lambda & \text{if } a = 1 \\ 0 & \text{otherwise.} \end{cases}$$

The parameter $\lambda > 0$ is called the activity. Interchanging the values ± 1 on \mathcal{L}_{odd} we obtain an isomorphic attractive model to which our techniques can be applied, although the interaction takes the value $+\infty$ so that the finite energy condition does not hold as it stands. However, there are still enough admissible configurations to satisfy all needs of the Burton–Keane theorem and our other applications of the finite energy property; more details will be provided in the Appendix. We therefore can state the following theorem.

Theorem 6.3: For the hard-core lattice gas on \mathbb{Z}^2 at any activity $\lambda > 0$ there exist at most two extremal Gibbs measures.

APPENDIX

Here we explain in more detail how our arguments can be extended to obtain Theorems 6.1 and 6.3.

Comments on the proof of Theorem 6.1: (1) *Connections and contours. A basic consequence of the planarity assumption (L4) is that (\mathcal{L}, \sim) admits a conjugate matching graph (\mathcal{L}, \sim) . As indicated by the notation, this conjugate graph has the same set of vertices, but the relation $x^* \sim y$ holds if either $x \sim y$ or x and y are distinct points (on the border) of the same face of (\mathcal{L}, \sim) . (Note that this matching dual is in general not planar. An interesting exception is the triangular lattice **T**, which is self-matching.) The edges of (\mathcal{L}, \sim) are then used to define the concept of *connectedness. The construction implies that every path in (\mathcal{L}, \sim) is also a *path (i.e., a path in (\mathcal{L}, \sim) , and that the outer boundary of any cluster is a *path, and vice versa. (The latter property holds for arbitrary matching pairs of graphs as defined in Kesten,¹¹ for example. However, we also used repeatedly the former property which does not extend to general matching pairs. In particular, this means that our results do not apply to the Ising model on the matching conjugate of \mathbb{Z}^2 having nearest-neighbor interactions *and* diagonal interactions.)

Another consequence of planarity is that we can draw contours separating clusters from *clusters. Such contours can either be visualized by broken lines passing through the edges of (\mathcal{L}, \sim) , or simply as a pair consisting of a quasipath and an adjacent *quasipath, namely the two faces of the contour.

(2) Half-planes and boundary lines. A half-plane π in \mathcal{L} is still defined as the intersection of \mathcal{L} with a set of the form $\{x \in \mathbf{R}^2 : x_i \ge k\}, k \in \mathbf{Z}, i \in \{1,2\}$, or with \le instead of \ge . However, the "boundary line" l is now in general not a straight line but rather the set $l = \{x \in \pi : x \sim y \text{ for some } y \notin \pi\} = \partial(\pi^c)$. In particular, l is not necessarily a line of fixed points for the reflection $R \in \mathcal{R}$ mapping π onto its conjugate half plane π' . Rather, for each $x \in l$ we have either Rx = x or $Rx \sim x$. For example, for $\mathcal{L} = \mathbf{T}$, the triangular lattice, π_{up} and π_{down} have a common straight boundary line, but the boundaries of π_{right} and π_{left} are not straight; besides a common part on the vertical axis they also contain the adjacent points (1,k) and (-1,k), $k \in 2\mathbf{Z}$, respectively. For the honeycomb lattice \mathbf{H} , π_{up} and π_{down} have again a common straight boundary line, but π_{right} have no common points.

Nevertheless, it is easy to see that Lemma 2.3 (and thus also Lemma 5.2) are still valid, and these are the only results in which fixed points of reflections show up. In all other places one has only to observe that the axes l_{hor} and l_{vert} get a different meaning according to which half-space is considered; so one has to distinguish between an "upper" horizontal axis $l_{\text{hor,up}}$ (being the boundary "line" of π_{up}) and a "lower" horizontal axis $l_{\text{hor,down}}$, and similarly between $l_{\text{vert,left}}$ and $l_{\text{vert,right}}$.

(3) Construction of connections and paths. At various places we needed to establish prescribed connections or to construct specific paths. For example, the key idea of Lemma 3.4 was to extend -*semicircuits in $\tilde{\pi}$ to the boundary line of π . In the present setup, this will in general require a finite -path rather than a single -spin, so that one has to adapt the definition of A_n accordingly. In view of (L3) this is obviously possible, and one will only end up with a higher power of δ . Similarly, the -semicircuit σ in the proof of Lemma 4.2 has in general to be redefined using the geometry of \mathcal{L} , and the same is the case for the points (k, 0) in the definition of A_k in the proof of Corollary 3.2, the paths $p_{k,n}$ in the proof of Lemma 4.3, and the sets $\Delta_{k,n}$ in the proof of Lemma 5.4; see also comment (5).

(4) *Flip-reflection invariance*. In the standard Ising model on \mathbb{Z}^2 it is true that the phases μ^+ and μ^- are invariant under all $R \in \mathcal{R}$ and related to each other by the spin flip *T*. However, we did not make use of this fact, cf. the comments after Corollary 3.3. We only needed that $\mu^+ = \mu^- \circ T$ for all $R \in \mathcal{R}$ (implying that μ^+ and μ^- are periodic, and that any flip-reflection invariant μ is different from these phases; the latter was used in Lemmas 3.1 and 4.3). This, however, already holds whenever the interaction is only invariant under simultaneous flip-reflections, as stated in assumption (H2). This property is also sufficient for flip-reflection domination and the

H.-O. Georgii and Y. Higuchi

point-to-semicircuit lemma, as their proofs only use the composed mapping $R \circ T$ for $R \in \mathcal{R}$.

(5) *Translations*. Since the lattice and the interaction are in general only preserved by the translation subgroup ϑ_x , $x \in 2\mathbb{Z}^2$, we have to confine ourselves to this class of translations. We did this already in the proof of the butterfly lemma and its Corollary 3.2, and we can obviously do so in the proof of Lemma 4.2. The only statements needing discussion are Proposition 5.1 and Lemma 5.4. The former now only asserts that each Gibbs measure is periodic with period 2. Accordingly, in Lemma 5.4 and below we have to replace ϑ_{hor} by ϑ_{hor}^2 . In addition, the minimal distance between distinct lattice points can be less than 1, and the origin does not necessarily belong to \mathcal{L} . So, a_n has to be defined as the abscissa of the rightmost point in the boundary line of $\pi_{n,up}$ which belongs to $I_{\pi_{n,up}}^{+*}$, and $d_n(\omega, \omega') = a_n(\omega) - a_n(\omega') = a_n(\omega) - \alpha_n(\vartheta_{hor}^2 \omega') + 2$. In general, we then have only the inclusion

$$F \subset \{d_n > -2 \text{ eventually}\} \cup \{d_n < 2 \text{ eventually}\},$$

and we need to derive a contradiction from the assumption that $|d_n| < 2$ eventually almost surely. This means that we have to prescribe the configurations for the two layers on larger sets than $\Delta_{k,n}$ (depending on *n* and both $a_n(\omega)$ and $a_n(\omega')$) to obtain the inequality $\nu(|d_n| \ge 2) \ge \delta \nu(|d_{n+1}| < 2)$ for some $\delta > 0$. While this is tedious to write down in full generality, it should be clear how it can be done.

Comments on the proof of Theorem 6.3: Just as in the case of the Ising antiferromagnet, we replace \mathbb{Z}^2 by its translate $\mathcal{L} = \mathbb{Z}^2 + (\frac{1}{2}, \frac{1}{2})$. So we make sure that all reflections $R \in \mathcal{R}$ map \mathcal{L}_{even} into \mathcal{L}_{odd} and vice versa. Nevertheless, below it will be convenient to ignore the shift by $(\frac{1}{2}, \frac{1}{2})$ and to characterize the lattice points by integer coordinates. Performing a spin flip on \mathcal{L}_{odd} we obtain an isomorphic model which is defined by setting

$$U_{x,y}(a,b) = \begin{cases} \infty & \text{if } a = \epsilon(x), b = \epsilon(y) \\ 0 & \text{otherwise,} \end{cases}$$
$$V_x(a) = \begin{cases} -\log \lambda & \text{if } a = \epsilon(x) \\ 0 & \text{otherwise,} \end{cases}$$

where $\epsilon(x) = 1$ if $x \in \mathcal{L}_{even}$, and $\epsilon(x) = -1$ otherwise. This model satisfies both (H1) and (H2). However, the finite energy condition is violated because $U_{x,y}$ takes the value $+\infty$. Let us see how this obstacle can be overcome. The basic observation is that the "vacuum configuration" $-\epsilon$ can occur in any finite region with positive probability.

(1) In the proof of the Burton-Keane theorem, the finite energy property is used to connect different +clusters with positive probability. This is still possible because for any box Δ , any $x \in \Delta$, any finite number of points $x_1, \ldots, x_k \in \partial \Delta$, and any ω with $\omega(x_1) = \cdots = \omega(x_k) = +1$ we have

$$\mu_{\Delta}^{\omega}(x \text{ is } + \text{connected to } x_1, \dots, x_k) > 0.$$

(2) A different use of the finite energy property is made in the proofs of Corollary 3.2 and Lemma 4.3: the events A_k there involve the existence of both + and -paths. To adapt the proof of Corollary 3.2 to the present case we redefine A_0 as the event that a prescribed point $x \in \mathcal{L}_{odd}$ belongs to a two-sided infinite +path with its two halves staying in π_{up} and π_{down} , respectively, and a neighbor point $y \in \mathcal{L}_{even}$ belongs to an infinite -cluster; for $k \in 2\mathbb{Z}$ we set $A_k = \vartheta_{hor}^{-k}A_0$. A +spin at x then does not interfere with a -spin at y. Therefore, if Δ is a sufficiently large box and $u_1, u_2, u_3 \in \partial \Delta$ are three points belonging, respectively, to infinite +, +, -clusters meeting Δ , we can find paths p_1, p_2 in Δ from x to u_1 resp. u_2 and a path p_3 from y to u_3 such that y is the only site of p_3 which is adjacent to $p_1 \cup p_2$. The configuration in Δ which is equal to +1 on $p_1 \cup p_2$, -1 on p_3 , and $-\epsilon$ otherwise then has positive conditional probability given the configuration in Δ^c . This shows that $\mu(A_0) > 0$. The proof of Lemma 4.3 can be adapted in a similar manner. (3) In Lemma 4.2 we used the finite energy property to make sure that $\mu_{n,-}^+(\omega \equiv -1 \text{ on } p) > 0$, where $p = \{0\} \times \{-n+1,...,0\}$. To obtain the same result here we simply set $\Delta = p \cup \partial p \setminus \{x_2 = -n\}$ and observe that

$$\mu_{\Lambda}^{\omega}(\omega \equiv -1 \text{ on } p, \omega \equiv -\epsilon \text{ on } \Delta p) > 0$$

whenever $\omega(0, -n) = -1$.

(4) Uniform lower bounds for conditional probabilities were used twice, in the proofs of the shift lemma and the contour fluctuation lemma. In the proof of Lemma 3.4, it is sufficient to replace the set $\{x,y\}$ by $\Delta(x) \cup \Delta(y)$, where $\Delta(x) = \{k-1,k,k+1\} \times \{n-1,n\}$ when x = (k,n). This is because for $\omega(k,n+1) = -1$ we have the estimate

$$\mu^{\omega}_{\Delta(x)}(\omega(x) = -1, \quad \omega \equiv -\epsilon \text{ on } \Delta(x) \setminus \{x\}) \ge \delta \equiv \frac{\lambda \wedge 1}{(1+\lambda)^6}.$$

More care is needed in the proof of Lemma 5.4 where we used a uniform estimate for the conditional probability of $B_{k,n}$ given $A_{k,n}$. First, according to comment (5) on the proof of Theorem 6.1 we have to specify the abscissas $a_n(\omega), a_n(\omega')$ by two parameters $k, k' \in \mathbb{Z}$ with $|k-k'| \leq 1$. Note, however, that the point $(a_n(\omega), n)$ necessarily belongs to \mathcal{L}_{even} because otherwise $\omega = \epsilon$ at the adjacent points $(a_n(\omega), n+1)$ and $(a_n(\omega)+1, n+1)$; but this is excluded by the hard-core interaction. Therefore we have in fact k=k', and we can consider the events $A_{k,n}$ as before. Next we redefine $\Delta_{k,n}$ as the set $\{k-1, \dots, k+2\} \times \{n-1, n\}$, and $B_{k,n}$ as the event that $\omega(k,n) = \omega(k+1,n) = 1$ (as before), $\omega'(k,n) = \omega'(k+1,n) = -1$ (in variation of the former definition), and anything else occurs at the remaining sites of $\Delta_{k,n}$ we find

$$\mu^{\omega}_{\Delta_{k,n}} \otimes \mu^{\mu'}_{\Delta_{k,n}}(B_{k,n}) \geq \frac{\lambda}{(1+\lambda)^8} \equiv \delta$$

as above. We can thus argue as before.

- ²Y. Higuchi, "On the absence of non-transiation invariant Gibbs states for the two-dimensional Ising model," in *Random Fields, Esztergom (Hungary)*, edited by J. Fritz, J. L. Lebowitz, and D. Szász (North-Holland, Amsterdam, 1979), Vol. I, pp. 517–534.
- ³L. Russo, "The infinite cluster method in the two-dimensional Ising model," Commun. Math. Phys. **67**, 251–266 (1979).
- ⁴A. Messager and S. Miracle-Solé, "Equilibrium states of the two-dimensional Ising model in the two-phase region," Commun. Math. Phys. 40, 187–196 (1975).
- ⁵R. Burton and M. Keane, "Density and uniqueness in percolation," Commun. Math. Phys. 121, 501–505 (1989).
- ⁶H.-O. Georgii, O. Häggström, and C. Maes, "The random geometry of equilibrium phases," in *Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, New York, 2000).
- ⁷H.-O. Georgii, Gibbs Measures and Phase Transitions (de Gruyter, Berlin, 1988).
- ⁸A. Coniglio, C. R. Nappi, F. Peruggi, and L. Russo, "Percolation and phase transitions in the Ising model," Commun. Math. Phys. **51**, 315–323 (1976).
- ⁹A. Gandolfi, M. Keane, and L. Russo, "On the uniqueness of the infinite occupied cluster in dependent two-dimensional site percolation," Ann. Prob. **16**, 1147–1157 (1988).
- ¹⁰H. Fukuda, "The two-dimensional Ising model and Gibbs states, an approach with percolation method," Master thesis, Osaka University (in Japanese).
- ¹¹H. Kesten, Percolation Theory for Mathematicians (Birkhauser, Boston, 1982).

¹M. Aizenman, "Translation invariance and instability of phase coexistence in the two-dimensional Ising system," Commun. Math. Phys. **73**, 83–94 (1980).