

ADVANCED ANALYSIS – WiSe 2019/20

Exercise sheet 1

22.10.2019

Emanuela Giacomelli – emanuela.giacomelli@math.lmu.de

Exercise 1. [10 points]

Let f_n be an increasing sequence of summable functions on the measure space (Ω, Σ, μ) and let $f(x)$ be defined as

$$f(x) := \lim_{n \rightarrow +\infty} f_n(x).$$

Prove that f is measurable and

$$\lim_{n \rightarrow +\infty} \int_{\Omega} f_n(x) d\mu(x) = \int_{\Omega} f(x) d\mu(x).$$

Exercise 2. [10 points]

Find a simple condition on $f_n(x)$ so that

$$\sum_{n=0}^{\infty} \int_{\Omega} f_n(x) \mu(dx) = \int_{\Omega} \left\{ \sum_{n=0}^{\infty} f_n(x) \right\} \mu(dx).$$

Exercise 3. [10 points]

Let $1 \leq p \leq 2$ and let $0 < b < a$. Prove that

$$(a+b)^p + (a-b)^p \geq 2a^p + p(p-1)a^{p-2}b^2.$$

Exercise 4. [10 points]

Recall that if f and g are functions in $L^p(\Omega)$ with $1 \leq p \leq 2$, then

$$\|f+g\|_p^p + \|f-g\|_p^p \geq (\|f\|_p + \|g\|_p)^p + \|\|f\|_p - \|g\|_p\|^p, \quad (1)$$

$$(\|f+g\|_p + \|f-g\|_p)^p + \|\|f+g\|_p - \|f-g\|_p\|^p \leq 2^p(\|f\|_p^p + \|g\|_p^p). \quad (2)$$

Assume that f and g lie on the unit sphere in L^p , i.e., $\|f\|_p = \|g\|_p = 1$. Assume also that $\|f-g\|_p$ is small. Draw a picture of the situation. Then, using exercise 3, explain why (2) shows that the unit sphere is *uniformly convex*. Explain also why (1) shows that the unit sphere is *uniformly smooth*, i.e., it has no corners.