

# Groups quasi-isometric to symmetric spaces

Bruce Kleiner\* and Bernhard Leeb<sup>‡</sup>

November 10, 1998

## Abstract

We determine the structure of finitely generated groups which are quasi-isometric to symmetric spaces of noncompact type, allowing Euclidean de Rham factors. If  $X$  is a symmetric space of noncompact type with no Euclidean de Rham factor, and  $\Gamma$  is a finitely generated group quasi-isometric to the product  $\mathbb{E}^k \times X$ , then there is an exact sequence  $1 \rightarrow H \rightarrow \Gamma \rightarrow L \rightarrow 1$  where  $H$  contains a finite index copy of  $\mathbb{Z}^k$  and  $L$  is a uniform lattice in the isometry group of  $X$ .<sup>1</sup>

## 1 Introduction

If  $X$  is a symmetric space with no Euclidean de Rham factor, then any finitely generated group  $\Gamma$  quasi-isometric to  $X$  is a finite extension of a uniform lattice in  $Isom(X)$ . This result is a direct corollary of the main results of [KlLe97b] together with earlier work in the rank 1 cases [Tuk88, Gro81a, Hin90, Pan89, Ga92, CJ94], and was first announced in June 1994 at MSRI, and in [KlLe97a]. This result does not extend to symmetric spaces with a nontrivial Euclidean factor: it was observed by Epstein, Gersten, and Mess that any extension of a Fuchsian group by  $\mathbb{Z}$  is quasi-isometric to  $\mathbb{H}^2 \times \mathbb{R}$ , and such extensions are typically not finite extensions of lattices in  $Isom(\mathbb{H}^2 \times \mathbb{R})$ . In this paper we treat the case of groups quasi-isometric to symmetric spaces with a Euclidean de Rham factor.

**Theorem 1.1** *Let  $X$  be a symmetric space of noncompact type with no Euclidean de Rham factor, and let  $Nil$  be a simply connected nilpotent Lie group equipped with a left-invariant Riemannian metric. Suppose  $\Gamma$  is a finitely generated group quasi-isometric to  $Nil \times X$ . Then there is an exact sequence*

$$1 \longrightarrow H \longrightarrow \Gamma \xrightarrow{p} L \longrightarrow 1 \tag{1.1}$$

*where  $H$  is a finitely generated group quasi-isometric to  $Nil$  and  $L$  is a uniform lattice in the isometry group of  $X$ , and this sequence is unique up to isomorphism.*

---

\*Supported by a Sloan foundation fellowship, and NSF grants DMS-95-05175 and DMS-96-26911.

<sup>‡</sup>Supported by SFB 256 (Bonn).

<sup>1</sup>1991 Mathematics Subject Classification: 20F32, 53C35, 53C21

Furthermore, given any quasi-isometry  $\Gamma \xrightarrow{\phi} Nil \times X$ , there is a quasi-isometry  $L \xrightarrow{\bar{\phi}} X$  so that the diagram

$$\begin{array}{ccc} \Gamma & \xrightarrow{p} & L \\ \phi \downarrow & & \bar{\phi} \downarrow \\ Nil \times X & \xrightarrow{\pi_2} & X \end{array} \quad (1.2)$$

commutes up to bounded error. In particular,  $H$  is undistorted<sup>2</sup> in  $\Gamma$ .

When  $Nil$  is the trivial group then  $\Gamma$  is a finite extension of a uniform lattice in  $Isom(X)$ , and when  $Nil \simeq \mathbb{R}^k$  then  $H$  is virtually abelian of rank  $k$  by [Gro81b, Pan83]. The case when  $X$  is the hyperbolic plane and  $Nil \simeq \mathbb{R}$  is due to Rieffel [Rie93].

We further refine Theorem 1.1 when  $Nil \simeq \mathbb{R}^n$ .

**Theorem 1.2** *Let  $X$  be as in Theorem 1.1. Then any finitely generated group  $\Gamma$  quasi-isometric to  $\mathbb{R}^n \times X$  contains a finite index subgroup  $\Gamma_1 \subset \Gamma$  which is a central extension of the form*

$$1 \longrightarrow \mathbb{Z}^n \longrightarrow \Gamma_1 \longrightarrow L_1 \longrightarrow 1 \quad (1.3)$$

where  $L_1$  is a finite extension of a lattice in  $Isom(X)$ .

In general, one cannot arrange that the group  $L_1$  is a lattice in  $Isom(X)$  rather than a finite extension of a lattice. Examples of Raghunathan [Rag84] show that this is impossible in general even when  $n = 0$ .

Theorem 1.2 raises the question of which central extensions (1.3) are quasi-isometric to  $\mathbb{R}^n \times X$ . Theorem 1.4 below gives a homological answer to this.

**Definition 1.3** *An extension  $1 \rightarrow K \rightarrow G \xrightarrow{p} Q \rightarrow 1$  of finitely generated groups is **quasi-isometrically trivial** if there is a quasi-isometry  $G \xrightarrow{\phi} K \times Q$  so that the diagram*

$$\begin{array}{ccc} G & \xrightarrow{p} & Q \\ \phi \downarrow & & id_Q \downarrow \\ K \times Q & \xrightarrow{\pi_2} & Q \end{array} \quad (1.4)$$

commutes up to bounded error.

The central extension (1.3) is quasi-isometrically trivial by the second part of Theorem 1.1. The next result gives a general characterisation of quasi-isometrically trivial extensions.

---

<sup>2</sup>The inclusion of  $H$  in  $\Gamma$  is biLipschitz with respect to the word metrics.

**Theorem 1.4** (See section 7 for the definition of  $L^\infty$  cohomology for CW complexes.)  
 Let

$$1 \rightarrow \mathbb{Z}^n \rightarrow G \rightarrow Q \rightarrow 1 \tag{1.5}$$

be a central extension of finitely generated groups, and let  $\alpha \in H^2(Q; \mathbb{Z}^n)$  be the associated cohomology class. Let  $K$  be a CW-complex with finite 1-skeleton which is an Eilenberg-MacLane space for  $Q$ , and identify  $\alpha$  with a class in  $H^2(K; \mathbb{Z}^n) \simeq H^2(Q; \mathbb{Z}^n)$ . Then the extension (1.5) is quasi-isometrically trivial iff  $\alpha$  is in the image of the homomorphism  $H_{L^\infty}^2(K; \mathbb{Z}^n) \rightarrow H^2(K; \mathbb{Z}^n)$ , and any lift  $\hat{\alpha} \in H_{L^\infty}^2(K; \mathbb{Z}^n)$  of  $\alpha$  pulls back to zero in  $H_{L^\infty}^2(\tilde{K}; \mathbb{Z}^n)$ , where  $\tilde{K}$  denotes the universal cover of  $K$ .

*Remark.* Using bounded cohomology instead of  $L^\infty$  cohomology, Gersten [Ger92] gave a sufficient condition for a central extension by  $\mathbb{Z}$  to be quasi-isometric to a trivial extension.

An earlier version of this paper was posted on the AMS preprint server in October 1996.

We gratefully acknowledge support by the RiP-program at the Mathematisches Forschungsinstitut Oberwolfach.

## Contents

1	Introduction . . . . .	1
2	Preliminaries . . . . .	3
3	Projecting quasi-actions to the factors . . . . .	5
4	Straightening cocompact quasi-actions on irreducible symmetric spaces	5
5	A Growth estimate for small elements in nondiscrete cocompact subgroups of $Isom(X)$ . . . . .	7
	5.1 Parabolic isometries of symmetric spaces . . . . .	7
	5.2 The growth estimate . . . . .	7
6	Proof Theorem 1.1 . . . . .	10
7	Proof of Theorem 1.2 . . . . .	10
8	Geometry of central extensions by $\mathbb{Z}^n$ . . . . .	12
	Bibliography . . . . .	15

## 2 Preliminaries

In this section we recall some basic definitions and notation. See [Gro93] for more discussion and background.

**Definition 2.1** A map  $f : X \rightarrow Y$  between metric spaces is an  $(L, A)$  **quasi-isometry** if for every  $x_1, x_2 \in X$

$$L^{-1}d(x_1, x_2) + A \leq d(x_1, x_2) \leq Ld(x_1, x_2) + A,$$

and for every  $y \in Y$  we have  $d(y, f(X)) < A$ . Two quasi-isometries  $f_1, f_2 : X \rightarrow Y$  are **equivalent** if  $d(f_1, f_2) < \infty$ .

If  $\Gamma$  is a finitely generated group, then any two word metrics on  $\Gamma$  are biLipschitz to one another by  $id_\Gamma : \Gamma \rightarrow \Gamma$ . We will implicitly endow our finitely generated groups with word metrics.

**Definition 2.2** An  $(L, A)$ -**quasi-action** of a group  $\Gamma$  on a metric space  $Z$  is a map  $\rho : \Gamma \times Z \rightarrow Z$  so that  $\rho(\gamma, \cdot) : Z \rightarrow Z$  is an  $(L, A)$  quasi-isometry for every  $\gamma \in \Gamma$ ,  $d(\rho(\gamma_1, \rho(\gamma_2, z)), \rho(\gamma_1\gamma_2, z)) < A$  for every  $\gamma_1, \gamma_2 \in \Gamma, z \in Z$ , and  $d(\rho(e, z), z) < A$  for every  $z \in Z$ .

We will denote the self-map  $\rho(\gamma, \cdot) : Z \rightarrow Z$  by  $\rho(\gamma)$ .  $\rho$  is **discrete** if for any point  $z \in Z$  and any radius  $R > 0$ , the set of all  $\gamma \in \Gamma$  such that  $\rho(\gamma, z)$  is contained in the ball  $B_R(z)$  is finite.  $\rho$  is **cobounded** if  $Z$  coincides with a finite tubular neighborhood of the ‘‘orbit’’  $\rho(\Gamma)z \subset Z$  for every  $z$ . If  $\rho$  is a discrete cobounded quasi-action of a finitely generated group  $\Gamma$  on a geodesic metric space  $Z$ , it follows easily that the map  $\Gamma \rightarrow Z$  given by  $\gamma \mapsto \rho(\gamma, z)$  is a quasi-isometry for every  $z \in Z$ .

**Definition 2.3** Two quasi-actions  $\rho$  and  $\rho'$  are **equivalent** if there exists a constant  $D$  so that  $d(\rho(\gamma), \rho'(\gamma)) < D$  for all  $\gamma \in \Gamma$ .

**Definition 2.4** Let  $\rho$  and  $\rho'$  be a quasi-actions of  $\Gamma$  on  $Z$  and  $Z'$  respectively, and let  $\phi : Z \rightarrow Z'$  be a quasi-isometry. Then  $\rho$  is **quasi-isometrically conjugate to  $\rho'$  via  $\phi$**  if there is a  $D$  so that  $d(\phi \circ \rho(\gamma), \rho'(\gamma) \circ \phi) < D$  for all  $\gamma \in \Gamma$ .

**Lemma 2.5** (cf [Gro87, 8.2.K]) Let  $X$  be a Hadamard manifold of dimension  $\geq 2$  with sectional curvature  $\leq K < 0$ , and let  $\partial_\infty X$  denote the geometric boundary of  $X$  with the cone topology. Recall that every quasi-isometry  $\Phi : X \rightarrow X$  induces a boundary homeomorphism  $\partial_\infty \Phi : \partial_\infty X \rightarrow \partial_\infty X$ .

1. If  $\rho : \Gamma \times X \rightarrow X$  is a quasi-action on  $X$ , then  $\rho$  is discrete (respectively cobounded) iff  $\partial_\infty \rho$  acts properly discontinuously (respectively cocompactly) on the space of distinct triples in  $\partial_\infty X$ .
2. Given  $(L, A)$  there is a  $D$  so that if  $\phi_k, \psi$  are  $(L, A)$  quasi-isometries, then  $\partial_\infty \phi_k$  converges uniformly to  $\partial_\infty \psi$  iff  $\limsup d(\phi_k x, \psi x) < D$  for every  $x \in X$ . In particular, if  $\phi_1, \phi_2 : X \rightarrow X$  are  $(L, A)$  quasi-isometries with the same boundary mappings, then  $d(\phi_1, \phi_2) < D$ .

*Proof.* Let  $\partial^3 X \subset \partial_\infty X \times \partial_\infty X \times \partial_\infty X$  denote the subspace of distinct triples. The uniform negative curvature of  $X$  implies that there is a  $D_0$  depending only on  $K$  such that

(a) For every  $x \in X$  there is a triple  $(\xi_1, \xi_2, \xi_3) \in \partial^3 X$  such that  $d(x, \overline{\xi_i \xi_j}) < D_0$  for every  $1 \leq i \neq j \leq 3$ , where  $\overline{\xi_i \xi_j}$  denotes the geodesic with ideal endpoints  $\xi_i, \xi_j$ . Moreover for every  $C$  the set  $\{(\xi_1, \xi_2, \xi_3) \mid d(x, \overline{\xi_i \xi_j}) < C \text{ for all } 1 \leq i \neq j \leq 3\}$  has compact closure in  $\partial^3 X$ .

and

(b) For every  $(\xi_1, \xi_2, \xi_3) \in \partial^3 X$  there is a point  $x \in X$  so that  $d(x, \overline{\xi_i \xi_j}) < D_0$  for each  $1 \leq i \neq j \leq 3$ . And for every  $C$  there is a  $C'$  depending only on  $C$  and  $K$  so that  $\{x \in X \mid d(x, \overline{\xi_i \xi_j}) < C \text{ for every } 1 \leq i \neq j \leq 3\}$  has diameter  $< C'$ .

1 and 2 follow easily from this. □

### 3 Projecting quasi-actions to the factors

Let  $Nil$  and  $X$  be as in Theorem 1.1 and decompose  $X$  into irreducible factors:

$$X = \prod_{i=1}^l X_i \tag{3.1}$$

Suppose  $\rho$  is a quasi-action of the finitely generated group  $\Gamma$  on  $Nil \times X$ . We denote by  $p : Nil \times X \rightarrow X$  the canonical projection. By applying [KlLe97b, Theorem 1.1.2]<sup>3</sup> to each quasi-isometry  $\rho(\gamma)$  we construct quasi-actions  $\rho_i$  of  $\Gamma$  on  $X_i$  so that

$$d(p \circ \rho(\gamma), \prod_{i=1}^k \rho_i(\gamma) \circ p) < D$$

for all  $\gamma \in \Gamma$  and some positive constant  $D$ .

### 4 Straightening cocompact quasi-actions on irreducible symmetric spaces

The following result is a direct consequence of [Pan89, Théorème 1] and [KlLe97b, Theorem 1.1.3].

**Fact 4.1** *Let  $X$  be an irreducible symmetric space other than a real or complex hyperbolic space. Then every quasi-action on  $X$  is equivalent to an isometric action.*

*Proof.* Let  $\rho$  be a quasi-action of a group  $\Gamma$  on  $X$ . By the results just cited, there is an isometry  $\bar{\rho}(\gamma)$  at finite distance from the quasi-isometry  $\rho(\gamma)$  for every  $\gamma \in \Gamma$ . This isometry is unique and its distance from  $\rho(\gamma)$  is uniformly bounded<sup>4</sup> in terms of the constants of the quasi-action. So  $\bar{\rho}$  is an isometric action equivalent to  $\rho$ .  $\square$

We recall that the real and complex hyperbolic spaces of all dimensions admit quasi-isometries which are not equivalent to isometries [Pan89].

**Fact 4.2** *Any cobounded quasi-action  $\rho$  on a real or complex hyperbolic space is quasi-isometrically conjugate to an isometric action.*

This result is proven in [Tuk88] in the real-hyperbolic case. Using Pansu's theory of Carnot differentiability one can carry out Tukia's arguments for all rank-one symmetric spaces other than hyperbolic plane, cf. [Pan89, sec. 11]. Another proof for the complex-hyperbolic case can be found in [Chow96].

**Fact 4.3** *Let  $\rho$  be a cobounded quasi-action of a group  $\Gamma$  on  $\mathbb{H}^2$ . Then  $\rho$  is quasi-isometrically conjugate to a cocompact isometric action of  $\Gamma$  on  $\mathbb{H}^2$ .*

---

<sup>3</sup>Although Theorem 1.1.2 is only formulated in the case that  $Nil \simeq \mathbb{R}^n$ , the same proof works in general provided one uses [Pan83] to conclude that all asymptotic cones of  $Nil$  are homeomorphic to  $\mathbb{R}^k$  where  $k = Dim(Nil)$ .

<sup>4</sup>The uniformity in the rank one case follows from Lemma 2.5.

*Proof.* We recall that every quasi-isometry  $\phi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  induces a quasi-symmetric homeomorphism  $\partial_\infty \phi : \partial_\infty \mathbb{H}^2 \rightarrow \partial_\infty \mathbb{H}^2$ , see [TuVa82]; moreover the quasi-symmetry constant of  $\partial_\infty \phi$  can be estimated in terms of the quasi-isometry constants of  $\phi$ . Since equivalent quasi-isometries yield the same boundary homeomorphism, every quasi-action  $\rho$  on  $\mathbb{H}^2$  induces a genuine action  $\partial_\infty \rho$  on  $\partial_\infty \mathbb{H}^2$  by uniformly quasi-symmetric homeomorphisms.

Let  $\bar{\Gamma}$  be the quotient of  $\Gamma$  by the kernel of the action  $\partial_\infty \rho$ , and let  $\pi : \Gamma \rightarrow \bar{\Gamma}$  be the canonical epimorphism. If two elements  $\gamma_1, \gamma_2 \in \Gamma$  have the same boundary map then  $d(\rho(\gamma_1), \rho(\gamma_2))$  is uniformly bounded by Lemma 2.5. Hence we may obtain a quasi-action  $\bar{\rho}$  of  $\bar{\Gamma}$  on  $\mathbb{H}^2$  by choosing  $\gamma \in \pi^{-1}(\bar{\gamma})$  for each  $\bar{\gamma} \in \bar{\Gamma}$ , and setting  $\bar{\rho}(\bar{\gamma}) = \rho(\gamma)$ . If  $\bar{\tau}$  is an isometric action of  $\bar{\Gamma}$  on  $\mathbb{H}^2$  and  $\phi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  quasi-isometrically conjugates  $\bar{\rho}$  into  $\bar{\tau}$ , then  $\phi$  will quasi-isometrically conjugate  $\rho$  into the isometric action  $\tau : \Gamma \times \mathbb{H}^2 \rightarrow \mathbb{H}^2$  given by  $\tau(\gamma) = \bar{\tau}(\pi(\gamma))$ . Hence it suffices to treat the case when  $\bar{\Gamma} = \Gamma$ , and so we will assume that  $\partial_\infty \rho$  is an effective action.

**Lemma 4.4** *The quasi-action  $\rho$  is discrete if and only if the action  $\partial_\infty \rho$  on  $\partial_\infty \mathbb{H}^2$  is discrete in the compact-open topology.*

*Proof.* Suppose  $\partial_\infty \rho$  is discrete, and let  $(\gamma_i)$  be a sequence in  $\Gamma$  so that  $\rho(\gamma_i)$  maps a point  $p \in \mathbb{H}^2$  into a fixed ball  $B_R(p)$ . Then by a selection argument we may assume – after passing to a subsequence if necessary – that there is a quasi-isometry  $\phi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  so that for every  $q \in \mathbb{H}^2$  we have  $\limsup_i d(\rho(\gamma_i)(q), \phi(q)) < D$  for some  $D$ . Hence the boundary maps  $\partial_\infty \rho(\gamma_i)$  converge to  $\partial_\infty \phi$ , and so the sequence  $\partial_\infty \rho(\gamma_i)$  is eventually constant. Since  $\rho$  is effective we conclude that  $\gamma_i$  is eventually constant. Therefore  $\rho$  is a discrete quasi-action.

If  $\rho$  is a discrete quasi-action on  $\mathbb{H}^2$ , then  $\partial_\infty \rho$  is discrete by Lemma 2.5. □

*Proof of 4.3 continued.*

*Case 1:  $\partial_\infty \rho$  is discrete.* In this case,  $\rho$  is a discrete convergence group action (Lemma 2.5) and by the work of [CJ94, Ga92], there is a discrete isometric action  $\tau$  of  $\Gamma$  on  $\mathbb{H}^2$  so that  $\partial_\infty \rho$  is topologically conjugate to  $\partial_\infty \tau$ . Since  $\rho$  is cobounded,  $\partial_\infty \rho$  acts cocompactly on the set of distinct triples of points in  $\partial_\infty \mathbb{H}^2$  (lemma 2.5); therefore  $\partial_\infty \tau$  also acts cocompactly on the space of triples and so  $\tau$  is a discrete, cocompact, isometric action of  $\Gamma$  on  $\mathbb{H}^2$ . We now have two discrete, cobounded, quasi-actions of  $\Gamma$  on  $\mathbb{H}^2$ , so they are quasi-isometrically conjugate by some quasi-isometry  $\psi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ .

*Case 2:  $\partial_\infty \rho$  is nondiscrete.* By [Hin90, Theorem 4],  $\partial_\infty \rho$  is quasi-symmetrically conjugate to  $\partial_\infty \tau$ , where  $\tau$  is an isometric action on  $\mathbb{H}^2$ . The conjugating quasi-symmetric homeomorphism is the boundary of a quasi-isometry  $\psi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ , [TuVa82], which quasi-isometrically conjugates  $\partial_\infty \rho$  into the isometric action  $\tau$ . Applying Lemma 2.5 again, we conclude that  $\tau$  is cocompact. □

Section 3, and facts 4.1, 4.2 and 4.3 imply:

**Corollary 4.5** *Let  $X$  be a symmetric space of noncompact type without Euclidean factor. Then any cobounded quasi-action on  $X$  is quasi-isometrically conjugate to a cocompact isometric action on  $X$ .*

## 5 A Growth estimate for small elements in nondiscrete compact subgroups of $Isom(X)$

### 5.1 Parabolic isometries of symmetric spaces

Let  $X$  be a symmetric space of noncompact type, and let  $G = Isom(X)$ .

An isometry  $g \in G$  is **semisimple** if its displacement function  $\delta_g$  attains its infimum and **parabolic** otherwise.

**Lemma 5.1** *Let  $A \subset G$  be a finitely generated abelian group all of whose nontrivial elements are parabolic. Then  $A$  has a fixed point at infinity.*

*Proof.* Recall that the nearest point projection to a closed convex subset is well-defined and distance non-increasing. This implies that if  $C$  is a non-empty  $A$ -invariant closed convex set, then for all displacement functions  $\delta_a$ ,  $a \in A$ , we have  $\inf \delta_a = \inf \delta_a|_C$ . Hence for all  $n \in \mathbb{N}$ , the intersubsection of the sublevel sets  $\{p \mid \delta_{a_i}(p) \leq \inf \delta_{a_i} + 1/n\}$  is non-empty and contains a point  $p_n$ . We have  $\delta_{a_i}(p_n) \rightarrow \inf \delta_{a_i}$  for all  $a_i$ , and since the isometries  $a_i$  are parabolic the sequence  $\{p_n\}$  subconverges to an ideal boundary point  $\xi \in \partial_\infty X$ . It follows that the  $a_i$  fix  $\xi$ .  $\square$

**Lemma 5.2** *Let  $a_1, \dots, a_k \in Isom(X)$  be commuting parabolic isometries. Then there is a sequence of isometries  $\{g_n\} \subset G$  so that for every  $i$  the sequence  $g_n a_i g_n^{-1}$  subconverges to a semisimple isometry  $\bar{a}_i$ .*

*Proof.* From the proof of the previous lemma, there is a sequence of points  $\{p_n\} \subset X$  converging to an ideal point  $\xi$  so that  $\delta_{a_i}(p_n) \rightarrow \inf \delta_{a_i}$  for all  $a_i$ . Pick isometries  $g_n \in G$  such that  $g_n \cdot p_n = p_0$ . The conjugates  $g_n a_i g_n^{-1}$  have the same infimum displacement as  $a_i$ . Since

$$\delta_{g_n a_i g_n^{-1}}(p_0) = \delta_{a_i}(p_n) \rightarrow \inf \delta_{a_i} \quad ,$$

the  $g_n a_i g_n^{-1}$  subconverge to a semisimple isometry.  $\square$

We call an isometry  $g \neq e$  **purely parabolic**<sup>5</sup> if the identity is the only semisimple element in  $Ad_G(G) \cdot g$ .

### 5.2 The growth estimate

**Proposition 5.3** *Let  $X$  be a symmetric space of noncompact type with no Euclidean de Rham factors. Let  $\Gamma \subset G = Isom(X)$  be a finitely generated, nondiscrete, cocompact subgroup. Let  $U \subset Isom(X)$  be a neighborhood of the identity, and set*

$$f(k) := \#\{g \in \Gamma : |g|_\Gamma < k, g \in U\},$$

where  $|\cdot|_\Gamma$  denotes a word norm on  $\Gamma$ . Then  $f$  grows faster than any polynomial, i.e. for every  $d > 0$   $\limsup_{k \rightarrow \infty} \frac{f(k)}{k^d} = \infty$ .

<sup>5</sup>This is a geometric way of defining unipotent isometries.

*Proof.* Let  $\bar{\Gamma}^\circ$  denote the identity component of the closure of  $\Gamma$  in  $G$ .

*Case 1:  $\bar{\Gamma}^\circ$  is nilpotent.* Let  $A$  be the last non-trivial subgroup in the derived series of  $\bar{\Gamma}^\circ$ . Then  $A \subset \bar{\Gamma}$  is a connected abelian subgroup of positive dimension,  $A$  is normal in  $\bar{\Gamma}$ , and  $\Gamma \cap A$  is dense in  $A$ .

**Lemma 5.4** *For every  $\delta \in (0, 1)$  there is a  $\gamma \in \Gamma$  such that all eigenvalues of the automorphism  $Ad_G(\gamma)|_A : A \rightarrow A$  have absolute value  $< \delta$ .*

*Proof.* See section 5.1 for terminology.

*Step 1:  $A$  contains no semisimple isometries other than  $e$ .* Otherwise we can consider the intersection  $C$  of the minimum sets for the displacement functions  $\delta_a$  where  $a$  runs through all semisimple elements in  $A$ .  $C$  is a nonempty convex subset of  $X$  which splits metrically as  $C \cong \mathbb{E}^k \times Y$ . The flats  $\mathbb{E}^k \times \{y\}$  are the minimal flats preserved by all semisimple elements in  $A$ . Since  $\Gamma$  normalises  $A$  it follows that  $C$  is  $\Gamma$ -invariant. The cocompactness of  $\Gamma$  implies that  $C = X$  and  $k = 0$  because  $X$  has no Euclidean factor. This means that the semisimple elements in  $A$  fix all points, a contradiction.

*Step 2: All non-trivial isometries in  $A$  are purely parabolic.* If  $a \in A$ ,  $a \neq e$ , is not purely parabolic then there is a sequence of isometries  $g_n$  so that  $g_n a g_n^{-1}$  converges to a semisimple isometry  $\bar{a} \neq e$ . We can uniformly approximate the  $g_n$  by elements in  $\Gamma$ , i.e. there exist  $\gamma_n \in \Gamma$  and a bounded sequence  $k_n \in G$  subconverging to  $k \in G$  so that  $\gamma_n = k_n g_n$ . Then  $\gamma_n a \gamma_n^{-1} = k_n g_n a g_n^{-1} k_n^{-1}$  subconverges to the non-trivial semisimple element  $k \bar{a} k^{-1}$ . This contradicts step 1.

*Step 3:* Pick a basis  $\{a_1, \dots, a_k\}$  for  $A \simeq \mathbb{R}^k$ . By Lemma 5.2 there exist elements  $g_n \in G$  so that  $g_n a_i g_n^{-1} \rightarrow e$  for all  $a_i$ . We approximate the  $g_n$  as above by  $\gamma_n$  so that the sequence  $\gamma_n g_n^{-1}$  is bounded. Then  $\gamma_n a_i \gamma_n^{-1} \rightarrow e$  for all  $a_i$ . The lemma follows by setting  $\gamma = \gamma_n$  for sufficiently large  $n$ .  $\square$

*Proof of case 1 continued.* By Lemma 5.4, there is a  $\gamma \in \Gamma$ ,  $\gamma \neq e$ , and a norm  $\|\cdot\|_A$  on  $A$  such that for all  $a \in A$  we have

$$\|\gamma a \gamma^{-1}\|_A < \frac{1}{2} \|a\|_A.$$

Consider a neighborhood  $U$  of  $e$  in  $G$ . Let  $r > 0$  be small enough so that  $\{a \in A : \|a\|_A < r\} \subset U$  and pick  $\alpha \in \Gamma \cap A$  with  $\|\alpha\|_A < r/2$ . Then the elements

$$\gamma_{\epsilon_0 \dots \epsilon_{n-1}} = \alpha^{\epsilon_0} \cdot (\gamma \alpha \gamma^{-1})^{\epsilon_1} \dots (\gamma^{n-1} \alpha \gamma^{1-n})^{\epsilon_{n-1}}$$

for  $\epsilon_i \in \{0, 1\}$  are  $2^n$  pairwise distinct elements contained in  $\Gamma \cap U$  with word norm  $|\gamma_{\epsilon_0 \dots \epsilon_{n-1}}|_\Gamma < n^2(|\alpha|_\Gamma + |\gamma|_\Gamma)$ . This implies superpolynomial growth of  $f$ .

*Case 2:  $\bar{\Gamma}^\circ$  is not nilpotent.* Define an increasing sequence (the upper central series) of nilpotent Lie subgroups  $Z_i \subset \bar{\Gamma}^\circ$  inductively as follows: Set  $Z_0 = \{e\}$  and let  $Z_{i+1}$  be the inverse image in  $\bar{\Gamma}^\circ$  of the center in  $\bar{\Gamma}^\circ/Z_i$ . The dimension of  $Z_i$  stabilizes and we choose  $k$  so that  $\dim Z_k$  is maximal. Then the center of  $\bar{\Gamma}/Z_k$  is discrete and, since  $\bar{\Gamma}^\circ$  is not nilpotent, we have  $\dim Z_k < \dim \bar{\Gamma}$ . Proposition 5.3 now follows by applying the next lemma with  $H = \bar{\Gamma}$  and  $H_1 = Z_k$ .  $\square$



**Lemma 5.5** *Let  $H$  be a Lie group, let  $H_1 \triangleleft H$  be a closed normal subgroup so that  $\bar{H} := H/H_1$  is a positive dimensional Lie group with discrete center, and suppose  $\Gamma \subset H$  is a dense, finitely generated subgroup. If  $U$  is any neighborhood of  $e$  in  $H$ , then the function  $f(k) := \#\{g \in \Gamma : |g|_\Gamma \leq k, g \in U\}$  grows superpolynomially.*

*Proof.* The idea of the proof is to use the contracting property of commutators to produce a sequence  $\{\alpha_k\}$  in  $H \cap \Gamma$  which converges exponentially to the identity. The word norm  $|\alpha_k|_\Gamma$  grows exponentially with  $k$ , but the number of elements of  $\langle \alpha_1, \dots, \alpha_k \rangle$  in  $U$  also grows exponentially with  $k$ ; by comparing growth exponents we find that  $f$  grows superpolynomially.

Fix  $M \in \mathbb{N}$ , a positive real number  $\epsilon < 1/3$  and some left-invariant Riemannian metric on  $H$ . Since the differential of the commutator map  $(h, h') \mapsto [h, h']$  vanishes at  $(e, e)$  we can find a neighborhood  $V$  of  $e$  in  $H$  such that:

$$h, h' \in V \implies [h, h'] \in V \quad \text{and} \quad d([h, h'], e) < \frac{1}{2M}d(h, e) \quad (5.1)$$

Since the differential of the  $k$ -th power  $h \mapsto h^k$  at  $e$  is  $k \cdot id_{T_e H}$  for all  $k \in \mathbb{Z}$ , we can furthermore achieve that, whenever  $1 \leq k, k' \leq M$  and  $h, h^k, h^{k'} \in V$ , then

$$d(h^k, h^{k'}) \geq (|k - k'| - \epsilon) \cdot d(h, e) \quad (5.2)$$

By our assumption, there exist finitely many elements  $\gamma_1, \dots, \gamma_m \in \Gamma \cap V$  such that the centralizers  $Z_{\bar{H}}(\bar{\gamma}_j)$  of their images in  $\bar{H}$  have discrete intersubsection. We construct an infinite sequence of elements  $\alpha_i \in (\Gamma \cap V) \setminus H_1$  by picking  $\alpha_0 \in V$  arbitrarily and setting  $\alpha_{i+1} = [\alpha_i, h_{j(i)}] \notin H_1$  for suitably chosen  $1 \leq j(i) \leq m$ . Then

$$0 < d(\alpha_{i+1}, e) < \frac{1}{2M}d(\alpha_i, e) \quad (5.3)$$

by (5.1).

**Sublemma 5.6** *Pick  $n_0 \in \mathbb{N}$ . The  $M^n$  elements*

$$\gamma_{\epsilon_1 \dots \epsilon_n} = \alpha_{n_0+1}^{\epsilon_1} \cdots \alpha_{n_0+n}^{\epsilon_n} \quad \epsilon_i \in \{0, \dots, M-1\} \quad (5.4)$$

*are distinct.*

*Proof.* Assume that  $\gamma_{\epsilon_1 \dots \epsilon_n} = \gamma_{\epsilon'_1 \dots \epsilon'_n}$ ,  $\epsilon_l \neq \epsilon'_l$  and  $\epsilon_i = \epsilon'_i$  for all  $i < l$ . Then

$$\alpha_{n_0+l}^{\epsilon_l - \epsilon'_l} = \alpha_{n_0+l+1}^{\epsilon'_{l+1} - \epsilon_{l+1}} \cdots \alpha_{n_0+n}^{\epsilon'_n - \epsilon_n}.$$

On the other hand (5.2, 5.3) and the triangle inequality imply

$$d(\alpha_{n_0+l+1}^{\epsilon'_{l+1} - \epsilon_{l+1}} \cdots \alpha_{n_0+n}^{\epsilon'_n - \epsilon_n}, e) < M \cdot \sum_{j=1}^{\infty} \frac{1}{(2M)^j} \cdot d(\alpha_{n_0+l}, e) < \frac{1}{2}d(\alpha_{n_0+l}, e) < d(\alpha_{n_0+l}^{\epsilon_l - \epsilon'_l}, e),$$

a contradiction.  $\square$

To complete the proof of the lemma, we observe that the elements (5.4) have word norm  $|\gamma_{\epsilon_1 \dots \epsilon_n}|_\Gamma \leq \text{const}(n_0) \cdot 2^n$  and are contained in  $U$  if  $n_0$  is sufficiently large. This shows that  $f(k)$  grows polynomially of order at least  $\frac{\log(M)}{\log(2)}$  for all  $M$ , hence the claim.  $\square$

## 6 Proof Theorem 1.1

Let  $\rho_0 : \Gamma \times \Gamma \rightarrow \Gamma$  be the isometric action of  $\Gamma$  on itself by left translation, and let  $\phi : \Gamma \rightarrow Nil \times X$  be a quasi-isometry. Then there is a quasi-action  $\rho$  of  $\Gamma$  on  $Nil \times X$  such that  $\phi$  quasi-isometrically conjugates  $\rho_0$  into  $\rho$ . According to section 3,  $\rho$  projects (up to bounded error) to a cobounded quasi-action  $\bar{\rho}$  of  $\Gamma$  on  $X$ .  $\bar{\rho}$  is quasi-isometrically conjugate to a cocompact isometric action  $\hat{\rho}$ , cf. Corollary 4.5. Pick  $x \in X$ ,  $y \in Nil \times \{x\}$ , and  $R > 0$ . Since the quasi-action  $\rho$  covers  $\bar{\rho}$ , we know that for all  $\gamma \in \Gamma$  with  $\hat{\rho}(\gamma) \cdot x \in B_R(x)$ , the distance  $d(\rho(\gamma) \cdot y, Nil \times \{x\})$  is uniformly bounded. The map  $\Gamma \rightarrow Nil \times X$  given by  $g \mapsto \rho(\gamma) \cdot y$  being a quasi-isometry, we conclude that the function

$$N(k) := \#\{\gamma \in \Gamma \mid |\gamma|_\Gamma < k, \hat{\rho}(\gamma) \cdot x \in B_R(x)\} \quad (6.1)$$

grows at most as fast as the volume of balls in  $Nil$ , i.e. it is  $< Ck^d$  for some  $C, d \in \mathbb{R}$ . Proposition 5.3 implies that  $L := \hat{\rho}(\Gamma)$  is a discrete subgroup in  $Isom(X)$  and hence a uniform lattice. The kernel  $H$  of the action  $\hat{\rho}$  is then a finitely generated group quasi-isometric to the fiber  $Nil$ , since it clearly (quasi)-acts discretely and coboundedly on the fiber.

To see that the sequence (1.1) is unique up to isomorphism, let

$$1 \rightarrow H' \rightarrow \Gamma \xrightarrow{p'} L' \rightarrow 1$$

be an exact sequence with  $L' \subset Isom(X)$  a uniform lattice and  $H'$  a group quasi-isometric to  $Nil$ . Then by [Gro81b, Pan83]  $H'$  is a virtually nilpotent group. Now if  $\Gamma \xrightarrow{f} \Gamma$  is an isomorphism then  $p'(H) \subset L'$  is a normal, finitely generated, virtually nilpotent subgroup; it follows that  $p'(f(H))$  is trivial. Similarly  $p(f^{-1}(H'))$  is trivial and we conclude that  $f$  induces an isomorphism of the two exact sequences.

We now prove the last statement of Theorem 1.1. When we restrict  $\bar{\rho}$  to  $H$  we get a quasi-action which is equivalent to the trivial action of  $H$  on  $X$ . Hence  $\bar{\rho}$  induces a quasi-action  $\eta$  of  $L = \Gamma/H$  on  $X$ , which is discrete and cobounded. The action  $\eta_0$  of  $L$  on itself by left translations is also discrete and cobounded, so  $g \mapsto \eta(g)(\pi_2(\phi(e)))$  defines a quasi-isometry  $L \xrightarrow{\bar{\phi}} X$ . It follows that the diagram

$$\begin{array}{ccc} \Gamma & \xrightarrow{p} & L \\ \phi \downarrow & & \bar{\phi} \downarrow \\ Nil \times X & \xrightarrow{\pi_2} & X \end{array} \quad (6.2)$$

commutes up to bounded error since  $\phi$  quasi-isometrically conjugates  $\rho_0$  into  $\rho$ ,  $\rho$  projects to  $\bar{\rho}$ , and  $d(\bar{\rho}(\gamma H), \eta(\gamma H))$  is uniformly bounded (independent of  $\gamma$ ).  $\square$

## 7 Proof of Theorem 1.2

**Overview.** If  $\Gamma$  is quasi-isometric to  $\mathbb{R}^n \times X$  where  $X$  is a symmetric space with no Euclidean de Rham factor, then by Theorem 1.1,  $\Gamma$  fits into an exact sequence (1.1)

where  $H$  is an undistorted virtually  $\mathbb{Z}^n$  subgroup. We will use the undistortedness of  $H$  to pass to a finite index subgroup of  $\Gamma$  which is a central extension, cf. [Ger91].

If  $S$  is a subset of a group  $G$ , we will use the notation  $Z(S, G)$  to denote the centralizer of  $S$  in  $G$ , and  $Z(G)$  to denote the center of  $G$ .

*Proof of Theorem 1.2.* By Theorem 1.1 we get an exact sequence

$$1 \longrightarrow H \longrightarrow \Gamma \xrightarrow{p} L \longrightarrow 1$$

where  $H$  is a finitely generated group quasi-isometric to  $\mathbb{Z}^n$ , and  $L \subset \text{Isom}(X)$  is a uniform lattice. Applying the second part of the theorem we can get a quasi-isometry  $\Gamma \xrightarrow{f} \mathbb{Z}^n \times L$  so that

$$\begin{array}{ccc} \Gamma & \xrightarrow{p} & L \\ f \downarrow & & id \downarrow \\ \mathbb{Z}^n \times L & \xrightarrow{\pi_2} & L \end{array} \quad (7.1)$$

commutes up to bounded error. Clearly  $f(H) \subset \mathbb{Z}^n \times L$  has finite Hausdorff distance from  $\mathbb{Z}^n \times \{e\} \subset \mathbb{Z}^n \times L$ , so  $H$  is undistorted<sup>6</sup> in  $\Gamma$ . By [Gro81b, Pan83] that  $H$  contains a finite index copy of  $\mathbb{Z}^n$ .

Next we will identify a finite index abelian subgroup of  $H$  which is normal in  $\Gamma$ . Let  $T$  be the subgroup of “translations” in  $H$ , i.e.

$$T = \{h \in H \mid [H : Z(h, H)] < \infty\}. \quad (7.2)$$

Clearly  $T$  is a characteristic subgroup of  $H$ , and has finite index in  $H$ ; in particular  $T$  is finitely generated. Note that  $Z(T)$ , the center of  $T$ , has finite index in  $T$  since if  $T = \langle t_1, \dots, t_k \rangle$ , then  $Z(T) = \cap_i Z(t_i, T)$  is a finite intersection of finite index subgroups of  $T$ . Hence  $Z(T)$  is a finitely generated abelian group of the form  $\mathbb{Z}^n \oplus A$  where  $A$  is a finite abelian group. Note  $Z(T)$  is normal in  $\Gamma$  since it is characteristic in  $H$ , and  $H$  is normal in  $\Gamma$ .

**Lemma 7.1** *The centralizer of  $Z(T)$  in  $\Gamma$ ,  $Z(Z(T), \Gamma)$ , has finite index in  $\Gamma$ .*

The proof uses properties of translation numbers, see [Gro81a, pp. 189-191]. The paper [Ger91] uses a similar setup.

**Definition 7.2** *Let  $G$  be a finitely generated group, and let  $|\cdot|_G$  be a word norm on  $G$ . Then the **translation length** of  $g \in G$  is*

$$\delta_G(g) := \lim_{k \rightarrow \infty} \frac{|g^k|_G}{k}.$$

*The limit exists since  $k \mapsto |g^k|_G$  is a subadditive function.*

---

<sup>6</sup>A finitely generated subgroup of a finitely generated group is undistorted if the inclusion homomorphism is a quasi-isometric embedding.

The translation length is conjugacy invariant, vanishes on torsion elements, and changes by at most a bounded factor if one passes to a different word metric. If a homomorphism  $H \rightarrow G$  of finitely generated groups is a quasi-isometric embedding then the pullback of  $\delta_G$  to  $H$  is equivalent to  $\delta_H$ .

*Proof of Lemma 7.1.* We know that  $Z(T)$  is undistorted in  $\Gamma$  since  $Z(T)$  has finite index in  $H$  and  $H$  is undistorted in  $\Gamma$ . Hence  $\delta_\Gamma$  restricts to a function on  $Z(T)$  which is equivalent to  $\delta_{Z(T)}$ . The latter function clearly factors through the homomorphism  $Z(T) \rightarrow \mathbb{Z}^n$  whose kernel is the torsion subgroup  $A \subset Z(T)$ . Hence  $\delta_{Z(T)} : Z(T) \rightarrow \mathbb{R}$  is a proper function on  $Z(T)$  which is invariant under conjugacy by elements of  $\Gamma$ . Therefore the action of  $\Gamma$  on  $Z(T)$  by conjugacy factors through a finite group, and we conclude that  $Z(Z(T), \Gamma)$  has finite index in  $\Gamma$ .  $\square$

*Proof of Theorem 1.2 concluded.* Let  $\Gamma_1 := Z(Z(T), \Gamma)$ , let  $H_1 \subseteq Z(T) \subseteq \Gamma_1 \cap H$  be a finite index subgroup of  $Z(T)$  isomorphic to  $\mathbb{Z}^n$ , and set  $L_1 := \Gamma_1/H_1$ . Then clearly  $L_1$  is a finite extension of a uniform lattice in  $Isom(X)$ , and hence

$$1 \rightarrow H_1 \rightarrow \Gamma_1 \rightarrow L_1 \rightarrow 1$$

is an exact sequence as in (1.3).  $\square$

## 8 Geometry of central extensions by $\mathbb{Z}^n$

The objective of this section is Proposition 8.2, which provides criteria for recognizing quasi-isometrically trivial central extensions.

**Definition 8.1** *Let  $X$  be a CW-complex. A cellular  $k$ -cochain  $\alpha \in C^k(X; \mathbb{Z}^n)$  is **bounded** if its values on the  $k$ -cells of  $X$  are uniformly bounded. The collection of bounded cochains forms a subcomplex  $C_{L^\infty}^*(X; \mathbb{Z}^n)$  of  $C^*(X; \mathbb{Z}^n)$ , and its cohomology is  $H_{L^\infty}^*(X; \mathbb{Z}^n)$ .*

Note that the homomorphism  $H_{L^\infty}^i(X; \mathbb{Z}^n) \rightarrow H^i(X; \mathbb{Z}^n)$  is surjective if  $X$  has a finite  $i$ -skeleton, and injective if  $X$  has a finite  $i - 1$ -skeleton.

If  $G$  is a finitely generated group, then we may find a CW-complex  $X$  with finite 1-skeleton which is an Eilenberg-MacLane space for  $G$ . We will be interested in elements of  $H^2(G; \mathbb{Z}^n)$  in the image of the monomorphism  $H_{L^\infty}^2(X; \mathbb{Z}^n) \rightarrow H^2(X; \mathbb{Z}^n)$  whose lift to  $H_{L^\infty}^2(X; \mathbb{Z}^n)$  lies in the kernel of the pullback homomorphism  $H_{L^\infty}^2(X; \mathbb{Z}^n) \rightarrow H_{L^\infty}^2(\tilde{X}; \mathbb{Z}^n)$ . Note that the subgroup of  $H^2(G; \mathbb{Z}^n)$  defined this way is independent of the choice of  $X$ ; for if  $X_1$  and  $X_2$  are two Eilenberg-MacLane spaces for  $G$  with finite 1-skeleton, then we can find a cellular homotopy equivalence  $X_1 \xrightarrow{f} X_2$ , and this will induce a  $G$ -equivariant map  $C_{L^\infty}^1(\tilde{X}_2; \mathbb{Z}^n) \rightarrow C_{L^\infty}^1(\tilde{X}_1; \mathbb{Z}^n)$ .

**Proposition 8.2** *Let*

$$1 \rightarrow \mathbb{Z}^n \xrightarrow{i} G \xrightarrow{p} Q \rightarrow 1 \tag{8.1}$$

*be a central extension of finitely generated groups. Then the following are equivalent:*

1. The extension is quasi-isometrically trivial, i.e. there is a quasi-isometry  $G \xrightarrow{f} \mathbb{Z}^n \times Q$  so that the diagram

$$\begin{array}{ccc} G & \xrightarrow{p} & Q \\ f \downarrow & & \text{id} \downarrow \\ \mathbb{Z}^n \times Q & \xrightarrow{\pi_Q} & Q \end{array} \quad (8.2)$$

commutes up to bounded error.

2. There is a Lipschitz section  $s : Q \rightarrow G$  of  $p$ .

3. If  $K$  is an Eilenberg-MacLane space for  $Q$ , and  $K$  has a finite 1-skeleton, then the cohomology class in  $H^2(K; \mathbb{Z}^n)$  associated with the central extension (8.1) is an  $L^\infty$  class which lies in the kernel of the pullback to the universal cover  $H_{L^\infty}^2(K; \mathbb{Z}^n) \rightarrow H_{L^\infty}^2(\tilde{K}; \mathbb{Z}^n)$ .

*Proof.* (1  $\implies$  2). Suppose  $f$  makes diagram (8.2) commute up to bounded error, and let  $f^{-1}$  be a quasi-inverse<sup>7</sup> for  $f$ . Define  $s_0 : Q \rightarrow G$  to be the composition  $Q \rightarrow \{e\} \times Q \rightarrow \mathbb{Z}^n \times Q \xrightarrow{f^{-1}} G$ . The approximate commutativity of (8.2) implies that  $d(p \circ s_0, \text{id}_Q) < \infty$ . Define a section  $s : Q \rightarrow G$  of  $p$  by letting  $s(q)$  be a point in  $p^{-1}(q)$  closest to  $s_0(q)$ , for all  $q \in Q$ . By Lemma 8.3 below, we have  $d(s, s_0) < \infty$ , and so  $s$  is Lipschitz since  $s_0$  is Lipschitz and  $d(q_1, q_2) \geq 1$  for distinct elements  $q_1, q_2 \in Q$ .

**Lemma 8.3** *If  $H \triangleleft G$  are finitely generated groups, then the coset distance metric on  $G/H$  is equivalent<sup>8</sup> to any word metric on  $G/H$ .*

*Proof.* Let  $\Sigma \subset G$  be a symmetric finite generating set, and let  $\bar{\Sigma} \subset G/H$  be the image of  $\Sigma$  under  $G \rightarrow G/H$ . Then there is a canonical 1-Lipschitz map between the Cayley graphs  $\text{Cay}(G, \Sigma)$  and  $\text{Cay}(G/H, \bar{\Sigma})$ . Paths in  $\text{Cay}(G/H, \bar{\Sigma})$  can be lifted to paths in  $\text{Cay}(G, \Sigma)$  of the same length which join the corresponding cosets of  $H$ .  $\square$

(2  $\implies$  1). If  $s : Q \rightarrow G$  is a Lipschitz section of  $p$ , we may define a map  $\pi_{\mathbb{Z}^n} : G \rightarrow \mathbb{Z}^n$  by the formula  $\pi_{\mathbb{Z}^n}(g)s(p(g)) = g$ , i.e.  $\pi_{\mathbb{Z}^n}$  is the unique map  $G \rightarrow \mathbb{Z}^n$  which sends  $s(Q)$  to  $e \in \mathbb{Z}^n$ , and which is equivariant with respect to left translation by elements of  $\mathbb{Z}^n$ .

**Lemma 8.4**  *$\pi_{\mathbb{Z}^n}$  is Lipschitz.*

*Proof.* Note that if  $g_1, g_2 \in G$ ,  $h \in \mathbb{Z}^n$ , and  $g_2 = g_1 h$ , then  $\pi_{\mathbb{Z}^n}(g_2) = \pi_{\mathbb{Z}^n}(g_1)h$ , so  $d_{\mathbb{Z}^n}(\pi_{\mathbb{Z}^n}(g_1), \pi_{\mathbb{Z}^n}(g_2)) = d_{\mathbb{Z}^n}(e, h)$ . The properness of the distance function  $d_{\mathbb{Z}^n}(\cdot, e)$  implies that there is a function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  so that for all  $h \in \mathbb{Z}^n$ ,

$$d_{\mathbb{Z}^n}(h, e) \leq \delta(d_G(h, e)). \quad (8.3)$$

To prove Lemma 8.4, it suffices to find an  $L$  such that  $d_{\mathbb{Z}^n}(\pi_{\mathbb{Z}^n}(g_1), \pi_{\mathbb{Z}^n}(g_2)) \leq L$  whenever  $d_G(g_1, g_2) = 1$ . Consider the unique  $g_3 \in g_1 \mathbb{Z}^n$  which satisfies  $\pi_{\mathbb{Z}^n}(g_3) =$

<sup>7</sup> $d(f^{-1} \circ f, \text{id}_G)$  and  $d(f \circ f^{-1}, \text{id}_{\mathbb{Z}^n \times Q})$  are both finite.

<sup>8</sup>The two metrics have uniformly bounded ratio.

$\pi_{\mathbb{Z}^n}(g_2)$ , i.e.  $g_3 \in g_1\mathbb{Z}^n \cap (\pi_{\mathbb{Z}^n}(g_2)s(Q))$ . Then  $d_G(g_3, g_2) \leq C$  for some constant  $C$  because the composition  $s \circ p$  is Lipschitz. Applying triangle inequalities and (8.3), we get

$$\begin{aligned} d_{\mathbb{Z}^n}(\pi_{\mathbb{Z}^n}(g_1), \pi_{\mathbb{Z}^n}(g_2)) &= d_{\mathbb{Z}^n}(\pi_{\mathbb{Z}^n}(g_1), \pi_{\mathbb{Z}^n}(g_3)) \\ &\leq \delta(d_G(g_1, g_3)) \leq \delta(1 + C). \end{aligned}$$

□

To finish the proof that (2  $\implies$  1), note that we have a bijection  $\hat{f} : \mathbb{Z}^n \times Q \rightarrow G$  given by  $\hat{f}(h, q) = hs(q)$ .  $\hat{f}$  is clearly  $Lip(s)$ -Lipschitz in the  $Q$  direction. That  $\hat{f}$  is Lipschitz in the  $\mathbb{Z}^n$  direction follows from the fact that  $\mathbb{Z}^n$  is a central subgroup of  $G$ :

$$\begin{aligned} d_G(\hat{f}(h_1, q), \hat{f}(h_2, q)) &= d_G(h_1s(q), h_2s(q)) \\ &= d_G(h_1h_2^{-1}, e) \leq d_{\mathbb{Z}^n}(h_1h_2^{-1}, e) = d_{\mathbb{Z}^n}(h_1, h_2). \end{aligned}$$

Letting  $f = \hat{f}^{-1}$ , we see that  $f = (\pi_{\mathbb{Z}^n}, p)$  is a biLipschitz bijection.

(2  $\iff$  3). This follows from the obstruction theoretic interpretation of the characteristic class of the extension. Let  $K$  be a CW complex with finite 1-skeleton and one vertex, and which is an Eilenberg-MacLane space for  $Q$ . Let  $P \rightarrow K$  be a principal  $T^n$ -bundle with characteristic class  $[\alpha] \in H^2(K; \mathbb{Z}^n)$ , so that the exact homotopy sequence  $\pi_1(T^n) \rightarrow \pi_1(P) \rightarrow \pi_1(K)$  for the fibration  $P \rightarrow K$  is isomorphic to (8.1). Let  $\sigma : Skel_1(K) \rightarrow P$  be a section of  $P$  over the 1-skeleton of  $K$ . In the fiber over the point  $Skel_0(K)$ , choose a bouquet of  $n$  circles with vertex at  $\sigma(Skel_0(K))$ , which gives a standard basis for the fundamental group of the fiber. Let  $M \subset P$  be the 1-complex consisting of the union of this bouquet of circles with the bouquet  $\sigma(Skel_1(K)) \subset P$ .

Let  $\hat{P} \rightarrow \hat{K}$  be the pullback of the bundle  $P \rightarrow K$  under the covering projection  $\hat{K} \rightarrow K$ , let  $\hat{\sigma} : Skel_1(\hat{K}) \rightarrow \hat{P}$  be the pullback of  $\sigma$ , and let  $\hat{M} \subset \hat{P}$  be the inverse image of  $M$  under the covering  $\hat{P} \rightarrow P$ . Finally, let  $\tilde{P} \rightarrow \hat{P}$  be the universal covering, and let  $\tilde{M} \subset \tilde{P}$  be the inverse image of  $\hat{M}$  under  $\tilde{P} \rightarrow \hat{P}$ . Note that if we put path metrics on  $Skel_1(\hat{K})$  and  $\tilde{M}$ , then the projection map  $Skel_0(\tilde{M}) \rightarrow Skel_0(\hat{K})$  is naturally biLipschitz equivalent to  $G \xrightarrow{p} Q$ .

Now suppose 3 holds, and that  $\alpha \in C_{L^\infty}^2(K; \mathbb{Z}^n) \subset C^2(K; \mathbb{Z}^n)$ . We may assume that our section  $\sigma : Skel_1(K) \rightarrow P$  was chosen so that the associated cellular obstruction cocycle is  $\alpha$ . Then  $\hat{\alpha}$ , the image of  $\alpha$  under the map  $C_{L^\infty}^2(K; \mathbb{Z}^n) \rightarrow C_{L^\infty}^2(\hat{K}; \mathbb{Z}^n)$ , is the obstruction cocycle for  $\hat{\sigma} : Skel_1(\hat{K}) \rightarrow \hat{P}$ . By assumption,  $\hat{\alpha} = \delta\theta$  for some  $\theta \in C_{L^\infty}^1(\hat{K}; \mathbb{Z}^n)$ . Hence we may modify  $\hat{\sigma}$  using  $\theta$  to get a new section  $\hat{\sigma}_1 : Skel_1(\hat{K}) \rightarrow \hat{P}$  with trivial obstruction cocycle. In particular, if  $\tilde{P} \rightarrow \hat{P}$  is the universal covering map, then  $\hat{\sigma}_1$  lifts to a section  $\tilde{\sigma} : Skel_1(\tilde{K}) \rightarrow \tilde{P}$  of the  $\mathbb{R}$ -bundle  $\tilde{P} \rightarrow \tilde{K}$ . The fact that  $\theta$  is an  $L^\infty$ -cochain implies that  $\tilde{\sigma}$  restricts to a 1-Lipschitz map from  $Skel_0(\tilde{K})$  to  $Skel_0(\tilde{M})$ . Since the projection  $Skel_0(\tilde{M}) \rightarrow Skel_0(\tilde{K})$  is biLipschitz equivalent to  $G \rightarrow Q$ , we get a Lipschitz section of  $p$ , so 2 holds.

Conversely, suppose 2 holds. Then we get a Lipschitz section  $\tau : Skel_0(\tilde{K}) \rightarrow Skel_0(\tilde{M})$  of the projection  $Skel_0(\tilde{M}) \rightarrow Skel_0(\tilde{K})$ . We may extend  $\tau$  to a section  $\tilde{\sigma} : Skel_1(\tilde{K}) \rightarrow \tilde{P}$ , and let  $\hat{\sigma}_1 : Skel_1(\hat{K}) \rightarrow \hat{P}$  be the composition of  $\tilde{\sigma}$  with  $\tilde{P} \rightarrow \hat{P}$ .

**Lemma 8.5**  $\hat{\sigma}_1$  is obtained from  $\hat{\sigma}$  by applying a bounded cochain  $\theta \in C_{L^\infty}^1(\hat{K}; \mathbb{Z}^n)$ .

*Proof.* If  $e$  is a closed 1-cell in  $Skel_1(\tilde{K})$ , we want to show that the fixed endpoint homotopy classes of the two sections  $\hat{\sigma}|_e : e \rightarrow \hat{P}$  and  $\hat{\sigma}_1|_e : e \rightarrow \hat{P}$  (as maps into the inverse image of  $e$  in  $\hat{P}$ ) agree up to bounded error. If  $\gamma : [0, 1] \rightarrow e$  is a characteristic map for  $e$ , lift the path  $\hat{\sigma} \circ \gamma : [0, 1] \rightarrow \hat{M} \subset \hat{P}$  to a path  $\tilde{\gamma} : [0, 1] \rightarrow \tilde{M} \subset \tilde{P}$  starting at  $\tilde{\sigma} \circ \gamma(0)$ . Then

$$\begin{aligned} d_{\tilde{M}}(\tilde{\gamma}(1), \tilde{\sigma} \circ \gamma(0)) &\leq d_{\tilde{M}}(\tilde{\gamma}(1), \tilde{\gamma}(0)) + d_{\tilde{M}}(\tilde{\gamma}(0), \tilde{\sigma} \circ \gamma(0)) \\ &= 1 + d_{\tilde{M}}(\tau(\gamma(0)), \tau(\gamma(1))) \\ &\leq 1 + L_\tau \end{aligned}$$

where  $L_\tau$  is the Lipschitz constant of  $\tau$ . But then  $\tilde{\gamma}(1) = (\tilde{\sigma} \circ \gamma(1))h$  for some  $h \in \mathbb{Z}^n$ , and we can bound  $d_{\mathbb{Z}^n}(h, e)$  by a constant  $C$  depending on  $L_\tau$ , cf. (8.3). In other words, the fixed endpoint homotopy classes of  $\hat{\sigma}|_e$  and  $\hat{\sigma}_1|_e$  (as maps from  $e$  to the inverse image of  $e$  in  $\hat{P}$ ) differ by some  $h \in \mathbb{Z}^n$  where  $\|h\|_{\mathbb{Z}^n} < C$ .  $\square$

It follows that 3 holds. This completes the proof of Proposition 8.2.  $\square$

## References

- [CJ94] A. Casson and D. Jungreis, *Convergence groups and Seifert fibered 3-manifolds*, Inv. Math. **118** no. 3 (1994), 441–456.
- [Chow96] R. Chow, *Groups quasi-isometric to complex hyperbolic space*, Trans. AMS, **348** (1996), no. 5, pp. 1757-1769.
- [Esk96] A. Eskin, *Quasi-isometric rigidity of nonuniform lattices in higher rank symmetric spaces*, J. AMS, **11**, (1998), no. 2, 321–361.
- [Ga92] D. Gabai, *Convergence groups are Fuchsian groups*, Ann. Math. **136** no. 3 (1992), 447–510.
- [Ger92] S. Gersten, *Bounded cocycles and combings of groups*, Int. J. Alg. Comp., **2**, (1992), no. 3, 307-326.
- [Ger91] S. Gersten, H. Short, *Rational subgroups of biautomatic groups*, Ann. of Math., (2), **134**, (1991), no. 1, 125–158.
- [GDH90] E. Ghys, P. de la Harpe, *Sur les groupes hyperboliques d'apres Mikhael Gromov*, Progress in Mathematics, **83**, Birkhäuser.
- [Gro93] M. Gromov, *Asymptotic invariants for infinite groups*, in: Geometric group theory, London Math. Soc. lecture note series 182, 1993.
- [Gro81a] M. Gromov, *Hyperbolic manifolds, groups, and actions*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference, 183-213, Ann. of Math. Stud. 97.
- [Gro81b] M. Gromov, *Groups of polynomial growth and expanding maps*, Publ. IHES, **53** (1981), pp. 53-73.

- [Gro87] M. Gromov, *Hyperbolic groups*, 75-263, In: Essays in group theory, MSRI Publ. 8, Springer, 1987.
- [Hin90] A. Hinkkanen, *The structure of certain quasi-symmetric groups*, Mem. Amer. Math. Soc., **83**, (1990), no. 422, 1-83.
- [KaLe96] M. Kapovich and B. Leeb, *Quasi-isometries preserve the geometric decomposition of Haken manifolds*, Invent. Math., **128**, (1997), no. 2, 393-416.
- [KILe97a] B. Kleiner and B. Leeb, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, C. R. Acad. Sci. Paris, **324**, (1997), no. 6, 639-643.
- [KILe97b] B. Kleiner and B. Leeb, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, Publ. IHES, No. 86, (1997), 115-197.
- [Pan83] P. Pansu, *Croissance des boules et des géodésiques fermées dans les nilvariétés*, Erg. Thy. Dyn. Sys., **3**, (1983), no. 3, 415-445.
- [Pan89] P. Pansu, *Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math., **129** (1989), 1-60.
- [Rag84] M. S. Ragnathan, *Torsion in cocompact lattices in coverings of  $\text{Spin}(2, n)$* , Math. Ann., **266**, (1984), no. 4, 403-419.
- [Rie93] E. Rieffel, *Groups coarse quasi-isometric to  $\mathbb{H}^2 \times \mathbb{R}$* , PhD Thesis, UCLA, 1993.
- [Sch95] R. Schwartz, *The quasi-isometry classification of rank one lattices*, Publ. of IHES, vol. **82** (1995) 133- 168.
- [Tuk88] P. Tukia, *Homeomorphic conjugates of Fuchsian groups*, J. Reine Angew. Math. **391** (1988), 1-54.
- [TuVa82] P. Tukia, J. Väisälä, *Quasiconformal extension from dimension  $n$  to  $n + 1$* , Ann. Math., **115**, (1982), 331-348.

Bruce Kleiner  
 Department of Mathematics  
 University of Utah  
 Salt Lake City, UT 84112-0090  
 bkleiner@math.utah.edu

Bernhard Leeb  
 Department of Mathematics  
 University of Mainz  
 leeb@Mathematik.Uni-Mainz.DE