

# Clifford algebra, geometric algebra, and applications

Douglas Lundholm and Lars Svensson

Department of Mathematics  
KTH Royal Institute of Technology  
SE-100 44 Stockholm, Sweden

## **Abstract**

These are lecture notes for a course on the theory of Clifford algebras, with special emphasis on their wide range of applications in mathematics and physics. Clifford algebra is introduced both through a conventional tensor algebra construction (then called geometric algebra) with geometric applications in mind, as well as in an algebraically more general form which is well suited for combinatorics, and for defining and understanding the numerous products and operations of the algebra. The various applications presented include vector space and projective geometry, orthogonal maps and spinors, normed division algebras, as well as simplicial complexes and graph theory.

Revised version date: June 7, 2016

## Preface

These lecture notes were originally prepared for a course held at the Department of Mathematics, KTH, during the spring of 2009, and intended for advanced undergraduates and Ph.D. students in mathematics and mathematical physics. They are primarily based on the lecture notes of L.S. from a similar course in 2000 [43], and the M.Sc. thesis of D.L. in 2006 [30]. Additional material has been adopted from [27, 28, 12, 21] as well as from numerous other sources that can be found in the references. A substantial revision of the notes was made in conjunction with giving the course again at KTH in the spring of 2016.

When preparing these notes, we have tried to aim for

- efficiency in the presentation, made possible by the introduction, already at an early stage, of a combinatorial approach to Clifford algebra and its standard operations.
- completeness, in the sense that we have tried to include most of the standard theory of Clifford algebras, along with proofs of all stated theorems, or (when in risk of a long digression) reference to where a proof can be found.
- diversity in the range of applications presented — from pure algebra and combinatorics to discrete and continuous geometry and physics.

In order to make the presentation rather self-contained and accessible also to undergraduate students, we have included an appendix with some basic notions in algebra, as well as some additional material which may be unfamiliar also to graduate students. Statements requiring a wider mathematical knowledge are placed as remarks in the text, and are not required for an understanding of the main material. There are also certain sections and proofs, marked with an asterisk, that go slightly outside the scope of the course and consequently could be omitted by the student (but which could be of interest for a researcher in the field).

We are grateful to our students for valuable input and suggestions for improvements, as well as to John Baez for helpful correspondence concerning division algebras. D.L. would also like to thank the organizers of ICCA8 in Campinas 2008, as well as the funds of the Royal Swedish Academy of Sciences, the Swedish Research Council (grants ? and 2013-4734) and the Knut and Alice Wallenberg Foundation (grants KAW 2005.0098 and KAW 2010.0063) for financial support.

D.L. and L.S.  
Stockholm, June 7, 2016

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Grassmann's idea of extensive quantities . . . . .	1
1.2	Clifford's application of the extensive algebra . . . . .	2
<b>2</b>	<b>Foundations</b>	<b>6</b>
2.1	The geometric algebra $\mathcal{G}(V, q)$ over a quadratic space . . . . .	6
2.1.1	Orthogonal bases and signatures . . . . .	9
2.1.2	The pseudoscalar . . . . .	12
2.2	Combinatorial Clifford algebra $\mathcal{Cl}(X, R, r)$ . . . . .	13
2.2.1	Equivalence of $\mathcal{G}$ and $\mathcal{Cl}$ . . . . .	15
2.2.2	Grade structure . . . . .	17
2.3	Standard operations . . . . .	19
2.3.1	Products and involutions . . . . .	19
2.3.2	Linear independence . . . . .	22
2.3.3	Duality . . . . .	23
<b>3</b>	<b>Vector space geometry</b>	<b>26</b>
3.1	Blades and subspaces . . . . .	26
3.2	Linear functions . . . . .	30
3.3	Projections and rejections . . . . .	34
3.4	Some projective geometry . . . . .	35
3.4.1	The cross ratio in $\mathbb{P}^1(\mathbb{R})$ . . . . .	36
3.4.2	Cross ratios and quadrics in $\mathbb{P}^2(\mathbb{R})$ . . . . .	37
3.4.3	Pascal's theorem . . . . .	39
3.4.4	*Polarizations in quadrics . . . . .	39
<b>4</b>	<b>Discrete geometry</b>	<b>42</b>
4.1	Simplicial complexes . . . . .	42
4.1.1	Chain maps and complex morphisms . . . . .	43
4.1.2	An index theorem and Sperner's lemma . . . . .	44
4.1.3	Homology . . . . .	45
4.2	The number of spanning trees in a graph . . . . .	46
4.3	Fermionic operators . . . . .	48
4.4	Supersymmetric quantum mechanics on a lattice . . . . .	49
4.5	Infinite-dimensional Clifford algebra . . . . .	49
4.6	Embedded discrete geometry . . . . .	51
4.7	*Generalized infinite-dimensional Clifford algebra . . . . .	52
<b>5</b>	<b>Classification of real and complex geometric algebras</b>	<b>54</b>
5.1	A prototype example . . . . .	54
5.2	Matrix algebra classification . . . . .	55
5.3	Graded tensor products and the mother algebra . . . . .	61
<b>6</b>	<b>Groups</b>	<b>63</b>
6.1	Groups in $\mathcal{G}$ and their actions on $\mathcal{G}$ . . . . .	63
6.2	The Cartan–Dieudonné Theorem . . . . .	65
6.3	The Lipschitz group . . . . .	71
6.4	Properties of the Pin and Spin groups . . . . .	73

6.5	The bivector Lie algebra $\mathcal{G}^2$ . . . . .	75
6.6	Examples in lower dimensions . . . . .	80
6.6.1	The euclidean line . . . . .	81
6.6.2	The anti-euclidean line . . . . .	81
6.6.3	The degenerate line . . . . .	81
6.6.4	The euclidean plane . . . . .	82
6.6.5	The anti-euclidean plane . . . . .	83
6.6.6	The lorentzian plane . . . . .	83
6.6.7	The space algebra . . . . .	84
6.6.8	The spacetime algebra . . . . .	86
6.6.9	*The Dirac algebra . . . . .	89
6.7	*Norm functions and factorization identities . . . . .	91
<b>7</b>	<b>Euclidean and conformal geometry</b>	<b>93</b>
7.1	Linearization of the Euclidean group . . . . .	93
7.2	Conformal algebra . . . . .	94
<b>8</b>	<b>Representation theory</b>	<b>95</b>
8.1	Real, complex and quaternionic representations . . . . .	95
8.2	Fermionic representations . . . . .	98
8.3	Examples . . . . .	100
8.3.1	Normed division algebras . . . . .	100
8.3.2	Vector fields on spheres . . . . .	102
<b>9</b>	<b>Spinors</b>	<b>104</b>
9.1	Spin representations . . . . .	104
9.2	Spinors as elements of ideals . . . . .	107
9.3	Mixed-action spinors . . . . .	107
<b>10</b>	<b>Some Clifford analysis on <math>\mathbb{R}^n</math></b>	<b>108</b>
<b>11</b>	<b>The algebra of Minkowski spacetime</b>	<b>109</b>
11.1	Relative space . . . . .	109
11.2	Dirac-Hestenes spinors . . . . .	109
11.3	Electromagnetic fields . . . . .	109
<b>A</b>	<b>Appendix</b>	<b>112</b>
A.1	Notions in algebra . . . . .	112
A.1.1	Basic notions . . . . .	112
A.1.2	Direct products and sums . . . . .	116
A.1.3	Lists . . . . .	117
A.1.4	The free associative $R$ -algebra generated by $X$ . . . . .	117
A.1.5	Quotients . . . . .	118
A.1.6	Tensor products of modules . . . . .	119
A.1.7	Sums and products of algebras . . . . .	120
A.1.8	The tensor algebra $\mathcal{T}(V)$ . . . . .	120
A.2	The concept of a category . . . . .	121
A.2.1	Some examples of categories . . . . .	121
A.2.2	Universal objects in a category . . . . .	122
A.3	Expansion of the inner product . . . . .	122

A.4	Extension of morphisms . . . . .	123
A.5	Matrix theorems . . . . .	124
	<b>References</b>	<b>127</b>
	<b>Index</b>	<b>130</b>

# 1 Introduction

Let us start with an introduction, in terms of modern language, to the ideas of Hermann Günther Grassmann (1809-1877) and William Kingdon Clifford (1845-1879). For a detailed account of the history of the subject, see e.g. [28].

## 1.1 Grassmann's idea of extensive quantities

Let us consider our ordinary physical space which we represent as  $\mathbb{R}^3$ . An ordered pair  $(P, Q)$  in  $\mathbb{R}^3 \times \mathbb{R}^3$  is called a *line segment*, and we think of it as directed from the point  $P$  to the point  $Q$ . We call two line segments equivalent if one can be transformed into the other by a translation. This leads to the original formal notion of a *vector* as an equivalence class of directed line segments. Note that such a vector can be characterized by a *direction*, in the form of a line through the origin and parallel to the line segment, an *orientation*, i.e. the direction along the line in which it points, and a *magnitude*, or length of the segment, represented by a positive number. We are very familiar with how to add, subtract and rescale vectors. But can they be multiplied? We of course know about the scalar product and the cross product, but we shall introduce a new product called the *outer product*.




Figure 1.1: A directed line segment  $(P, Q)$  and a pair of plane surface areas  $A$  and  $A'$  in the planes  $P$  and  $P'$ , respectively.

Let  $A$  denote a plane surface area, in a plane  $P$ , endowed with an “orientation” or “signature” represented by a rotation arrow as in Figure 1.1. If  $A'$  denotes another plane surface area in a plane  $P'$ , also endowed with a rotation arrow, then we shall say that  $A$  is equivalent to  $A'$  if and only if the planes  $P$  and  $P'$  are parallel, the areas of  $A$  and  $A'$  are the same, and if the rotation arrow is directed in the same way after translating  $A'$  to  $A$ , i.e.  $P'$  to  $P$ . Equivalence classes of such directed surface areas are to be called *2-blades* (or *simple 2-vectors*).

We observe that a 2-blade  $A$  can be represented as a parallelogram spanned by two vectors  $a$  and  $b$  in the same plane as  $A$ , according to Figure 1.2, and call this oriented parallelogram area the outer product of  $a$  and  $b$  and denote it by  $a \wedge b$ . If the area of  $A$  is zero then we write  $A = 0$ . Hence,  $a \wedge a = 0$ . By  $-A$  we denote the equivalence class of plane surface areas with the same area and in the same plane as  $A$ , but with an oppositely directed rotation arrow. Thus, e.g.  $-(a \wedge b) = b \wedge a$ .

We would now like to define the sum of two 2-blades  $A$  and  $B$ . If  $A$  and  $B$




Figure 1.2: A 2-blade  $A$  represented by an outer product  $a \wedge b$ .




Figure 1.3: Geometric interpretation of  $b \wedge a = -a \wedge b$ .

are represented by plane surface areas in the planes  $P$  and  $Q$ , respectively, then we can translate  $Q$  such that  $P \cap Q$  contains a line  $L$ . We then realize that we can choose a vector  $c$ , parallel to  $L$ , such that  $A = a \wedge c$  and  $B = b \wedge c$  for some vectors  $a$  in  $P$ , and  $b$  in  $Q$ . Now, we define

$$A + B = a \wedge c + b \wedge c := (a + b) \wedge c,$$

which has a nice geometric representation<sup>1</sup> according to Figure 1.4.

Analogously, outer products of three vectors,  $a \wedge b \wedge c$ , *trivectors* or *3-blades*, are defined as equivalence classes of “directed” volumes with the same volume as the parallel epiped generated by  $a$ ,  $b$  and  $c$ , and with the orientation given by the handedness of the frame  $\{a, b, c\}$  (see Figure 1.5). Furthermore, although we have chosen to work with ordinary three-dimensional space for simplicity, the notion of a blade generalizes to arbitrary dimensions, where a  $k$ -blade can be thought of as representing a  $k$ -dimensional linear subspace equipped with an orientation and a magnitude.

## 1.2 Clifford’s application of the extensive algebra

Say that we are given three symbols  $\{e_1, e_2, e_3\}$  and that we construct a game with these symbols by introducing the following set of rules:

- We may form words of the symbols by writing them next to each other (the empty word is denoted 1).
- We may also form linear combinations of such words, i.e. multiply/scale each word by a number, and add scaled words together – writing a plus sign between different words, and summing up the numbers in front of equal words.

<sup>1</sup>Note however that this picture unfortunately does not generalize straightforwardly to higher dimensions.




Figure 1.4: Geometric interpretation of the sum  $A + B$  of a pair of 2-blades.




Figure 1.5: A 3-blade  $a \wedge b \wedge c$ .

- Furthermore, if two different symbols appear next to each other in a word then we may swap places and scale the word by  $-1$ , i.e.  $e_i e_j = -e_j e_i$ ,  $i \neq j$ .
- Lastly, we impose a rule that if a symbol appears next to the same symbol then we may strike out them both, i.e.  $e_1 e_1 = e_2 e_2 = e_3 e_3 = 1$ .

For example,

$$e_1 e_2 e_1 e_3 = -e_1 e_1 e_2 e_3 = -e_2 e_3.$$

Similarly, we see that the only words that can appear after applying the reduction rules are

$$1, e_1, e_2, e_3, e_1 e_2, e_1 e_3, e_2 e_3, \text{ and } e_1 e_2 e_3,$$

and that what we have obtained is just a game of adding and multiplying linear combinations of such words.

As an example, consider a linear combination of the original symbols,

$$a = a_1 e_1 + a_2 e_2 + a_3 e_3,$$

where  $a_1, a_2, a_3$  are ordinary (say, real) numbers. Then the square of this linear combination is

$$\begin{aligned} a^2 &:= aa = (a_1 e_1 + a_2 e_2 + a_3 e_3)(a_1 e_1 + a_2 e_2 + a_3 e_3) \\ &= a_1^2 e_1 e_1 + a_2^2 e_2 e_2 + a_3^2 e_3 e_3 \\ &\quad + (a_1 a_2 - a_2 a_1) e_1 e_2 + (a_1 a_3 - a_3 a_1) e_1 e_3 + (a_2 a_3 - a_3 a_2) e_2 e_3 \\ &= (a_1^2 + a_2^2 + a_3^2) 1. \end{aligned}$$

Similarly, if  $b = b_1 e_1 + b_2 e_2 + b_3 e_3$  is another linear combination, then

$$\begin{aligned} ab &= (a_1 b_1 + a_2 b_2 + a_3 b_3) 1 \\ &\quad + (a_1 b_2 - a_2 b_1) e_1 e_2 + (a_1 b_3 - a_3 b_1) e_1 e_3 + (a_2 b_3 - a_3 b_2) e_2 e_3, \end{aligned} \tag{1.1}$$






Figure 1.6: The standard unit blades of  $\mathbb{R}^3$ .

where we recognize the number in front of the empty word (the *scalar part* of  $ab$ ) as the ordinary scalar product  $a \cdot b$  of spatial vectors  $a = (a_1, a_2, a_3)$ ,  $b = (b_1, b_2, b_3)$  in a cartesian coordinate representation, while the remaining part of the expression resembles the cross product  $a \times b$ , but in terms of the words  $e_1e_2$  instead of  $e_3$  etc.

The geometric interpretation of the above construction in terms of our ordinary space  $\mathbb{R}^3$  is that  $e_1, e_2, e_3$  represent orthonormal basis vectors, while  $e_1e_2, e_2e_3, e_3e_1$  represent orthonormal basis 2-blades corresponding to the coordinate planes, with a right-handed orientation and unit area (see Figure 1.6). The full word  $e_1e_2e_3$  represents a unit 3-blade, i.e. a unit volume with right-handed orientation, while the empty word could be thought of as representing an empty blade, corresponding to the trivial zero-dimensional subspace. From this interpretation follows that the non-scalar part of (1.1) is a sum of coordinate 2-blades, with each component given by the corresponding component of the cross product. Recalling that the direction, orientation and magnitude of the vector  $a \times b$  is determined by the direction and orientation of the vectors  $a, b$  and the area of the parallelogram spanned by  $a$  and  $b$  – similarly to how we defined a 2-blade above – it follows that the sum of these blades actually is equal to the blade  $a \wedge b$ , and the expression (1.1) becomes simply

$$ab = a \cdot b + a \wedge b.$$

Let us look at an example of why this representation of objects in space could be useful. Consider the  $e_1e_2$ -plane spanned by the orthogonal unit vectors  $e_1$  and  $e_2$ . A rotation in this plane by an angle  $\theta$  can of course be understood as a linear map  $R$  determined by

$$\begin{aligned} e_1 &\mapsto R(e_1) = \cos \theta e_1 + \sin \theta e_2, \\ e_2 &\mapsto R(e_2) = -\sin \theta e_1 + \cos \theta e_2. \end{aligned} \tag{1.2}$$

Now, note that

$$(e_1e_2)^2 = e_1e_2e_1e_2 = -e_1e_1e_2e_2 = -1,$$

which means that the unit blade  $e_1e_2$  behaves like a complex imaginary unit in this regard, and we can rewrite (1.2) as<sup>2</sup>

$$R(e_1) = \cos \theta e_1 + \sin \theta e_1e_2 = e_1(\cos \theta + \sin \theta e_1e_2) = e_1e^{\theta e_1e_2},$$

<sup>2</sup>Just like for complex numbers, we could define the exponential expression in terms of sine and cosine, or alternatively, by the usual power series expansion.

and

$$\mathbf{R}(e_2) = -\sin \theta e_2^2 e_1 + \cos \theta e_2 = e_2(\sin \theta e_1 e_2 + \cos \theta) = e_2 e^{\theta e_1 e_2}.$$

This is of course not much more insightful than the usual representation of a rotation in the plane in terms of complex numbers. However, note that we can also write

$$\begin{aligned} e_1 e^{\theta e_1 e_2} &= e_1 e^{\frac{\theta}{2} e_1 e_2} e^{\frac{\theta}{2} e_1 e_2} = e_1 \left( \cos \frac{\theta}{2} + \sin \frac{\theta}{2} e_1 e_2 \right) e^{\frac{\theta}{2} e_1 e_2} \\ &= \left( \cos \frac{\theta}{2} - \sin \frac{\theta}{2} e_1 e_2 \right) e_1 e^{\frac{\theta}{2} e_1 e_2} = e^{-\frac{\theta}{2} e_1 e_2} e_1 e^{\frac{\theta}{2} e_1 e_2}, \end{aligned}$$

and similarly,

$$e_2 e^{\theta e_1 e_2} = e^{-\frac{\theta}{2} e_1 e_2} e_2 e^{\frac{\theta}{2} e_1 e_2}.$$

The point with the above rewriting is that the resulting expression also applies to the basis vector  $e_3$ , which is orthogonal to the plane and thus unaffected by the rotation, i.e.

$$\mathbf{R}(e_3) = e_3 = e_3 e^{-\frac{\theta}{2} e_1 e_2} e^{\frac{\theta}{2} e_1 e_2} = e^{-\frac{\theta}{2} e_1 e_2} e_3 e^{\frac{\theta}{2} e_1 e_2},$$

where we used that  $e_3 e_1 e_2 = e_1 e_2 e_3$ . By linearity, the rotation  $\mathbf{R}$  acting on *any* vector  $x \in \mathbb{R}^3$  can then be written

$$x \mapsto \mathbf{R}(x) = e^{-\frac{\theta}{2} e_1 e_2} x e^{\frac{\theta}{2} e_1 e_2} = R^{-1} x R. \quad (1.3)$$

The object  $R := e^{\frac{\theta}{2} e_1 e_2}$  is called a *rotor*, and it encodes in a compact way both the plane of rotation and the angle. Also note that by the generality of the construction, and by choosing a basis accordingly, *any* rotation in  $\mathbb{R}^3$  can be expressed in this form by some rotor  $R = e^{a \wedge b}$ , where the blade  $a \wedge b$  represents both the plane in which the rotation is performed, as well as the direction and angle of rotation. As we will see (in Section 6), similar expressions hold true for general orthogonal transformations in arbitrary dimensions, and in euclidean space as well as in lorentzian spacetime.

## 2 Foundations

In this section we define geometric algebra and work out a number of its basic properties. We first consider the definition that is most common in the mathematical literature, where it is introduced as a quotient space on the tensor algebra of a vector space with a quadratic form. We see that this leads, in the finite-dimensional real or complex case, to the equivalent definition of an algebra with generators  $\{e_1, \dots, e_n\}$  satisfying the so-called *anticommutation relations*

$$e_i e_j + e_j e_i = 2g_{ij}$$

for some given metric tensor (symmetric matrix)  $g$ . For physicists, this is perhaps the most well-known definition.

We go on to consider an alternative definition of geometric algebra based on its purely algebraic and combinatorial features. The resulting algebra, which we call *Clifford algebra* due to its higher generality but less direct connection to geometry, allows us to introduce common operations and prove fundamental identities in a simple and straightforward way compared to the traditional tensor algebraic approach.

### 2.1 The geometric algebra $\mathcal{G}(V, q)$ over a quadratic space

The conventional definition of geometric algebra is carried out in the context of vector spaces endowed with an inner product, or more generally a quadratic form. We consider here a vector space  $V$  of arbitrary dimension over some field<sup>3</sup>  $\mathbb{F}$ .

**Definition 2.1.** A *quadratic form*  $q$  on a vector space  $V$  is a map  $q: V \rightarrow \mathbb{F}$  such that

- i)  $q(\alpha v) = \alpha^2 q(v) \quad \forall \alpha \in \mathbb{F}, v \in V,$  and
- ii) the map  $(v, w) \mapsto q(v + w) - q(v) - q(w)$  is linear in both  $v$  and  $w$ .

The corresponding bilinear form  $\beta_q(v, w) := \frac{1}{2}(q(v + w) - q(v) - q(w))$  is called the *polarization* of  $q$  (this normalization is convenient, but in the case  $\text{char } \mathbb{F} = 2$  we leave out the factor  $\frac{1}{2}$  in this definition). The pair  $(V, q)$  or  $V \xrightarrow{q} \mathbb{F}$  is called a *quadratic space*.

Many vector spaces are naturally endowed with quadratic forms, as the following examples show.

**Example 2.1.** If  $V$  has a bilinear form  $\langle \cdot, \cdot \rangle$  then  $q(v) := \langle v, v \rangle$  is a quadratic form and  $\beta_q$  is the symmetrization of  $\langle \cdot, \cdot \rangle$ . This could be positive definite (i.e. an inner product), or indefinite (a metric tensor of arbitrary signature).

**Example 2.2.** If  $V$  is a normed vector space over  $\mathbb{R}$ , with norm denoted by  $|\cdot|$ , and where the parallelogram identity  $|x+y|^2 + |x-y|^2 = 2|x|^2 + 2|y|^2$  holds, then  $q(v) := |v|^2$  is a quadratic form and  $\beta_q$  is an inner product on  $V$ . This classic fact is sometimes called the Jordan–von Neumann theorem (see Exercise 2.1).

<sup>3</sup>See Appendix A.1 if the notion of a field, ring, tensor, or any other algebraic notion, is unfamiliar.

Let

$$\mathcal{T}(V) := \bigoplus_{k=0}^{\infty} \bigotimes^k V = \text{Span}_{\mathbb{F}} \{v_1 \otimes v_2 \dots \otimes v_k : v_i \in V, k = 0, 1, 2, \dots\}$$

denote the tensor algebra over  $V$ , the elements of which are finite sums of tensors of arbitrary finite grades over  $V$  (see A.1.8). Consider the two-sided ideal generated by all elements of the form<sup>4</sup>  $v \otimes v - q(v)$  for vectors  $v$ , that is

$$\mathcal{I}_q(V) := \left\{ \sum_k A_k \otimes (v \otimes v - q(v)) \otimes B_k \quad : \quad v \in V, A_k, B_k \in \mathcal{T}(V) \right\}. \quad (2.1)$$

We define the geometric algebra over  $V$  by quoting out (see A.1.5) this ideal from  $\mathcal{T}(V)$ , so that, in the resulting algebra, the square of a vector  $v$  will be equal to the scalar  $q(v)$ .

**Definition 2.2.** The *geometric algebra*  $\mathcal{G}(V, q)$  over the vector space  $V$  with quadratic form  $q$  is defined by

$$\mathcal{G}(V, q) := \mathcal{T}(V) / \mathcal{I}_q(V).$$

When it is clear from the context what vector space or quadratic form we are working with, we shall often denote  $\mathcal{G}(V, q)$  by  $\mathcal{G}(V)$ , or just  $\mathcal{G}$ .

The product in  $\mathcal{G}$ , called the *geometric* or *Clifford product*, is inherited from the tensor product in  $\mathcal{T}(V)$  and we denote it by juxtaposition (or  $\cdot$  if absolutely necessary),

$$\begin{aligned} \mathcal{G} \times \mathcal{G} &\rightarrow \mathcal{G}, \\ (A, B) &\mapsto AB := [A \otimes B] = A \otimes B + \mathcal{I}_q. \end{aligned}$$

Note that this product is bilinear and associative. Furthermore,

$$v^2 = [v \otimes v] = [v \otimes v - q(v)1] + q(v)1_{\mathcal{G}} = 0 + q(v),$$

(we usually leave out the unit  $1_{\mathcal{G}} = [1]$ ) and

$$q(v+w) = (v+w)^2 = v^2 + vw + wv + w^2 = q(v) + vw + wv + q(w),$$

so that, together with the definition of  $\beta_q$ , we immediately find the following identities on  $\mathcal{G}$  for all  $v, w \in V$  — characteristic for a Clifford algebra:

$$v^2 = q(v) \quad \text{and} \quad vw + wv = 2\beta_q(v, w). \quad (2.2)$$

It is at this point not obvious what the structure of the algebra  $\mathcal{G}(V, q)$  as defined above is. For example, we expect to see that the vector space we started with lies embedded in it,  $V \hookrightarrow \mathcal{G}$  (as is clearly the case for the tensor algebra  $\mathcal{T}(V)$  we started with), but it is in fact not even immediately clear that  $\mathcal{G}$  becomes non-trivial. These formal issues will have to wait until the next subsection where we construct the algebra explicitly. However, we should already now emphasize that one of the most important consequences of the above definition of the geometric algebra is the following universality property:

---

<sup>4</sup>It is common (mainly in the mathematical literature as opposed to the physics literature) to choose a different sign convention here, resulting in reversed signature in many of the following expressions. One argument for the convention we use here is that e.g. squares of vectors in euclidean spaces become positive instead of negative.

**Theorem 2.1** (Universality of  $\mathcal{G}$ ). *Let  $V \xrightarrow{q} \mathbb{F}$  be an arbitrary quadratic space, and let  $\mathcal{A}$  be an associative algebra over  $\mathbb{F}$  with a unit denoted by  $1_{\mathcal{A}}$ . If  $f: V \rightarrow \mathcal{A}$  is linear and*

$$f(v)^2 = q(v)1_{\mathcal{A}} \quad \forall v \in V, \quad (2.3)$$

*then there exists a unique  $\mathbb{F}$ -algebra homomorphism*

$$F: \mathcal{G}(V, q) \rightarrow \mathcal{A}$$

*such that  $F([v]) = f(v)$  for all  $v \in V$ .*

*Remark.* An algebra homomorphism  $\varphi: \mathcal{A} \rightarrow \mathcal{A}'$  between two unital associative  $\mathbb{F}$ -algebras is a map satisfying the following:

$$\begin{aligned} F(\alpha) &= \alpha, & \forall \alpha \in \mathbb{F} \text{ (image of } \mathbb{F} \text{ in } \mathbb{F} \hookrightarrow \mathcal{A}^{(l)}), \\ F(x+y) &= F(x) + F(y), & \forall x, y \in \mathcal{A}, \\ F(xy) &= F(x)F(y), & \forall x, y \in \mathcal{A}. \end{aligned}$$

*Proof.* We start by extending  $f$  to a map on the noncommutative polynomial algebra,

$$\varphi: \mathbb{F}\{V\} \rightarrow \mathcal{A}$$

via  $\varphi(1) = 1_{\mathcal{A}}$ ,

$$\varphi(v_1 v_2 \dots v_m) := f(v_1) f(v_2) \dots f(v_m), \quad v_i \in V,$$

and extending  $\mathbb{F}$ -linearly. Let us use the notation  $+'$  and  $\cdot'$  for addition and multiplication in  $\mathbb{F}\{V\}$ . Then follows that for  $x, y \in V$ ,

$$\begin{aligned} \varphi((x +' y) -' (x + y)) &= \varphi(x +' y) - \varphi(x + y) = \varphi(x) + \varphi(y) - f(x + y) \\ &= f(x) + f(y) - f(x) - f(y) = 0. \end{aligned}$$

Furthermore, we have that if  $x \in V$  and  $t \in \mathbb{F}$ ,

$$\varphi((t \cdot' x) -' (tx)) = \varphi(t \cdot' x) - \varphi(tx) = t\varphi(x) - f(tx) = tf(x) - tf(x) = 0.$$

Hence,  $\varphi$  is zero on the ideal  $\hat{D}$  that is quoted out from  $\mathbb{F}\{V\}$  in order to obtain the tensor algebra  $\mathcal{T}(V)$  (see Appendix A.1.8). This means that we have a unique map

$$\Phi: \mathcal{T}(V) \rightarrow \mathcal{A}$$

which is an  $\mathbb{F}$ -algebra homomorphism satisfying  $\Phi(v) = f(v)$  for all  $v \in V$ . Now we also have that for all  $v \in V$ ,

$$\Phi(v \otimes v - q(v)) = \Phi(v \otimes v) - q(v) = \Phi(v)\Phi(v) - q(v) = f(v)^2 - q(v) = 0,$$

and hence  $\Phi$  is zero on the ideal  $\mathcal{I}_q(V)$  that is quoted out from  $\mathcal{T}(V)$  to obtain  $\mathcal{G}(V, q)$ . But then since our definitions were again necessary we find that there exists a unique  $\mathbb{F}$ -algebra homomorphism

$$F: \mathcal{G}(V, q) \rightarrow \mathcal{A},$$

such that  $F([v]) = f(v)$  for all  $v \in V$ , which proves the theorem.  $\square$

### 2.1.1 Orthogonal bases and signatures

So far we have not made any assumptions on the dimension of  $V$ . We will come back to the infinite-dimensional case after discussing the more general Clifford algebra. Here we will familiarize ourselves with the properties of quadratic forms on finite-dimensional spaces, which will lead to a better understanding of the structure of geometric algebras. For the remainder of this section we will therefore assume that  $\dim V = n < \infty$ .

**Definition 2.3.** A basis  $\{e_1, \dots, e_n\}$  of  $(V, q)$  is said to be *orthogonal* or *canonical* if

$$q(e_i + e_j) = q(e_i) + q(e_j) \quad \forall i \neq j,$$

i.e. if  $\beta_q(e_i, e_j) = 0$  for all  $i \neq j$ . The basis is called *orthonormal* if we also have that  $q(e_i) \in \{-1, 0, 1\}$  for all  $i$ .

Orthogonal bases are essential for our understanding of geometric algebra, and the following theorem asserts their existence.

**Theorem 2.2.** *If  $\text{char } \mathbb{F} \neq 2$  then there exists an orthogonal basis of  $(V, q)$ .*

*Proof.* If  $q(x) = 0 \forall x \in V$  then every basis is orthogonal. Hence, we can assume that there exists  $x \in V$  with  $q(x) \neq 0$ . Let  $e$  be such an element and define

$$\begin{aligned} W = e^\perp &= \{x \in V : \beta_q(x, e) = 0\}, \\ \mathbb{F}e &= \{te : t \in \mathbb{F}\}. \end{aligned}$$

Since  $\beta_q(te, e) = t\beta_q(e, e)$ , it follows that  $W \cap \mathbb{F}e = 0$  and that  $\dim W < \dim V$ . Take an arbitrary  $x \in V$  and form

$$\tilde{x} = x - \frac{\beta_q(x, e)}{q(e)} e.$$

Then we have  $\tilde{x} \in W$  because

$$\beta_q(\tilde{x}, e) = \beta_q(x, e) - \frac{\beta_q(x, e)}{q(e)} \beta_q(e, e) = 0.$$

This shows the orthogonal decomposition  $V = W \oplus \mathbb{F}e$  with respect to  $q$ .

We now use induction over the dimension of  $V$ . For  $\dim V = 1$  the theorem is trivial, so let us assume that the theorem is proved for all vector spaces of strictly lower dimension than  $\dim V = n$ . But then there exists an orthogonal basis  $\{e_1, \dots, e_{n-1}\}$  of  $(W, q|_W)$ . Hence,  $\{e_1, \dots, e_{n-1}, e\}$  is an orthogonal basis of  $(V, q)$ .  $\square$

Because this rather fundamental theorem is not valid for fields of characteristic two (such as  $\mathbb{Z}_2$ ), we will always assume that  $\text{char } \mathbb{F} \neq 2$  when talking about *geometric algebra*. General fields and rings will be treated by the general Clifford algebra  $\mathcal{Cl}$ , which is introduced in the next subsection.

**Proposition 2.3.** *If  $\mathbb{F} = \mathbb{R}$  and  $q(e) > 0$  (or  $q(e) < 0$ ) for all elements  $e$  of an orthogonal basis  $E$ , then  $q(v) = 0$  implies  $v = 0$ .*

*Proof.* This becomes obvious when expanding  $v$  in the basis  $E$ .  $\square$

**Theorem 2.4** (Sylvester's Law of Inertia). *Assume that  $\mathbb{F} = \mathbb{R}$ . If  $E$  and  $F$  are two orthogonal bases of  $V$  and we set*

$$\begin{aligned} E^+ &:= \{e \in E : q(e) > 0\}, \\ E^- &:= \{e \in E : q(e) < 0\}, \\ E^0 &:= \{e \in E : q(e) = 0\}, \end{aligned}$$

and similarly for  $F^{+, -, 0}$ , then

$$\begin{aligned} |E^+| &= |F^+|, \\ |E^-| &= |F^-|, \\ \text{Span } E^0 &= \text{Span } F^0. \end{aligned}$$

*Proof.* Let  $V^\perp = \{x \in V : \beta_q(x, y) = 0 \ \forall y \in V\}$ . It is obvious that  $\text{Span } E^0 \subseteq V^\perp$  because  $E$  is an orthogonal basis. In order to show that  $V^\perp \subseteq \text{Span } E^0$ , we take an arbitrary  $x \in V^\perp$  and write

$$x = x^0 + x^+ + x^-, \quad x^{0,+, -} \in \text{Span } E^{0,+, -}.$$

Now,  $0 = \beta_q(x, x^+) = \beta_q(x^+, x^+) \Rightarrow x^+ = 0$ , and  $0 = \beta_q(x, x^-) = \beta_q(x^-, x^-) \Rightarrow x^- = 0$ , so that  $x = x^0$ , which shows that  $V^\perp = \text{Span } E^0$ . Analogously, we find that  $V^\perp = \text{Span } F^0$ .

In order to prove that  $|E^+| = |F^+|$ , we assume that  $|E^+| < |F^+|$ . But then we must have  $|E^-| > |F^-|$  in which case  $E^0 \cup F^+ \cup E^-$  has to be linearly dependent. Thus, there exist  $x^0 \in \text{Span } E^0$ ,  $y^+ \in \text{Span } F^+$ ,  $x^- \in \text{Span } E^-$ , not all zero, such that  $x^0 + y^+ + x^- = 0$ . This implies

$$\begin{aligned} \beta_q(x^0 + x^-, x^0 + x^-) &= \beta_q(-y^+, -y^+) \\ \Rightarrow \beta_q(x^-, x^-) &= \beta_q(y^+, y^+) \Rightarrow x^- = y^+ = 0 \Rightarrow x^0 = 0 \end{aligned}$$

This contradiction shows that  $|E^+| \geq |F^+|$ . Analogously, we find  $|E^+| \leq |F^+|$ , which proves the theorem.  $\square$

This means that there is a unique *signature*  $(s, t, u) := (|E^+|, |E^-|, |E^0|)$  associated to  $(V, q)$ . The space  $(V, q)$  is called *degenerate* if and only if  $E^0 \neq \emptyset$ . For the complex (and in general non-real) case we have the following simpler version:

**Theorem 2.5.** *If  $E$  and  $F$  are orthogonal bases of  $V$  with  $\mathbb{F} = \mathbb{C}$  and*

$$\begin{aligned} E^\times &:= \{e \in E : q(e) \neq 0\}, \\ E^0 &:= \{e \in E : q(e) = 0\}, \end{aligned}$$

and similarly for  $F^{\times, 0}$ , then

$$\begin{aligned} |E^\times| &= |F^\times|, \\ \text{Span } E^0 &= \text{Span } F^0. \end{aligned}$$

Furthermore, if  $E^0 = \emptyset$  (i.e.  $q$  is nondegenerate) then there exists an orthonormal basis  $\hat{E}$  with  $q(e) = 1 \ \forall e \in \hat{E}$ .

From the above follows that we can talk about *the* signature of a quadratic form or a metric tensor without ambiguity. We use the short-hand notation  $\mathbb{R}^{s,t,u}$  to denote the  $(s+t+u)$ -dimensional real vector space with a quadratic form of signature  $(s,t,u)$ , while  $\mathbb{C}^n$  is understood to be the complex  $n$ -dimensional space with a nondegenerate quadratic form. When  $u=0$  or  $t=u=0$  we may simply write  $\mathbb{R}^{s,t}$  or  $\mathbb{R}^s$ . A space of type  $\mathbb{R}^{n,0}$  is called *euclidean* and  $\mathbb{R}^{0,n}$  *anti-euclidean*, while the spaces  $\mathbb{R}^{1,n}$  ( $\mathbb{R}^{n,1}$ ) are called *(anti-)lorentzian*. Note that, within real and complex spaces we can always find bases that are orthonormal.

*Remark.* The general condition for orthonormal bases to exist is that the field  $\mathbb{F}$  is a so called *spin field*. This means that every  $\alpha \in \mathbb{F}$  can be written as  $\alpha = \beta^2$  or  $-\beta^2$  for some  $\beta \in \mathbb{F}$ . The fields  $\mathbb{R}$ ,  $\mathbb{C}$  and  $\mathbb{Z}_p$  for  $p$  a prime with  $p \equiv 3 \pmod{4}$  (Exercise 2.3), are spin, but e.g.  $\mathbb{Q}$  is not.

Consider now the geometric algebra  $\mathcal{G}$  over a real or complex space  $V$ . If we pick an orthonormal basis  $E = \{e_1, \dots, e_n\}$  of  $V$  then it follows from Definition 2.2 and Equation (2.2) that  $\mathcal{G}$  is the free associative algebra generated by  $E$  modulo the relations

$$e_i^2 = q(e_i) \in \{-1, 0, 1\} \quad \text{and} \quad e_i e_j = -e_j e_i, \quad i \neq j. \quad (2.4)$$

**Example 2.3.** Consider the case  $V = \mathbb{R}^2$ , i.e. the euclidean plane. The corresponding geometric algebra  $\mathcal{G}(\mathbb{R}^2)$ , also called the *plane algebra*, is generated by an orthonormal basis  $\{e_1, e_2\}$ , with  $e_1^2 = e_2^2 = 1$  and  $e_1 e_2 = -e_2 e_1$ . Starting with the unit 1 and multiplying from the left or right with  $e_1$  and  $e_2$  we see that the process stops at  $e_1 e_2$  and it follows that  $\mathcal{G}(\mathbb{R}^2)$  is spanned by four elements:

$$1, e_1, e_2, e_1 e_2.$$

Similarly, (as seen already in the introduction) the *space algebra*  $\mathcal{G}(\mathbb{R}^3)$  is spanned by eight elements,

$$1, e_1, e_2, e_3, e_1 e_2, e_2 e_3, e_1 e_3, e_1 e_2 e_3,$$

generated by an orthonormal basis  $\{e_1, e_2, e_3\}$  of vectors which are mutually anticommuting and squaring to 1.

In the same manner we conclude that a general geometric algebra  $\mathcal{G}(V)$  is spanned by  $2^n$  elements of the form

$$\{E_{i_1 i_2 \dots i_k}\}_{1 \leq i_1 < i_2 < \dots < i_k \leq n}, \quad \text{with} \quad E_{i_1 i_2 \dots i_k} := e_{i_1} e_{i_2} \dots e_{i_k}, \quad (2.5)$$

and it will follow from the construction of  $\mathcal{Cl}$  in the next subsection that there cannot be any linear relations among these elements, so that (2.5) is actually a basis of  $\mathcal{G}$ .

*Remark.* In the case that  $q=0$ , the resulting algebra of anticommuting, nilpotent elements is called a *Grassmann algebra*. When associated to a vector space  $V$  it is also commonly referred to as the *exterior algebra* of  $V$  and is often denoted  $\wedge^* V$ , and the multiplication by  $\wedge$  (wedge). Note that  $\mathcal{G}(V, q) \cong \wedge^* V$  holds as an  $\mathbb{F}$ -algebra isomorphism only when  $q=0$ , but remains as an isomorphism of (graded) vector spaces also for non-trivial quadratic forms  $q$ .



### 2.1.2 The pseudoscalar

One element in  $\mathcal{G}$  deserves special attention, namely the so-called *pseudoscalar* (sometimes called *the (oriented) volume element*) formed by the product of all elements of an orthonormal basis,

$$I := e_1 e_2 \dots e_n. \quad (2.6)$$

Note that this definition is basis independent up to orientation when  $q$  is non-degenerate. Indeed, let  $\{Re_1, \dots, Re_n\}$  be another orthonormal basis with the same orientation, where  $R \in \text{SO}(V, q)$ , the group of linear orientation-preserving transformations which leave  $q$  invariant<sup>5</sup>. Then, due to the anticommutativity of the  $e_i$ 's and  $Re_i$ 's, and hence orthogonality of the rows and columns of the matrix of  $R$ ,<sup>6</sup>

$$\begin{aligned} Re_1 Re_2 \dots Re_n &= \sum_{j_1, \dots, j_n} R_{j_1 1} R_{j_2 2} \dots R_{j_n n} e_{j_1} e_{j_2} \dots e_{j_n} \\ &= \sum_{\pi \in S_n} \text{sign}(\pi) R_{1\pi(1)} R_{2\pi(2)} \dots R_{n\pi(n)} e_1 e_2 \dots e_n \\ &= \det R e_1 e_2 \dots e_n = I, \end{aligned}$$

where  $S_n$  denotes the symmetric group of order  $n$ , i.e. all permutations of  $n$  elements. Note that, by selecting a certain pseudoscalar for  $\mathcal{G}$  we also impose a certain orientation on  $V$ . There is no such thing as an absolute orientation; instead all statements concerning orientation will be made relative to the chosen one.

The square of the pseudoscalar is given by (and gives information about) the signature and dimension of  $(V, q)$ . For  $\mathcal{G}(\mathbb{R}^{s,t,u})$  we have that

$$I^2 = (-1)^{\frac{1}{2}n(n-1)+t} \delta_{u,0}, \quad \text{where } n = s + t + u. \quad (2.7)$$

We say that  $\mathcal{G}$  is *degenerate* if the quadratic form is degenerate, or equivalently if  $I^2 = 0$ . For odd  $n$ ,  $I$  commutes with all elements in  $\mathcal{G}$ , while for even  $n$ ,  $I$  anticommutes with all vectors  $v \in V$  (Exercise 2.6).

**Example 2.4.** The pseudoscalar in  $\mathcal{G}(\mathbb{R}^2)$  is  $I = e_1 e_2$ . Here,  $I^2 = -1$ , so that the subalgebra spanned by  $\{1, I\}$  is isomorphic to  $\mathbb{C}$ . Also, any vector  $v \in \mathbb{R}^2$  can be written

$$v = ae_1 + be_2 = e_1(a + bI) = |v|e_1 e^{\varphi I},$$

for some  $a, b, \varphi \in \mathbb{R}$ , which is analogous to the polar form of a complex number.

**Example 2.5.** In  $\mathcal{G}(\mathbb{R}^3)$  the pseudoscalar is given by  $I = e_1 e_2 e_3$ . Again,  $I^2 = -1$  and  $\{1, I\}$  spans a subalgebra isomorphic to the complex numbers. However, there are infinitely many other such subalgebras embedded in  $\mathcal{G}$  since for every choice of  $e = \alpha e_1 + \beta e_2 + \gamma e_3$  on the unit sphere  $\alpha^2 + \beta^2 + \gamma^2 = 1$ , the element

$$eI = \alpha e_2 e_3 + \beta e_3 e_1 + \gamma e_1 e_2 \quad (2.8)$$

also satisfies  $(eI)^2 = eIeI = e^2 I^2 = -1$ .

<sup>5</sup>Such transformations will be introduced and discussed in Section 6.

<sup>6</sup>For this identity it might be helpful to switch to the exterior product introduced in the next subsection (where the concept of pseudoscalar also extends in greater generality).

**Example 2.6.** Let us also consider an example with non-euclidean signature. The simplest case is the anti-euclidean line, i.e.  $\mathbb{R}^{0,1}$ . A unit basis element (also pseudoscalar)  $i \in \mathbb{R}^{0,1}$  satisfies  $i^2 = -1$ , and

$$\mathcal{G}(\mathbb{R}^{0,1}) = \text{Span}_{\mathbb{R}}\{1, i\} = \mathbb{C}.$$

**Example 2.7.** An even more interesting case is the anti-euclidean plane,  $\mathbb{R}^{0,2}$ . Let  $\{i, j\}$  denote an orthonormal basis, i.e.  $i^2 = j^2 = -1$  and  $ij = -ji$ . Denoting the pseudoscalar by  $k$ , we have  $k = ij$  and

$$i^2 = j^2 = k^2 = ijk = -1,$$

which are the defining relations of the *quaternion algebra*  $\mathbb{H}$ . It follows that  $\mathcal{G}(\mathbb{R}^{0,2})$  and  $\mathbb{H}$  are isomorphic as  $\mathbb{R}$ -algebras.

**Exercise 2.1.** Prove the Jordan-von Neumann theorem (see Example 2.2). *Hint:* First prove additivity and linearity over  $-1$ , then linearity over  $\mathbb{Z}$ ,  $\mathbb{Q}$ , and finally  $\mathbb{R}$ .

**Exercise 2.2.** Prove Theorem 2.5.

**Exercise 2.3.** Show that  $\mathbb{Z}_p$  is a spin field when  $p$  is a prime such that  $p \equiv 3 \pmod{4}$ .

**Exercise 2.4.** Verify formula (2.7) for the square of the pseudoscalar.

**Exercise 2.5.** Find an  $\mathbb{R}$ -algebra isomorphism

$$\mathcal{G}(\mathbb{R}^2) \rightarrow \mathbb{R}^{2 \times 2} = \{2 \times 2 \text{ matrices with entries in } \mathbb{R}\}.$$

*Hint:* Find (preferably very simple)  $2 \times 2$  matrices  $\gamma_1, \gamma_2$  which anti-commute and satisfy  $\gamma_1^2 = \gamma_2^2 = 1_{2 \times 2}$ .

**Exercise 2.6.** The *center*  $Z(\mathcal{A})$  of an algebra  $\mathcal{A}$  consists of those elements which commute with all of  $\mathcal{A}$ . Show that, for odd-dimensional  $V$  (or  $q = 0$ ), the center of  $\mathcal{G}(V)$  is  $Z(\mathcal{G}) = \text{Span}_{\mathbb{F}}\{1, I\}$ , while for even dimensions (and  $q \neq 0$ ), the center consists of the scalars  $\mathbb{F}$  only.

**Exercise 2.7.** Assume (as usual) that  $\dim V < \infty$  and let  $|\cdot|$  be any norm on  $\mathcal{G}(V)$ . Prove that  $e^x := \sum_{k=0}^{\infty} \frac{x^k}{k!}$  converges and that  $e^x e^{-x} = e^{-x} e^x = 1$  for all  $x \in \mathcal{G}$ .

## 2.2 Combinatorial Clifford algebra $\mathcal{Cl}(X, R, r)$

We now take a temporary step away from the comfort of fields and vector spaces and instead consider the purely algebraic features of geometric algebra that were uncovered in the previous subsection.

Note that we could, for an  $n$ -dimensional space  $V$  over  $\mathbb{F}$ , roughly write

$$\mathcal{G}(V) = \text{Span}_{\mathbb{F}}\{E_A\}_{A \subseteq \{1, 2, \dots, n\}}, \quad (2.9)$$

with  $E_A$  defined in (2.5), and that the geometric product of these basis elements behaves as

$$E_A E_B = \tau(A, B) E_{A \Delta B}, \quad \text{where } \tau(A, B) \in \mathbb{F} \text{ (usually 1, -1, or 0)}, \quad (2.10)$$

and  $A \Delta B := (A \cup B) \setminus (A \cap B)$  denotes the *symmetric difference* between the sets  $A$  and  $B$ . Motivated by these expressions we consider the following generalization.

**Definition 2.4.** Let  $X$  be any finite set and  $R$  a commutative ring with unit. Let  $r: X \rightarrow R$  be an arbitrary function which is to be thought of as a *signature* on  $X$ . The *Clifford algebra* over  $(X, R, r)$  is defined as the set<sup>7</sup>

$$\mathcal{Cl}(X, R, r) := \bigoplus_{\mathcal{P}(X)} R,$$

i.e. the free  $R$ -module generated by  $\mathcal{P}(X)$ , the set of all subsets of  $X$ . We may use the shorter notation  $\mathcal{Cl}(X)$ , or just  $\mathcal{Cl}$ , when the current choice of  $X$ ,  $R$  and  $r$  is clear from the context. Furthermore, we call  $R$  the *scalars* of  $\mathcal{Cl}$ .

**Example 2.8.** A typical element of  $\mathcal{Cl}(\{x, y, z\}, \mathbb{Z}, r)$ , i.e. the Clifford algebra over a set of three elements with integer scalars, could for example look like

$$5\emptyset + 3\{x\} + 2\{y\} - \{x, y\} + 12\{x, y, z\}. \quad (2.11)$$

We have not yet defined a multiplication on  $\mathcal{Cl}$ . In addition to being  $R$ -bilinear and associative, we would like the product to satisfy  $\{x\}^2 = r(x)\emptyset$  for  $x \in X$ ,  $\{x\}\{y\} = -\{y\}\{x\}$  for  $x \neq y \in X$  and  $\emptyset A = A\emptyset = A$  for all subsets  $A \in \mathcal{P}(X)$ . In order to arrive at such a product we make use of the following

**Lemma 2.6.** *Given  $X$  a finite set and  $r: X \rightarrow R$ , there exists a map  $\tau: \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow R$  such that*

- i)  $\tau(\{x\}, \{x\}) = r(x) \quad \forall x \in X$ ,
- ii)  $\tau(\{x\}, \{y\}) = -\tau(\{y\}, \{x\}) \quad \forall x, y \in X : x \neq y$ ,
- iii)  $\tau(\emptyset, A) = \tau(A, \emptyset) = 1 \quad \forall A \in \mathcal{P}(X)$ ,
- iv)  $\tau(A, B)\tau(A \Delta B, C) = \tau(A, B \Delta C)\tau(B, C) \quad \forall A, B, C \in \mathcal{P}(X)$ ,
- v)  $\tau(A, B) \in \{-1, 1\} \quad \text{if } A \cap B = \emptyset$ .

*Proof.* We proceed by induction on the cardinality  $|X|$  of  $X$ . For  $X = \emptyset$  the lemma is trivial, so let  $z \in X$  and assume the lemma holds for  $Y := X \setminus \{z\}$ . Hence, there is a  $\tau': \mathcal{P}(Y) \times \mathcal{P}(Y) \rightarrow R$  which has the properties (i)-(v) above. If  $A \subseteq Y$  we write  $A' = A \cup \{z\}$  and, for  $A, B$  in  $\mathcal{P}(Y)$  we extend  $\tau'$  to  $\tau: \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow R$  in the following way:

$$\begin{aligned} \tau(A, B) &:= \tau'(A, B) \\ \tau(A', B) &:= (-1)^{|B|} \tau'(A, B) \\ \tau(A, B') &:= \tau'(A, B) \\ \tau(A', B') &:= r(z)(-1)^{|B|} \tau'(A, B) \end{aligned}$$

Now it is straightforward to verify that (i)-(v) holds for  $\tau$ , which completes the proof.  $\square$

<sup>7</sup>Again, see Appendix A.1 if the notation is unfamiliar.

**Definition 2.5.** Define the *Clifford product*

$$\begin{aligned} Cl(X) \times Cl(X) &\rightarrow Cl(X) \\ (A, B) &\mapsto AB \end{aligned}$$

by taking  $AB := \tau(A, B)A \triangle B$  for  $A, B \in \mathcal{P}(X)$  and extending linearly. We choose to use the  $\tau$  which is constructed as in the proof of Lemma 2.6 by consecutively adding elements from the set  $X$ . A unique such  $\tau$  may only be selected after imposing a certain order (orientation) on the set  $X$ .

Using Lemma 2.6 one easily verifies that this product has all the properties that we asked for above. For example, in order to verify associativity we note that

$$A(BC) = A(\tau(B, C)B \triangle C) = \tau(A, B \triangle C)\tau(B, C)A \triangle (B \triangle C), \quad (2.12)$$

while

$$(AB)C = \tau(A, B)(A \triangle B)C = \tau(A, B)\tau(A \triangle B, C)(A \triangle B) \triangle C. \quad (2.13)$$

Associativity now follows from (iv) and the associativity of the symmetric difference (Exercise 2.8). As is expected from the analogy with  $\mathcal{G}$ , we also have the property that different basis elements of  $Cl$  commute up to a sign:

**Proposition 2.7.** *If  $A, B \in \mathcal{P}(X)$  then*

$$AB = (-1)^{\frac{1}{2}|A|(|A|-1) + \frac{1}{2}|B|(|B|-1) + \frac{1}{2}|A \triangle B|(|A \triangle B|-1)} BA.$$

*Proof.* By the property (v) in Lemma 2.6 it is sufficient to prove this for  $A = \{a_1\}\{a_2\} \dots \{a_k\}$ ,  $B = \{b_1\}\{b_2\} \dots \{b_l\}$ , where  $a_i$  are disjoint elements in  $X$  and likewise for  $b_i$ . If  $A$  and  $B$  have  $m$  elements in common then

$$AB = (-1)^{(k-m)l + m(l-1)} BA = (-1)^{kl-m} BA$$

by property (ii). But then we are done, since

$$\frac{1}{2}(-k(k-1) - l(l-1) + (k+l-2m)(k+l-2m-1)) \equiv kl + m \pmod{2}.$$

□

### 2.2.1 Equivalence of $\mathcal{G}$ and $Cl$

We are now ready to make the formal connection between  $\mathcal{G}$  and  $Cl$ . We work in the setting where both algebras have been defined and where orthogonal bases exist, i.e. finite-dimensional vector spaces over fields  $\mathbb{F}$  with  $\text{char } \mathbb{F} \neq 2$ . The idea is then to pick an orthogonal basis  $E = \{e_1, \dots, e_n\}$  of  $V$  and consider the Clifford algebra  $Cl(E, \mathbb{F}, q|_E)$ . Defining a canonical embedding  $i: V \rightarrow Cl$  by  $i(e_j) := \{e_j\}$  for  $j = 1, \dots, n$  and linear extension, we then observe by the properties given above that

$$\begin{aligned} i(v)^2 &= i(\sum_j v_j e_j) i(\sum_k v_k e_k) = \sum_{j,k} v_j v_k i(e_j) i(e_k) \\ &= \sum_{j,k} v_j v_k \{e_j\} \{e_k\} = \sum_j v_j^2 \{e_j\}^2 \\ &= \sum_j v_j^2 q|_E(e_j) \emptyset = \sum_j v_j^2 q(e_j) \emptyset = q(\sum_j v_j e_j) \emptyset = q(v) \emptyset. \end{aligned}$$

In Theorem 2.1 we noted that  $\mathcal{G}(V, q)$  solved a certain universality problem. We will now show that  $\mathcal{Cl}(E, \mathbb{F}, q|_E)$  has the same universality property, with  $E$  any choice of orthogonal basis for  $(V, q)$ .

**Theorem 2.8** (Universality of  $\mathcal{Cl}$ ). *Let  $V \xrightarrow{q} \mathbb{F}$  be a finite-dimensional quadratic space over the field  $\mathbb{F}$  with  $\text{char } \mathbb{F} \neq 2$ , and let  $\mathcal{A}$  be an associative algebra over  $\mathbb{F}$  with a unit denoted by  $1_{\mathcal{A}}$ . Moreover, let  $f: V \rightarrow \mathcal{A}$  be linear and such that  $f(v)^2 = q(v)1_{\mathcal{A}}$  for all  $v \in V$ . Take  $E$  to be any orthogonal basis for  $V$  and introduce  $\mathcal{Cl}(E, \mathbb{F}, q|_E)$ . Then there exists a unique  $\mathbb{F}$ -algebra homomorphism  $F: \mathcal{Cl}(E, \mathbb{F}, q|_E) \rightarrow \mathcal{A}$  such that  $F(i(v)) = f(v)$  for all  $v \in V$ .*

*Proof.* Let  $E = \{e_1, e_2, \dots, e_n\}$  be the orthogonal basis for  $(V, q)$ . For any subset  $A = \{1', 2', \dots, k'\} \subseteq \{1, 2, \dots, n\}$ , let  $e_A$  denote the element  $e_{1'}e_{2'} \dots e_{k'}$  in  $\mathcal{Cl}(E, \mathbb{F}, q|_E)$ , and put  $a_A := a_{1'}a_{2'} \dots a_{k'}$  in  $\mathcal{A}$  where  $a_i := f(e_i)$ . We also let  $e_{\emptyset} := 1$  in  $\mathcal{Cl}$  and  $a_{\emptyset} := 1_{\mathcal{A}}$ .

If  $v, w \in V$  are orthogonal, then

$$f(v+w)^2 = q(v+w)1_{\mathcal{A}} = q(v)1_{\mathcal{A}} + q(w)1_{\mathcal{A}} = f(v)^2 + f(w)^2,$$

but also

$$(f(v) + f(w))^2 = f(v)^2 + f(w)^2 + f(v)f(w) + f(w)f(v).$$

Hence  $f(v)f(w) + f(w)f(v) = 0$  and we have that

$$a_i a_j + a_j a_i = 0 \quad \forall i \neq j \quad \text{and} \quad a_i^2 = q(e_i)1_{\mathcal{A}} \quad \forall i. \quad (2.14)$$

We now define an  $\mathbb{F}$ -linear map by

$$\begin{aligned} F: \mathcal{Cl}(E, \mathbb{F}, q|_E) &\rightarrow \mathcal{A} \\ 1 = e_{\emptyset} &\mapsto a_{\emptyset} = 1_{\mathcal{A}} \\ e_A &\mapsto a_A, \end{aligned}$$

and linear extension, and we will verify that this is an  $\mathbb{F}$ -algebra homomorphism. Indeed, we have

$$F(e_A e_B) = F(\tau(A, B)e_{A \Delta B}) = \tau(A, B)a_{A \Delta B}$$

and also, by (2.14) and the properties of  $\tau$  in Lemma 2.6,

$$F(e_A)F(e_B) = a_A a_B = \tau(A, B)a_{A \Delta B},$$

so if  $x = \sum_{A \subseteq \{1, \dots, n\}} x_A e_A$  and  $y = \sum_B y_B e_B$  we therefore obtain

$$F(xy) = \sum_{A, B} x_A y_B F(e_A e_B) = \sum_{A, B} x_A y_B F(e_A)F(e_B) = F(x)F(y).$$

That  $F(x+y) = F(x) + F(y)$  is trivial. Noting that our definition of  $F$  above was necessary in order for it to be an algebra homomorphism, this proves the theorem.

We also note that the canonical map  $i: V \rightarrow \mathcal{Cl}(E, \mathbb{F}, q|_E)$  obviously is injective. The corresponding statement was not clear in the former universality theorem concerning  $\mathcal{G}(V, q)$ .  $\square$

From this universality property follows then that  $\mathcal{G}$  and  $\mathcal{Cl}$  are isomorphic.

**Corollary.** *If  $E$  is an orthonormal basis for  $(V, q)$  then  $\mathcal{Cl}(E, \mathbb{F}, q|_E)$  and  $\mathcal{G}(V, q)$  are isomorphic as  $\mathbb{F}$ -algebras, and in particular*

$$\dim_{\mathbb{F}} \mathcal{G}(V, q) = \dim_{\mathbb{F}} \mathcal{Cl}(E, \mathbb{F}, q|_E) = 2^{\dim V},$$

and the map  $j: V \rightarrow \mathcal{G}(V, q)$ ,  $v \mapsto [v]$ , is injective.

*Proof.* Consider the diagram

$$\begin{array}{ccc} & \mathcal{Cl}(E, \mathbb{F}, q|_E) \supseteq \text{id} & \\ & \nearrow i & \\ (V, q) & \varphi \downarrow \uparrow \psi & \\ & \searrow j & \\ & \mathcal{G}(V, q) \supseteq \text{id} & \end{array}$$

Since  $\mathcal{Cl}$  is universal, there exists a unique algebra homomorphism  $\alpha: \mathcal{Cl} \rightarrow \mathcal{Cl}$  such that  $\alpha \circ i = i$ , and since  $\alpha = \text{id}_{\mathcal{Cl}}$  satisfies this we know that  $\alpha$  has to be the identity map on  $\mathcal{Cl}$ . Similarly, by the universality of  $\mathcal{G}$  there is only one map  $\beta: \mathcal{G} \rightarrow \mathcal{G}$  satisfying  $\beta \circ j = j$ , so  $\beta = \text{id}_{\mathcal{G}}$ . Furthermore, there is also a unique  $\varphi: \mathcal{Cl} \rightarrow \mathcal{G}$  such that  $\varphi$  is an  $\mathbb{F}$ -algebra homomorphism and such that  $\varphi \circ i = j$ , and a unique  $\mathbb{F}$ -algebra homomorphism  $\psi: \mathcal{G} \rightarrow \mathcal{Cl}$  such that  $\psi \circ j = i$ . But since  $i$  is injective, we have then that also  $j$  must be injective. Moreover, since  $\psi \circ \varphi = \text{id}_{\mathcal{Cl}}$  and  $\varphi \circ \psi = \text{id}_{\mathcal{G}}$ , it follows that  $\varphi$  and  $\psi$  are isomorphisms and  $\psi^{-1} = \varphi$ .

The dimension of  $\mathcal{Cl}(E, \mathbb{F}, q|_E)$  is by construction  $2^{\dim V}$ .  $\square$

*Remark.* One may also formulate the above universality in the language of category theory and then use the fact that all initial objects in any category are isomorphic (see Appendix A.2).

We shall make this equivalence between  $\mathcal{G}$  and  $\mathcal{Cl}$  even more transparent by suppressing the unit  $\emptyset$  in expressions, writing simply  $e$  instead of  $\{e\}$  for singletons  $e \in E$ , and identifying the images  $\text{im } i = \text{im } j = V$ . For example, with an orthonormal basis  $\{e_1, e_2, e_3\}$  in  $\mathbb{R}^3$ , both  $\mathcal{G}$  and  $\mathcal{Cl}$  are then spanned by

$$\{1, e_1, e_2, e_3, e_1e_2, e_1e_3, e_2e_3, e_1e_2e_3\}, \quad (2.15)$$

and the two algebras can therefore be identified.

### 2.2.2 Grade structure

There is a natural grade structure on  $\mathcal{Cl}$  given by the cardinality of the subsets of  $X$ . Consider the following

**Definition 2.6.** The *subspace of  $k$ -vectors* in  $\mathcal{Cl}$ , or the *grade- $k$  part* of  $\mathcal{Cl}$ , is defined by

$$\mathcal{Cl}^k(X, R, r) := \bigoplus_{A \in \mathcal{P}(X) : |A|=k} R.$$

Of special importance are the *even* and *odd subspaces*,

$$\mathcal{Cl}^{\pm}(X, R, r) := \bigoplus_{\substack{k \text{ is} \\ \text{even} \\ \text{odd}}} \mathcal{Cl}^k(X, R, r).$$

This notation carries over to the corresponding subspaces of  $\mathcal{G}$  and we write  $\mathcal{G}^k$ ,  $\mathcal{G}^\pm$  etc. where for example  $\mathcal{G}^0 = \mathbb{F}$  are the scalars and  $\mathcal{G}^1 = V$  the vectors. The elements of  $\mathcal{G}^2$  are also called *bivectors*, while arbitrary elements of  $\mathcal{G}$  are conventionally called *multivectors*.

We then have a split of  $\mathcal{C}l$  into graded subspaces as

$$\begin{aligned} \mathcal{C}l(X) &= \mathcal{C}l^+ \oplus \mathcal{C}l^- \\ &= \mathcal{C}l^0 \oplus \mathcal{C}l^1 \oplus \mathcal{C}l^2 \oplus \dots \oplus \mathcal{C}l^{|X|}. \end{aligned} \quad (2.16)$$

Note that, under the Clifford product,  $\mathcal{C}l^\pm \cdot \mathcal{C}l^\pm \subseteq \mathcal{C}l^+$  and  $\mathcal{C}l^\pm \cdot \mathcal{C}l^\mp \subseteq \mathcal{C}l^-$  (by the properties of the symmetric difference). Hence, the even-grade elements  $\mathcal{C}l^+$  form a subalgebra of  $\mathcal{C}l$  (and we will in Section 5 see that it is in fact also isomorphic to a Clifford algebra, though over a space of one dimension less).

*Remark.* Strictly speaking, although the algebra  $\mathcal{C}l$  is  $\mathbb{Z}_2$ -graded in this way, it is not  $\mathbb{Z}$ -graded but *filtered* in the sense that

$$\left( \bigoplus_{i \leq k} \mathcal{C}l^i \right) \cdot \left( \bigoplus_{j \leq l} \mathcal{C}l^j \right) \subseteq \bigoplus_{m \leq k+l} \mathcal{C}l^m.$$

In  $\mathcal{C}l(X, R, r)$  we have the possibility of defining a unique pseudoscalar independently of the signature  $r$ , namely the set  $X$  itself. Note, however, that it can only be normalized if  $X^2 = \tau(X, X) \in R$  is invertible, which requires that  $r$  is nondegenerate (i.e. that its values in  $R$  are invertible). We will almost exclusively talk about pseudoscalars in the setting of nondegenerate vector spaces, so this will not be a problem.

**Example 2.9.** The space algebra

$$\mathcal{G}(\mathbb{R}^3) = \mathcal{G}^0 \oplus \mathcal{G}^1 \oplus \mathcal{G}^2 \oplus \mathcal{G}^3$$

can be decomposed into scalars, vectors, bivectors and trivectors (multiples of the pseudoscalar). Using the observation (2.8) for the relation between a vector and a bivector in the space algebra, we write an arbitrary multivector  $x \in \mathcal{G}(\mathbb{R}^3)$  as

$$x = \alpha + \mathbf{a} + \mathbf{b}I + \beta I,$$

with  $\alpha, \beta \in \mathbb{R}$ ,  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ . The subalgebra  $\mathcal{G}^+(\mathbb{R}^3)$  of scalars and bivectors is actually isomorphic to the quaternion algebra (see Example 2.7), as is easily verified by taking e.g.  $i := -e_1I, j := -e_2I, k := -e_3I$ .

**Exercise 2.8.** Prove that the symmetric difference  $\Delta$  is associative (for arbitrary sets — finite or infinite).

**Exercise 2.9.** Verify that conditions (i)-(v) hold for  $\tau$  constructed in the proof of Lemma 2.6.

**Exercise 2.10.** Let  $\mathcal{G} = \mathcal{G}(\mathbb{R}^3)$ ,  $C_0 = \mathcal{G}^0 \oplus \mathcal{G}^3$ , and  $C_1 = \mathcal{G}^1 \oplus \mathcal{G}^2$ . The complex numbers are as usual denoted by  $\mathbb{C}$ . Find (natural) maps

$$\begin{aligned} \alpha &: C_0 \rightarrow \mathbb{C} \\ \beta &: C_1 \rightarrow \mathbb{C}^3 \\ \gamma &: C_0 \times C_1 \rightarrow C_1 \end{aligned}$$

such that  $\alpha$  is an algebra isomorphism,  $\beta$  is a bijection, and such that the diagram below commutes

$$\begin{array}{ccccc} C_0 & \times & C_1 & \xrightarrow{\gamma} & C_1 \\ \alpha \downarrow & & \downarrow \beta & & \downarrow \beta \\ \mathbb{C} & \times & \mathbb{C}^3 & \xrightarrow{\delta} & \mathbb{C}^3 \end{array}$$

where  $\delta$  is the usual scalar multiplication on  $\mathbb{C}^3$ , i.e.

$$\delta(z, (z_1, z_2, z_3)) = (zz_1, zz_2, zz_3).$$

*Remark.* Hence, we can think of  $\mathbb{C}^3$  (as a complex vector space) as  $C_1$  interpreted as a vector space over the field  $C_0$ .

**Exercise 2.11.** Verify that there is an  $\mathbb{R}$ -algebra isomorphism  $\varphi : \mathcal{G}(\mathbb{R}^4) \rightarrow \mathbb{H}^{2 \times 2}$  such that

$$\varphi(e_1) = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \varphi(e_2) = \begin{bmatrix} 0 & -j \\ j & 0 \end{bmatrix}, \quad \varphi(e_3) = \begin{bmatrix} 0 & -k \\ k & 0 \end{bmatrix}, \quad \varphi(e_4) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

## 2.3 Standard operations

A key feature of Clifford algebras is that they contain a surprisingly large amount of structure. In order to conveniently access the power of this structure we need to introduce some powerful notation. Most of the following definitions will be made on  $\mathcal{Cl}$  for simplicity, but because of the equivalence between  $\mathcal{G}$  and  $\mathcal{Cl}$  they carry over to  $\mathcal{G}$  in a straightforward manner. The amount of new notation might seem heavy, and we therefore recommend the reader not to get stuck trying to learn everything at once, but to come back to this subsection later as a reference when the notation is needed.

### 2.3.1 Products and involutions

We will find it convenient to introduce the notation that for any proposition  $P$ ,  $\langle P \rangle$  will denote the number 1 if  $P$  is true and 0 if  $P$  is false.

**Definition 2.7.** For  $A, B \in \mathcal{P}(X)$  we define

$$\begin{aligned} A \wedge B &:= (A \cap B = \emptyset) AB && \text{outer/exterior/wedge product} \\ A \lrcorner B &:= (A \subseteq B) AB && \text{left inner/interior product} \\ A \rceil B &:= (A \supseteq B) AB && \text{right inner/interior product} \\ A * B &:= (A = B) AB && \text{scalar product} \\ \langle A \rangle_k &:= (|A| = k) A && \text{projection on grade } k \\ A^* &:= (-1)^{|A|} A && \text{grade involution} \\ A^\dagger &:= (-1)^{\binom{|A|}{2}} A && \text{reversion} \end{aligned}$$

and extend linearly to  $\mathcal{Cl}(X, R, r)$ .

The grade involution is also called the (*first*) *main involution*. It has the property

$$(xy)^* = x^*y^*, \quad v^* = -v \tag{2.17}$$



for all  $x, y \in Cl(X)$  and  $v \in Cl^1(X)$ , as is easily verified by expanding in linear combinations of elements in  $\mathcal{P}(X)$  and using that  $|A \triangle B| \equiv |A| + |B| \pmod{2}$ . The reversion earns its name from the property

$$(xy)^\dagger = y^\dagger x^\dagger, \quad v^\dagger = v, \quad (2.18)$$

and it is sometimes called the *second main involution* or the *principal antiautomorphism*. This reversing behavior follows directly from Proposition 2.7. We will find it convenient to have a name for the composition of these two involutions. Hence, we define the *Clifford conjugate*  $x^\square$  of  $x \in Cl(X)$  by  $x^\square := x^{\star\dagger}$  and observe the property

$$(xy)^\square = y^\square x^\square, \quad v^\square = -v. \quad (2.19)$$

Note that all the above involutions act by changing sign on some of the graded subspaces. We summarize the action of these involutions in Table 2.1. Note the periodicity.

	$Cl^0$	$Cl^1$	$Cl^2$	$Cl^3$	$Cl^4$	$Cl^5$	$Cl^6$	$Cl^7$
$\star$	+	-	+	-	+	-	+	-
$\dagger$	+	+	-	-	+	+	-	-
$\square$	+	-	-	+	+	-	-	+

Table 2.1: The action of involutions on graded subspaces of  $Cl$ .

**Example 2.10.** If  $x \in \mathcal{G}(\mathbb{R}^2)$  and  $x^\star = -x$ , then  $x$  has to be a vector since these are the only odd elements in this case. Similarly, we see that if  $x$  is a multivector in  $\mathcal{G}(\mathbb{R}^3)$  then the expression  $x^\dagger x$  cannot contain any bivector or trivector parts since the expression is self-reversing, i.e.  $(x^\dagger x)^\dagger = x^\dagger x^{\dagger\dagger} = x^\dagger x$ .

**Example 2.11.** In the case  $\mathcal{G}(\mathbb{R}^{0,1}) \cong \mathbb{C}$  we see that the grade involution (or Clifford conjugate) corresponds to the complex conjugate. On the other hand, in the quaternion algebra  $\mathcal{G}(\mathbb{R}^{0,2}) \cong \mathbb{H}$  (see Example 2.7), the Clifford conjugate corresponds to the quaternion conjugate since it changes sign on all non-scalar (i.e. imaginary) grades,

$$(\alpha + \beta i + \gamma j + \delta k)^\square = \alpha - \beta i - \gamma j - \delta k.$$

The scalar product has the symmetric property  $x \star y = y \star x$  for all  $x, y \in Cl$ . Therefore, it forms a symmetric bilinear map  $Cl \times Cl \rightarrow R$  which is seen to be degenerate if and only if  $Cl$  (i.e. the signature  $r$ ) is degenerate. Note that this map coincides with the bilinear form  $\beta_q$  when restricted to  $V = \mathcal{G}^1(V, q)$ , and that subspaces of different grade are orthogonal with respect to the scalar product.

The following identities relating the inner, outer, and scalar products<sup>8</sup> will turn out to be extremely useful:

<sup>8</sup>Another product that is often seen in the context of geometric algebra is the *inner product*, defined by

$$A \bullet B := (A \subseteq B \text{ or } A \supseteq B) AB = A \lrcorner B + A \jmath B - A \star B.$$

We will stick to the left and right inner products, however, because they admit a simpler handling of grades. For example, the corresponding identities in Proposition 2.9 would need to be supplied with grade restrictions. Also beware that the meaning of the symbols  $\lrcorner$  and  $\jmath$  is sometimes reversed in the literature.

**Proposition 2.9.** For all  $x, y, z \in Cl(X)$  we have

$$\begin{aligned}x \wedge (y \wedge z) &= (x \wedge y) \wedge z, \\x \lrcorner (y \lrcorner z) &= (x \lrcorner y) \lrcorner z, \\x \lrcorner (y \wedge z) &= (x \wedge y) \lrcorner z, \\x * (y \lrcorner z) &= (x \wedge y) * z,\end{aligned}$$

and

$$1 \wedge x = x \wedge 1 = 1 \lrcorner x = x \lrcorner 1 = x.$$

*Proof.* This follows directly from Definition 2.7 and basic set logic. For example, taking  $A, B, C \in \mathcal{P}(X)$  we have (consider drawing a Venn diagram)

$$\begin{aligned}A \lrcorner (B \lrcorner C) &= (B \subseteq C)(A \subseteq B \Delta C)ABC \\&= (B \subseteq C \text{ and } A \subseteq C \setminus B)ABC \\&= (A \cap B = \emptyset \text{ and } A \cup B \subseteq C)ABC \\&= (A \cap B = \emptyset)(A \Delta B \subseteq C)ABC \\&= (A \wedge B) \lrcorner C.\end{aligned}$$

The other identities are proven in an equally simple way.  $\square$

Note that the first identity in the above proposition states that the wedge product  $\wedge$  is associative. It is not difficult to see that the algebra  $(\mathcal{G}, \wedge)$  with this product is isomorphic (as graded algebras) to the exterior algebra  $\wedge^* V$ .

In order to be able to work efficiently with Clifford algebras it is crucial to understand how vectors behave under these operations.

**Proposition 2.10.** For all  $x, y \in Cl(X)$  and  $v \in Cl^1(X)$  we have

$$\begin{aligned}vx &= v \lrcorner x + v \wedge x, \\v \lrcorner x &= \frac{1}{2}(vx - x^*v) = -x^* \lrcorner v, \\v \wedge x &= \frac{1}{2}(vx + x^*v) = x^* \wedge v, \\v \lrcorner (xy) &= (v \lrcorner x)y + x^*(v \lrcorner y).\end{aligned}$$

The first three identities are shown simply by using linearity and set relations, while the fourth follows immediately from the second.

**Example 2.12.** For 1-vectors  $u, v \in Cl^1$  we have the basic relations

$$uv = \langle uv \rangle_0 + \langle uv \rangle_2 = u * v + u \wedge v, \quad (2.20)$$

$$u \lrcorner v = v \lrcorner u = u \lrcorner v = u * v = \frac{1}{2}(uv + vu), \quad (2.21)$$

and

$$u \wedge v = -v \wedge u = \frac{1}{2}(uv - vu), \quad (2.22)$$

while for bivectors  $B \in Cl^2$  and  $v \in Cl^1$  we have

$$vB = \langle vB \rangle_1 + \langle vB \rangle_3 = v \lrcorner B + v \wedge B, \quad (2.23)$$

$$v \lrcorner B = \frac{1}{2}(vB - Bv) = -B \lrcorner v, \quad (2.24)$$

$$v \wedge B = \frac{1}{2}(vB + Bv) = B \wedge v, \quad (2.25)$$

and  $v \lrcorner B = B \lrcorner v = v * B = 0$ . In the special case that  $B = u \wedge w$  is an outer product of two vectors  $u, w$  (i.e. a 2-blade), we find from taking the vector part of the the last identity in Proposition 2.10 the useful expansion

$$v \lrcorner (u \wedge w) = (v * u)w - (v * w)u. \quad (2.26)$$

This expansion is generalized in Exercise 2.19.

It is sometimes useful to expand the various products and involutions in terms of the grades involved. The following identities are left as exercises.

**Proposition 2.11.** *For all  $x, y \in Cl(X)$  we have*

$$\begin{aligned} x \wedge y &= \sum_{n, m \geq 0} \langle \langle x \rangle_n \langle y \rangle_m \rangle_{n+m}, \\ x \lrcorner y &= \sum_{0 \leq n \leq m} \langle \langle x \rangle_n \langle y \rangle_m \rangle_{m-n}, \\ x \lrcorner y &= \sum_{n \geq m \geq 0} \langle \langle x \rangle_n \langle y \rangle_m \rangle_{n-m}, \\ x \bullet y &= \sum_{n, m \geq 0} \langle \langle x \rangle_n \langle y \rangle_m \rangle_{|n-m|}, \\ x * y &= \langle xy \rangle_0, \\ x^* &= \sum_{n \geq 0} (-1)^n \langle x \rangle_n, \\ x^\dagger &= \sum_{n \geq 0} (-1)^{\binom{n}{2}} \langle x \rangle_n. \end{aligned}$$

### 2.3.2 Linear independence

In the general setting of a Clifford algebra with scalars in a ring  $R$ , we need to be careful about the notion of linear (in-)dependence. A subset  $\{x_1, x_2, \dots, x_m\}$  (with distinct elements  $x_j$ ) of  $Cl$  is called *linearly dependent* if there exist  $r_1, \dots, r_m \in R$ , not all zero, such that

$$r_1 x_1 + r_2 x_2 + \dots + r_m x_m = 0. \quad (2.27)$$

Note that a single nonzero 1-vector could be linearly dependent in this context (e.g. with  $R = \mathbb{Z}_4$ ,  $x_1 = 2e_1$  and  $r_1 = 2$ ). We will prove an important theorem concerning linear dependence where we use the following

**Lemma 2.12.** *If  $u_1, u_2, \dots, u_k$  and  $v_1, v_2, \dots, v_k$  are 1-vectors then*

$$(u_1 \wedge u_2 \wedge \dots \wedge u_k) * (v_k \wedge v_{k-1} \wedge \dots \wedge v_1) = \det \begin{bmatrix} u_1 * v_1 & \dots & u_1 * v_k \\ \vdots & & \vdots \\ u_k * v_1 & \dots & u_k * v_k \end{bmatrix}.$$

*Proof.* Since both sides of the expression are multilinear and alternating in both the  $u_i$ :s and the  $v_i$ :s, we need only consider ordered disjoint elements  $\{e_i\}$  in the basis of singleton sets in  $X$ . Both sides are zero, except in the case

$$\begin{aligned} (e_{i_1} e_{i_2} \dots e_{i_k}) * (e_{i_k} e_{i_{k-1}} \dots e_{i_1}) &= \\ &= r(e_{i_1}) r(e_{i_2}) \dots r(e_{i_k}) = \det [r(e_{i_p}) \delta_{p,q}]_{1 \leq p, q \leq k} \\ &= \det [e_{i_p} * e_{i_q}]_{1 \leq p, q \leq k}, \end{aligned}$$

so we are done.  $\square$

**Theorem 2.13.** *The 1-vectors  $\{x_1, x_2, \dots, x_m\}$  are linearly independent if and only if the  $m$ -vector  $\{x_1 \wedge x_2 \wedge \dots \wedge x_m\}$  is linearly independent.*

*\*Proof.* Assume that  $r_1x_1 + \dots + r_mx_m = 0$ , where, say,  $r_1 \neq 0$ . Then

$$\begin{aligned} r_1(x_1 \wedge \dots \wedge x_m) &= (r_1x_1) \wedge x_2 \wedge \dots \wedge x_m \\ &= (r_1x_1 + \dots + r_mx_m) \wedge x_2 \wedge \dots \wedge x_m = 0, \end{aligned}$$

since  $x_i \wedge x_i = 0$ .

Conversely, assume that  $rX = 0$  for  $r \neq 0$  in  $R$  and  $X = x_1 \wedge \dots \wedge x_m$ . We will use the basis minor theorem for arbitrary rings which can be found in the appendix (see Appendix A.5). Assume that  $x_j = x_{1j}e_1 + \dots + x_{nj}e_n$ , where  $x_{ij} \in R$  and  $e_i \in X$  are basis elements such that  $e_i^2 = 1$ . This assumption on the signature is no loss in generality, since this theorem only concerns the exterior algebra associated to the outer product. It will only serve to simplify our reasoning below. Collect the coordinates in a matrix

$$A := \begin{bmatrix} rx_{11} & x_{12} & \cdots & x_{1m} \\ rx_{21} & x_{22} & \cdots & x_{2m} \\ \vdots & \vdots & & \vdots \\ rx_{n1} & x_{n2} & \cdots & x_{nm} \end{bmatrix} \in R^{n \times m}, \quad m \leq n$$

and note that we can expand  $rX$  in a grade- $m$  basis as

$$rX = \sum_{E \subseteq X: |E|=m} (rX * E^\dagger)E = \sum_{E \subseteq X: |E|=m} (\det A_{E, \{1, \dots, m\}})E,$$

where we used Lemma 2.12. We find that the determinant of each  $m \times m$  minor of  $A$  is zero.

Now, let  $k$  be the rank of  $A$ . Then we must have  $k < m$ , and if  $k = 0$  then  $rx_{i1} = 0$  and  $x_{ij} = 0$  for all  $i = 1, \dots, n$ ,  $j > 1$ . But that would mean that  $\{x_1, \dots, x_m\}$  are linearly dependent. Therefore we assume that  $k > 0$  and, without loss of generality, that

$$d := \det \begin{bmatrix} rx_{11} & x_{12} & \cdots & x_{1k} \\ \vdots & \vdots & & \vdots \\ rx_{k1} & x_{k2} & \cdots & x_{kk} \end{bmatrix} \neq 0.$$

By Theorem A.5 (Basis minor) there exist  $r_1, \dots, r_k \in R$  such that

$$r_1rx_1 + r_2x_2 + \dots + r_kx_k + dx_m = 0.$$

Hence,  $\{x_1, \dots, x_m\}$  are linearly dependent.  $\square$

### 2.3.3 Duality

For our final set of operations, we will consider a nondegenerate geometric algebra  $\mathcal{G}$  with pseudoscalar  $I$ . The nondegeneracy implies that there exists a natural duality between the inner and outer products.

**Definition 2.8.** We define the *dual* of  $x \in \mathcal{G}$  by  $x^c := xI^{-1}$ . Furthermore, the *dual outer product* or *meet*, denoted  $\vee$ , is defined such that the diagram

$$\begin{array}{ccc} \mathcal{G} \times \mathcal{G} & \xrightarrow{\vee} & \mathcal{G} \\ (\cdot)^c \downarrow & & \downarrow (\cdot)^c \\ \mathcal{G} \times \mathcal{G} & \xrightarrow{\wedge} & \mathcal{G} \end{array}$$

commutes, i.e.  $(x \vee y)^c := x^c \wedge y^c$ , implying  $x \vee y = ((xI^{-1}) \wedge (yI^{-1}))I$ .

*Remark.* In  $\mathcal{Cl}(X)$ , the corresponding dual of  $A \in \mathcal{P}(X)$  is

$$A^c = AX^{-1} = \tau(X, X)^{-1} \tau(A, X) A \Delta X \propto A^c,$$

the complement of the set  $A$ . Hence, we actually find that the dual is a linearized version of a sign (or orientation) -respecting complement. This motivates our choice of notation.

**Proposition 2.14.** For all  $x, y \in \mathcal{G}$  we have

$$\begin{aligned} x \lrcorner y^c &= (x \wedge y)^c, \\ x \wedge y^c &= (x \lrcorner y)^c, \end{aligned} \tag{2.28}$$

and if  $x \in \mathcal{G}^k$  then  $x^c \in \mathcal{G}^{\dim V - k}$ .

*Proof.* Using Proposition 2.9 and the fact that  $xI = x \lrcorner I \forall x$ , we obtain

$$x \lrcorner (yI^{-1}) = x \lrcorner (y \lrcorner I^{-1}) = (x \wedge y) \lrcorner I^{-1} = (x \wedge y)I^{-1},$$

and from this follows also the second identity

$$(x \wedge y^c)I^{-1}I = (x \lrcorner y^{cc})I = (x \lrcorner y)I^{-2}I.$$

Lastly, the grade statement is obvious from the above definition and remark.  $\square$

**Example 2.13.** The cross product in  $\mathbb{R}^3$  can be defined as the dual of the outer product,

$$\mathbf{a} \times \mathbf{b} := (\mathbf{a} \wedge \mathbf{b})^c, \tag{2.29}$$

for  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ . From Proposition 2.14 and e.g. (2.26) we easily obtain the familiar relations

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \wedge (\mathbf{b} \wedge \mathbf{c}))^c = (\mathbf{a} \lrcorner (\mathbf{b} \wedge \mathbf{c}))^{cc} = -(\mathbf{a} * \mathbf{b})\mathbf{c} + (\mathbf{a} * \mathbf{c})\mathbf{b},$$

and

$$\mathbf{a} * (\mathbf{b} \times \mathbf{c}) = \mathbf{a} \lrcorner (\mathbf{b} \wedge \mathbf{c})^c = (\mathbf{a} \wedge \mathbf{b} \wedge \mathbf{c}) * I^\dagger,$$

which by Lemma 2.12 and a choice of basis becomes the usual determinant expression.

*Remark.* It is instructive to compare the exterior algebra  $(\mathcal{G}, \wedge)$  together with this duality operation to the language of differential forms and the Hodge  $*$  duality operation, which are completely equivalent (through the isomorphism  $(\mathcal{G}, \wedge) \cong \wedge^* V$ ). In that setting one often starts with the outer product and then uses a given metric tensor to define a dual. The inner product is then defined from the outer product and dual according to (2.28). The exact definition of the dual varies, but a common choice in the literature for differential forms is  $*x := (I^{-1}x)^\dagger$ , so that  $**x = IxI^\dagger$  (cp. e.g. [34]).

---

**Exercise 2.12.** Show that  $|A\triangle B| \equiv |A| + |B| \pmod{2}$  and verify Eqs. (2.17)-(2.19) from the definitions of these involutions.

**Exercise 2.13.** Prove the remaining identities in Proposition 2.9.

**Exercise 2.14.** Prove Proposition 2.10.

**Exercise 2.15.** Define a bilinear form on  $\mathcal{G}(V, q)$  via

$$\langle x, y \rangle_{\mathcal{G}} := \langle x^\dagger y \rangle_0,$$

and verify that this extends the scalar product on  $V = \mathbb{R}^n$  to a proper inner product on the full euclidean algebra  $\mathcal{G}(\mathbb{R}^n)$ . Also, verify that the operations  $\lrcorner$  and  $\wedge$  on  $\mathcal{G}$  are ‘reverse-adjoint’ to each other w.r.t. to the above bilinear form, in the sense that for all  $x, y, z \in \mathcal{G}$

$$\langle x, z \lrcorner y \rangle_{\mathcal{G}} = \langle z^\dagger \wedge x, y \rangle_{\mathcal{G}}.$$

**Exercise 2.16.** Prove Proposition 2.11.

**Exercise 2.17.** Show that the algebra  $(\mathcal{G}, \vee)$  with the meet product is associative with unit  $I$ . (The combined algebra  $(\mathcal{G}, \wedge, \vee)$  is sometimes called *the double Cayley algebra*.)

**Exercise 2.18.** Show that the cross product as defined in (2.29) gives rise to the familiar cartesian coordinate expression (and has the correct orientation) when expanding in an orthonormal basis.

**Exercise 2.19.** For  $v \in \mathcal{Cl}^1$  define the map

$$\begin{aligned} \partial_v : \mathcal{Cl} &\rightarrow \mathcal{Cl}, \\ x &\mapsto \partial_v(x) = v \lrcorner x. \end{aligned}$$

Verify that  $\partial_v \circ \partial_v = 0$  and that

$$\partial_v(xy) = \partial_v(x)y + x^* \partial_v(y)$$

as well as

$$\partial_v(x \wedge y) = \partial_v(x) \wedge y + x^* \wedge \partial_v(y)$$

for all  $x, y \in \mathcal{Cl}$  (such  $\partial_v$  is called an *anti-derivation*). Furthermore, note that this leads to the following useful formula:

$$v \lrcorner (a_1 \wedge a_2 \wedge \cdots \wedge a_m) = \sum_{k=1}^m (-1)^{k-1} (v * a_k) a_1 \wedge \cdots \wedge \check{a}_k \wedge \cdots \wedge a_m, \quad (2.30)$$

(where  $\check{\phantom{a}}$  denotes deletion) for any  $v, a_1, \dots, a_m \in \mathcal{Cl}^1$ . A generalized expansion for higher grades can be found in Appendix A.3.

**Exercise 2.20.** Let  $v \in \mathbb{R}^3$  and  $B \in \mathcal{G}^2(\mathbb{R}^3)$ . In each of the following cases, find  $w \in \mathcal{G}(\mathbb{R}^3)$  that satisfies

a)  $w \wedge w = 1 + v + B$

b)  $w \wedge (1 + v + B) = 1$

**Exercise 2.21.** Show that  $\mathcal{Cl}^2$  is closed under the commutator bracket, i.e.  $[A, B] := AB - BA \in \mathcal{Cl}^2$  for all  $A, B \in \mathcal{Cl}^2$ .

**Exercise 2.22.** Show that  $x_p \wedge x_r = (-1)^{pr} x_r \wedge x_p$  for all  $x_k \in \mathcal{Cl}^k$ .

### 3 Vector space geometry

We will now leave the general setting of combinatorial Clifford algebra for a moment and instead focus on the geometric properties of  $\mathcal{G}$  and its newly defined operations in the context of vector spaces and linear transformations.

#### 3.1 Blades and subspaces

The concept of blades is central for understanding the geometry encoded in a geometric algebra  $\mathcal{G}(V)$ .

**Definition 3.1.** A *blade*, or *simple multivector*, is an outer product of 1-vectors. We define the following:

$$\begin{aligned} \mathcal{B}_k &:= \{v_1 \wedge v_2 \wedge \cdots \wedge v_k \in \mathcal{G} : v_i \in V\} && \text{the set of } k\text{-blades} \\ \mathcal{B} &:= \bigcup_{k=0}^{\infty} \mathcal{B}_k && \text{the set of all blades} \\ \mathcal{B}^\bullet &:= \mathcal{B} \setminus \{0\} && \text{the nonzero blades} \\ \mathcal{B}^\times &:= \{B \in \mathcal{B} : B^2 \neq 0\} && \text{the invertible blades} \end{aligned}$$

The *orthogonal basis blades* associated to an orthogonal basis  $E = \{e_i\}_{i=1}^{\dim V}$  consist of the basis of  $\mathcal{G}$  generated by  $E$ , i.e.

$$\mathcal{B}_E := \{e_{i_1} \wedge e_{i_2} \wedge \cdots \wedge e_{i_k} \in \mathcal{G} : i_1 < i_2 < \cdots < i_k\}$$

(corresponding to  $\mathcal{P}(E)$  in  $\mathcal{Cl}$ ). We also include the unit 1 among the blades and call it the *0-blade*.

Note that  $\mathcal{B}_k \subseteq \mathcal{G}^k$  and that (e.g. by expanding in an orthogonal basis; see Exercise 3.1) we can expand a blade as a sum of geometric products,

$$a_1 \wedge a_2 \wedge \cdots \wedge a_k = \frac{1}{k!} \sum_{\pi \in S_k} \text{sign}(\pi) a_{\pi(1)} a_{\pi(2)} \cdots a_{\pi(k)}. \quad (3.1)$$

This expression is clearly similar to a determinant, except that this is a product of *vectors* instead of scalars.

The key property of blades is that they represent linear subspaces of  $V$ . This is made precise by the following

**Proposition 3.1.** *If  $A = a_1 \wedge a_2 \wedge \cdots \wedge a_k \neq 0$  is a nonzero  $k$ -blade and  $a \in V$  then*

$$a \wedge A = 0 \quad \Leftrightarrow \quad a \in \text{Span}\{a_1, a_2, \dots, a_k\}.$$

*Proof.* This follows directly from Theorem 2.13 since  $\{a_1, \dots, a_k\}$  are linearly independent, and  $a \wedge A = 0$  if and only if  $\{a, a_1, \dots, a_k\}$  are linearly dependent.  $\square$

Hence, to every nonzero  $k$ -blade  $A = a_1 \wedge a_2 \wedge \cdots \wedge a_k$  there corresponds a unique  $k$ -dimensional subspace

$$\bar{A} := \{a \in V : a \wedge A = 0\} = \text{Span}\{a_1, a_2, \dots, a_k\}. \quad (3.2)$$

Conversely, if  $\bar{A} \subseteq V$  is a  $k$ -dimensional subspace of  $V$ , then we can find a nonzero  $k$ -blade  $A$  representing  $\bar{A}$  by simply taking a basis  $\{a_i\}_{i=1}^k$  of  $\bar{A}$  and forming

$$A := a_1 \wedge a_2 \wedge \cdots \wedge a_k. \quad (3.3)$$

We thus have the geometric interpretation of blades as subspaces with an associated orientation (sign) and magnitude. Since every element in  $\mathcal{G}$  is a linear combination of orthogonal basis blades, we can think of every element as representing a linear combination of orthogonal subspaces. In the case of a nondegenerate algebra these basis subspaces are nondegenerate as well. On the other hand, any blade which represents a nondegenerate subspace can also be treated as a basis blade associated to an orthogonal basis. This will follow in the discussion below.

**Proposition 3.2.** *Every  $k$ -blade can be written as a geometric product of  $k$  vectors.*

*Proof.* Take a nonzero  $A = a_1 \wedge \cdots \wedge a_k \in \mathcal{B}^\bullet$ . Pick an orthogonal basis  $\{e_i\}_{i=1}^k$  of the subspace  $(\bar{A}, q|_{\bar{A}})$ . Then we can write  $a_i = \sum_j \beta_{ij} e_j$  for some  $\beta_{ij} \in \mathbb{F}$ , and  $A = \det [\beta_{ij}] e_1 e_2 \cdots e_k$  by (3.1).  $\square$

There are a number of useful consequences of this result.

**Corollary.** *For any blade  $A \in \mathcal{B}$ ,  $A^2$  is a scalar.*

*Proof.* Use the expansion of  $A$  above to obtain

$$A^2 = (\det [\beta_{ij}])^2 (-1)^{\frac{1}{2}k(k-1)} q(e_1)q(e_2) \cdots q(e_k) \in \mathbb{F}. \quad (3.4)$$

$\square$

**Corollary.** *If  $A \in \mathcal{B}^\times$  then  $A$  has an inverse  $A^{-1} = \frac{1}{A^2} A$ .*

**Example 3.1.** In  $\mathbb{R}^3$ , every bivector is also a 2-blade. This follows by duality, since if  $B$  is a bivector then  $B^c = BI^{-1} =: b$  is a vector. Choosing an orthonormal basis  $\{e_1, e_2, e_3\}$  such that, say,  $b = \beta e_3$ , we have  $B = bI = \beta e_1 \wedge e_2$ . The same situation is not true in  $\mathbb{R}^4$ , however, where we e.g. have a bivector

$$B := e_1 \wedge e_2 + e_3 \wedge e_4 = e_1 e_2 (1 - I)$$

such that  $B^2 = -(1 - I)^2 = -2(1 - I)$ , which is not a scalar. (Actually, one can show in general (see Exercise 6.12) that  $B^2 \in \mathbb{F}$  is both a necessary and sufficient condition for a bivector  $B$  to be a blade.)

Another useful consequence of Proposition 3.2 is the following

**Corollary.** *If  $A \in \mathcal{B}^\times$  then  $q$  is nondegenerate on  $\bar{A}$  and there exists an orthogonal basis  $E$  of  $V$  such that  $A \in \mathcal{B}_E$ .*

*Proof.* The first statement follows directly from (3.4). For the second statement note that, since  $q$  is nondegenerate on  $\bar{A}$ , we have  $\bar{A} \cap \bar{A}^\perp = 0$ . Take an orthogonal basis  $\{e_i\}_{i=1}^k$  of  $\bar{A}$ . For any  $v \in V$  we have that

$$v - \sum_i \beta_q(v, e_i) q(e_i)^{-1} e_i \in \bar{A}^\perp.$$

Thus,  $V = \bar{A} \oplus \bar{A}^\perp$  and we can extend  $\{e_i\}_{i=1}^k$  to an orthogonal basis of  $V$  consisting of one part in  $\bar{A}$  and one part in  $\bar{A}^\perp$ . By rescaling this basis we have  $A = e_1 \wedge \cdots \wedge e_k$ .  $\square$



*Remark.* Note that if we have an orthogonal basis of a subspace of  $V$  where  $g$  is degenerate, then it may not be possible to extend this basis to an orthogonal basis for all of  $V$ . Another way to state this is that one or more vectors in the corresponding product expansion will fail to be invertible. If the space is euclidean or anti-euclidean, though, orthogonal bases can always be extended (e.g. using the Gram-Schmidt algorithm; see Exercise 3.8).

**Example 3.2.** The simplest lorentzian space is  $\mathbb{R}^{1,1}$ , with an orthonormal basis conventionally denoted  $\{e_0, e_1\}$ ,  $e_0^2 = 1$ ,  $e_1^2 = -1$ . It has precisely two one-dimensional degenerate subspaces  $\mathbb{R}n_{\pm}$ , where

$$n_{\pm} := e_0 \pm e_1, \quad n_{\pm}^2 = 0,$$

but these cannot be orthogonal. In fact,

$$n_+ * n_- = e_0^2 - e_1^2 = 2,$$

but they of course span the space since they are linearly independent, and  $n_+ \wedge n_- = 2e_1e_0$ . In general, the degenerate part of a  $n$ -dimensional lorentzian space will be a  $(n-1)$ -dimensional cone, called the *null-* or *light-cone*.

It is useful to be able to work efficiently also with general bases of  $V$  and  $\mathcal{G}$  which need not be orthogonal. Let  $\{e_1, \dots, e_n\}$  be *any* basis of  $V$ . Then the set of blades formed out of this basis,  $\{e_{\mathbf{i}}\}_{\mathbf{i}}$ , is a basis of  $\mathcal{G}(V)$ , where we use a multi-index notation

$$\mathbf{i} = (i_1, i_2, \dots, i_k), \quad i_1 < i_2 < \dots < i_k, \quad 0 \leq k \leq n \quad (3.5)$$

and

$$e_{()} := 1, \quad e_{(i_1, i_2, \dots, i_k)} := e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k}. \quad (3.6)$$

Sums over  $\mathbf{i}$  are understood to be performed over all allowed such indices. If  $\mathcal{G}$  is nondegenerate then the scalar product  $(A, B) \mapsto A * B$  is also nondegenerate and we can find a so-called *reciprocal basis*  $\{e^1, \dots, e^n\}$  of  $V$  such that

$$e^i * e_j = \delta_j^i. \quad (3.7)$$

The reciprocal basis is easily verified (Exercise 3.2) to be given by

$$e^i = (-1)^{i-1} (e_1 \wedge \dots \wedge \check{e}_i \wedge \dots \wedge e_n) e_{(1, \dots, n)}^{-1}, \quad (3.8)$$

where  $\check{\phantom{x}}$  denotes a deletion. Furthermore, we have that  $\{e^{\mathbf{i}}\}_{\mathbf{i}}$  is a reciprocal basis of  $\mathcal{G}$ , where  $e^{(i_1, \dots, i_k)} := e^{i_k} \wedge \dots \wedge e^{i_1}$  (note the order). This follows since by Lemma 2.12 and (3.7),

$$e^{\mathbf{i}} * e_{\mathbf{j}} = (e^{i_k} \wedge \dots \wedge e^{i_1}) * (e_{j_1} \wedge \dots \wedge e_{j_l}) = \delta_{\mathbf{l}}^{\mathbf{k}} \det [e^{i_p} * e_{j_q}]_{1 \leq p, q \leq k} = \delta_{\mathbf{j}}^{\mathbf{i}}. \quad (3.9)$$

We now have the coordinate expansions

$$\begin{aligned} v &= \sum_{\mathbf{i}} (v * e^{\mathbf{i}}) e_{\mathbf{i}} = \sum_{\mathbf{i}} (v * e_{\mathbf{i}}) e^{\mathbf{i}} \quad \forall v \in V, \\ x &= \sum_{\mathbf{i}} (x * e^{\mathbf{i}}) e_{\mathbf{i}} = \sum_{\mathbf{i}} (x * e_{\mathbf{i}}) e^{\mathbf{i}} \quad \forall x \in \mathcal{G}(V), \end{aligned} \quad (3.10)$$

which also gives a geometric understanding of an arbitrary multivector in  $\mathcal{G}(V)$  as a linear combination of general (not necessarily orthogonal) subspaces of  $V$ .

**Example 3.3.** In the euclidean case an orthonormal basis  $\{e_i\}_i$ , satisfying  $e_i^2 = 1$  for all  $i$ , is its own reciprocal basis, i.e.  $e^i = e_i$ . In the lorentzian case  $\mathbb{R}^{1,n}$  the standard orthonormal basis  $\{e_0, e_1, \dots, e_n\}$ , with  $e_0^2 = 1$ , has as reciprocal basis  $\{e^0, e^1, \dots, e^n\}$  where  $e^0 = e_0$  and  $e^j = -e_j$ ,  $j = 1, 2, \dots, n$ . The non-orthogonal basis  $\{n_+, n_-\}$  in Example 3.2 has the reciprocal basis  $\{\frac{1}{2}n_-, \frac{1}{2}n_+\}$ .

In addition to being useful in coordinate expansions, the general and reciprocal bases also provide a geometric understanding of the dual operation because of the following

**Theorem 3.3.** *Assume that  $\mathcal{G}$  is nondegenerate. If  $A = a_1 \wedge \dots \wedge a_k \in \mathcal{B}^\bullet$  and we extend  $\{a_i\}_{i=1}^k$  to a basis  $\{a_i\}_{i=1}^n$  of  $V$  then*

$$A^c \propto a^{k+1} \wedge a^{k+2} \wedge \dots \wedge a^n,$$

where  $\{a^i\}_i$  is the reciprocal basis of  $\{a_i\}_i$ .

*Proof.* We obtain by induction on  $k$  that

$$\begin{aligned} & (a_k \wedge a_{k-1} \wedge \dots \wedge a_1) \lrcorner (a^1 \wedge \dots \wedge a^k \wedge a^{k+1} \wedge \dots \wedge a^n) \\ &= a_k \lrcorner ((a_{k-1} \wedge \dots \wedge a_1) \lrcorner (a^1 \wedge \dots \wedge a^{k-1} \wedge a^k \wedge \dots \wedge a^n)) \\ &= \{\text{induction assumption}\} = a_k \lrcorner (a^k \wedge \dots \wedge a^n) \\ &= a^{k+1} \wedge \dots \wedge a^n, \end{aligned}$$

where in the last step we used the expansion formula (2.30), plus orthogonality (3.7). It follows that

$$A^c = A \lrcorner I^{-1} \propto (a_k \wedge \dots \wedge a_1) \lrcorner (a^1 \wedge \dots \wedge a^k \wedge a^{k+1} \wedge \dots \wedge a^n) = a^{k+1} \wedge \dots \wedge a^n.$$

This can also be proved using an expansion of the inner product into sub-blades; see Appendix A.3.  $\square$

**Corollary.** *If  $A$  and  $B$  are blades then  $A^c$ ,  $A \wedge B$ ,  $A \vee B$  and  $A \lrcorner B$  are blades as well.*

The blade-subspace correspondence then gives us a geometric interpretation of these operations.

**Proposition 3.4.** *If  $A, B \in \mathcal{B}^\bullet$  are nonzero blades then  $\overline{A^c} = \bar{A}^\perp$  and*

$$\begin{aligned} A \wedge B \neq 0 &\Rightarrow \overline{A \wedge B} = \bar{A} + \bar{B} \text{ and } \bar{A} \cap \bar{B} = 0, \\ \bar{A} + \bar{B} = V &\Rightarrow \overline{A \vee B} = \bar{A} \cap \bar{B}, \\ A \lrcorner B \neq 0 &\Rightarrow \overline{A \lrcorner B} = \bar{A}^\perp \cap \bar{B}, \\ \bar{A} \subseteq \bar{B} &\Rightarrow A \lrcorner B = AB, \\ \bar{A} \cap \bar{B}^\perp \neq 0 &\Rightarrow A \lrcorner B = 0. \end{aligned}$$

The proofs of the statements in the above corollary and proposition are left as exercises. Some of them can be found in [43] and [21].

**Exercise 3.1.** Prove Eqn. (3.1), e.g. by noting that both sides are multilinear and alternating.

**Exercise 3.2.** Verify that  $\{e^i\}_i$  defined in (3.8) is a reciprocal basis w.r.t.  $\{e_i\}_i$ .

**Exercise 3.3.** Prove the corollary to Theorem 3.3.

**Exercise 3.4.** Prove Proposition 3.4.

**Exercise 3.5.** Show that the square of an arbitrary blade  $A = a_1 \wedge \dots \wedge a_k \in \mathcal{B}_k$  is  $A^2 = (-1)^{\frac{1}{2}k(k-1)} \det[a_i * a_j]_{1 \leq i, j \leq k}$ . Also, show in two different ways that

$$(a \wedge b)^2 = (a * b)^2 - a^2 b^2 \quad (3.11)$$

for  $a, b \in V$ .

**Exercise 3.6.** Let  $\{e_i\}_{i=1}^n$  be an arbitrary basis of a nondegenerate space. Show that  $\sum_i e_i e^i = n$  and that for an arbitrary grade- $r$  multivector  $A_r \in \mathcal{C}l^r$

$$\sum_i e_i (e^i \lrcorner A_r) = r A_r,$$

$$\sum_i e_i (e^i \wedge A_r) = (n - r) A_r,$$

and

$$\sum_i e_i A_r e^i = (-1)^r (n - 2r) A_r.$$

**Exercise 3.7.** Let  $v \in V^\times = \{u \in V : u^2 \neq 0\}$ . Show that the map

$$V \ni x \mapsto \frac{1}{2}(x + v^* x v^{-1}) \in V$$

is an orthogonal projection on  $\bar{v}^\perp = \{u \in V : u * v = 0\}$ .

**Exercise 3.8.** Consider the real euclidean geometric algebra  $\mathcal{G}(\mathbb{R}^n)$ . Let  $a_1, a_2, \dots$  be a sequence of 1-vectors in  $\mathcal{G}$ , and form the blades  $A_0 = 1, A_k = a_1 \wedge a_2 \wedge \dots \wedge a_k$ . Now, let  $b_k = A_{k-1}^\dagger A_k$ . Show that  $b_1, b_2, \dots$  are the vectors obtained using the Gram-Schmidt orthogonalization procedure from  $a_1, a_2, \dots$

## 3.2 Linear functions

Since  $\mathcal{G}$  is itself a vector space which embeds  $V$ , it is natural to consider the properties of linear functions on  $\mathcal{G}$ . There is a special class of such functions, called outermorphisms, which can be said to respect the graded/exterior structure of  $\mathcal{G}$  in a natural way. We will see that, just as the geometric algebra  $\mathcal{G}(V, q)$  is completely determined by the underlying vector space  $(V, q)$ , an outermorphism is completely determined by its behavior on  $V$ .

**Definition 3.2.** A linear map  $F: \mathcal{G} \rightarrow \mathcal{G}'$  is called an *outermorphism* or  $\wedge$ -*morphism* if

- i)  $F(1) = 1$ ,
- ii)  $F(\mathcal{G}^m) \subseteq \mathcal{G}'^m \quad \forall m \geq 0$ , (grade preserving)
- iii)  $F(x \wedge y) = F(x) \wedge F(y) \quad \forall x, y \in \mathcal{G}$ .

A linear transformation  $F : \mathcal{G} \rightarrow \mathcal{G}$  is called a *dual outermorphism* or  $\vee$ -*morphism* if

- i)  $F(I) = I$ ,
- ii)  $F(\mathcal{G}^m) \subseteq \mathcal{G}^m \quad \forall m \geq 0$ ,
- iii)  $F(x \vee y) = F(x) \vee F(y) \quad \forall x, y \in \mathcal{G}$ .

**Theorem 3.5.** *For every linear map  $f : V \rightarrow W$  there exists a unique outermorphism  $f_\wedge : \mathcal{G}(V) \rightarrow \mathcal{G}(W)$  such that  $f_\wedge(v) = f(v) \quad \forall v \in V$ .*

*Proof.* Take a general basis  $\{e_1, \dots, e_n\}$  of  $V$  and define, for  $1 \leq i_1 < i_2 < \dots < i_m \leq n$ ,

$$f_\wedge(e_{i_1} \wedge \dots \wedge e_{i_m}) := f(e_{i_1}) \wedge \dots \wedge f(e_{i_m}), \quad (3.12)$$

and extend  $f_\wedge$  to the whole of  $\mathcal{G}(V)$  by linearity. We also define  $f_\wedge(\alpha) := \alpha$  for  $\alpha \in \mathbb{F}$ . Hence, (i) and (ii) are satisfied. (iii) is easily verified by expanding in the induced basis  $\{e_i\}$  of  $\mathcal{G}(V)$ . Uniqueness is obvious since our definition was necessary.  $\square$

Uniqueness immediately implies the following.

**Corollary.** *If  $f : V \rightarrow V'$  and  $g : V' \rightarrow V''$  are linear then  $(g \circ f)_\wedge = g_\wedge \circ f_\wedge$ .*

**Corollary.** *If  $F : \mathcal{G}(V) \rightarrow \mathcal{G}(W)$  is an outermorphism then  $F = (F|_V)_\wedge$ .*

*Remark.* In the setting of  $\mathcal{Cl}$  this means that an outermorphism  $F : \mathcal{Cl}(X, R, r) \rightarrow \mathcal{Cl}(X', R, r')$  is completely determined by its values on the elements of  $X$ . Also note that, if  $E, F$  are two orthonormal bases of  $V$  and  $F = \{f_j = g(e_j)\}$  then

$$\mathcal{P}(E) \ni A_E \xrightarrow{g_\wedge} A_F \in \mathcal{P}(F)$$

so that  $g_\wedge : \mathcal{Cl}(E) \rightarrow \mathcal{Cl}(F)$  is an isomorphism of  $R$ -algebras.

We have seen that a nondegenerate  $\mathcal{G}$  results in a nondegenerate bilinear form  $x * y$ . This gives a canonical isomorphism

$$\theta : \mathcal{G} \rightarrow \mathcal{G}^* = \text{Lin}(\mathcal{G}, \mathbb{F})$$

between the elements of  $\mathcal{G}$  and the linear functionals on  $\mathcal{G}$  as follows. For every  $x \in \mathcal{G}$  we define a linear functional  $\theta(x)$  by  $\theta(x)(y) := x * y$ . Taking a general basis  $\{e_i\}_i$  of  $\mathcal{G}$  and using (3.9) we obtain a dual basis  $\{\theta(e^i)\}_i$  such that  $\theta(e^i)(e_j) = \delta_j^i$ . This shows that  $\theta$  is an isomorphism.

Now that we have a canonical way of moving between  $\mathcal{G}$  and its dual space  $\mathcal{G}^*$ , we can for every linear map  $F : \mathcal{G} \rightarrow \mathcal{G}$  define an *adjoint map*  $F^* : \mathcal{G} \rightarrow \mathcal{G}$  by

$$F^*(x) := \theta^{-1}(\theta(x) \circ F). \quad (3.13)$$

By definition, this has the expected and unique property

$$F^*(x) * y = x * F(y) \quad (3.14)$$

for all  $x, y \in \mathcal{G}$ . Note that if we restrict our attention to  $V = \mathcal{G}^1$  then this construction results in the usual adjoint (transpose)  $f^*$  of a linear map  $f : V \rightarrow V$ .

**Theorem 3.6** (Hestenes' Theorem). *Assume that  $\mathcal{G}$  is nondegenerate and let  $F: \mathcal{G} \rightarrow \mathcal{G}$  be an outermorphism. Then the adjoint  $F^*$  is also an outermorphism and*

$$\begin{aligned} x \lrcorner F(y) &= F(F^*(x) \lrcorner y), \\ F(x) \lrcorner y &= F(x \lrcorner F^*(y)), \end{aligned}$$

for all  $x, y \in \mathcal{G}$ .

*Proof.* We first prove that  $F^*$  is an outermorphism. The fact that  $F^*$  is grade preserving follows from (3.14) and the grade preserving property of  $F$ . Now take basis blades  $x = x_1 \wedge \cdots \wedge x_m$  and  $y = y_m \wedge \cdots \wedge y_1$  with  $x_i, y_j \in V$ . Then

$$\begin{aligned} F^*(x_1 \wedge \cdots \wedge x_m) * y &= (x_1 \wedge \cdots \wedge x_m) * F(y_m \wedge \cdots \wedge y_1) \\ &= (x_1 \wedge \cdots \wedge x_m) * (F(y_m) \wedge \cdots \wedge F(y_1)) \\ &= \det [x_i * F(y_j)]_{i,j} = \det [F^*(x_i) * y_j]_{i,j} \\ &= (F^*(x_1) \wedge \cdots \wedge F^*(x_m)) * (y_m \wedge \cdots \wedge y_1) \\ &= (F^*(x_1) \wedge \cdots \wedge F^*(x_m)) * y, \end{aligned}$$

where we have used Lemma 2.12. By linearity and nondegeneracy it follows that  $F^*$  is an outermorphism. The first identity stated in the theorem now follows quite easily from Proposition 2.9. For any  $z \in \mathcal{G}$  we have

$$\begin{aligned} z * (x \lrcorner F(y)) &= (z \wedge x) * F(y) = F^*(z \wedge x) * y \\ &= (F^*(z) \wedge F^*(x)) * y = F^*(z) * (F^*(x) \lrcorner y) \\ &= z * F(F^*(x) \lrcorner y). \end{aligned}$$

The nondegeneracy of the scalar product then gives the first identity. The second identity is proven similarly, using that  $(x \lrcorner y) * z = x * (y \wedge z)$ .  $\square$

From uniqueness of outermorphisms we also obtain the following

**Corollary.** *If  $f: V \rightarrow V$  is a linear transformation then  $(f^*)_{\wedge} = (f_{\wedge})^*$ .*

This means that we can simply write  $f_{\wedge}^*$  for the adjoint outermorphism of  $f$ .

Another powerful concept in geometric algebra (or exterior algebra) is the generalization of eigenvectors to so called *eigenblades*. For a function  $f: V \rightarrow V$ , a  $k$ -eigenblade with eigenvalue  $\lambda \in \mathbb{F}$  is a blade  $A \in \mathcal{B}_k$  such that

$$f_{\wedge}(A) = \lambda A. \quad (3.15)$$

Just as eigenvectors can be said to represent invariant 1-dimensional subspaces of a function, a  $k$ -blade with nonzero eigenvalue represents an invariant  $k$ -dimensional subspace. One important example of an eigenblade is the pseudoscalar  $I$ , which represents the whole invariant vector space  $V$ . Since  $f_{\wedge}$  is grade preserving, we must have  $f_{\wedge}(I) = \lambda I$  for some  $\lambda \in \mathbb{F}$  which we call the *determinant* of  $f$ , i.e.

$$f_{\wedge}(I) =: (\det f)I. \quad (3.16)$$

Expanding  $\det f = f_{\wedge}(I) * I^{-1}$  in a basis using Lemma 2.12, one finds that this agrees with the usual definition of the determinant of a linear function (see Exercise 3.13).

**Example 3.4.** Since

$$(\det f)I^2 = f_{\wedge}(I) * I = I * f_{\wedge}^*(I) = (\det f^*)I^2,$$

we immediately find that  $\det f = \det f^*$ . Also, by the corollary to Theorem 3.5 we have

$$(f \circ g)_{\wedge}(I) = f_{\wedge}(g_{\wedge}(I)) = f_{\wedge}((\det g)I) = (\det g)(\det f)I,$$

so that  $\det(fg) = \det f \cdot \det g$ .

**Definition 3.3.** For linear  $F: \mathcal{G} \rightarrow \mathcal{G}$  we define the *dual map*  $F^c: \mathcal{G} \rightarrow \mathcal{G}$  by  $F^c(x) := F(xI)I^{-1}$ , so that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{G} & \xrightarrow{F} & \mathcal{G} \\ (\cdot)^c \downarrow & & \downarrow (\cdot)^c \\ \mathcal{G} & \xrightarrow{F^c} & \mathcal{G} \end{array}$$

**Proposition 3.7.** *We have the following properties of the dual map:*

- i)  $F^{cc} = F$ ,
- ii)  $(F \circ G)^c = F^c \circ G^c$ ,
- iii)  $\text{id}^c = \text{id}$ ,
- iv)  $F(\mathcal{G}^s) \subseteq \mathcal{G}^t \Rightarrow F^c(\mathcal{G}^{\dim V - s}) \subseteq \mathcal{G}^{\dim V - t}$ ,
- v)  $F \wedge\text{-morphism} \Rightarrow F^c \vee\text{-morphism}$ ,
- vi)  $(F^*)^c = (F^c)^*$  if  $F$  is grade preserving,

for all linear  $F, G: \mathcal{G} \rightarrow \mathcal{G}$ .

The proofs are straightforward and left as exercises to the reader. As a special case of Theorem 3.6 we obtain, with  $y = I$  and a linear map  $f: V \rightarrow V$ ,

$$(\det f)xI = f_{\wedge}(f_{\wedge}^*(x)I), \quad (3.17)$$

so that

$$(\det f) \text{id} = f_{\wedge}^c \circ f_{\wedge}^* = f_{\wedge} \circ f_{\wedge}^{*c}. \quad (3.18)$$

If  $\det f \neq 0$  we then have a simple expression for the inverse;

$$f_{\wedge}^{-1} = (\det f)^{-1} f_{\wedge}^{*c}, \quad (3.19)$$

which is essentially the dual of the adjoint ( $F^{*c}$  is sometimes called the *adjugate*  $F^{\text{adj}}$  of  $F$ ).  $f^{-1}$  is obtained by simply restricting to  $V$ .

**Example 3.5.** Recall that an orthogonal transformation  $f \in O(V, q)$  (see (6.4)) satisfies  $f^{-1} = f^*$  and  $\det f = \pm 1$  (see also Exercise 6.3), so in this case (3.19) gives  $f_{\wedge}^c = \pm f_{\wedge}$ .

---

**Exercise 3.9.** Prove the second identity in Theorem 3.6.

**Exercise 3.10.** Prove Proposition 3.7.

**Exercise 3.11.** Let  $B \in \mathcal{G}^2$  be a bivector, and define the map  $\text{ad}_B : \mathcal{G} \rightarrow \mathcal{G}$  by  $\text{ad}_B(x) := [B, x] = Bx - xB$ . Show that  $\text{ad}_B(\mathcal{G}^m) \subseteq \mathcal{G}^m$  and that

$$\text{ad}_B(xy) = \text{ad}_B(x)y + x \text{ad}_B(y),$$

i.e. that  $\text{ad}_B$  is a grade-preserving derivation. Is  $\text{ad}_B$  an outermorphism?

**Exercise 3.12.** Let  $f : V \rightarrow V$  be linear and antisymmetric, i.e.

$$f(u) * v = -u * f(v) \quad \forall u, v \in V.$$

Assuming  $V$  is non-degenerate, show that there is a unique bivector  $B \in \mathcal{G}^2(V)$  such that  $f(v) = \text{ad}_B(v) \forall v \in V$ . Can you give a simple expression for  $B$ ? What if  $V$  is degenerate?

**Exercise 3.13.** Let  $\{a_1, \dots, a_n\}$  be an arbitrary basis of non-degenerate  $V$  and use the reciprocal basis to verify that  $\det f = f(I) * I^{-1}$  is the usual expression for the determinant. Use (3.1) to verify that the definition (3.16) makes sense also for degenerate spaces.

### 3.3 Projections and rejections

Let  $A$  be an invertible blade in a geometric algebra  $\mathcal{G}$ , and define a linear map  $P_A : \mathcal{G} \rightarrow \mathcal{G}$  through

$$P_A(x) := (x \lrcorner A)A^{-1}.$$

**Proposition 3.8.**  $P_A$  is an outermorphism.

*Proof.* We need to prove that  $P_A(x \wedge y) = P_A(x) \wedge P_A(y) \forall x, y \in \mathcal{G}$ . We observe that both sides are linear in both  $x$  and  $y$ . Hence, we can assume that  $x$  and  $y$  are basis blades. Also, since  $A$  was assumed to be invertible, we can by the corollary to Proposition 3.2, without loss of generality, assume that  $A$  is also among the orthogonal basis blades. We obtain

$$P_A(x \wedge y) = ((x \wedge y) \lrcorner A)A^{-1} = (x \cap y = \emptyset)(x \cup y \subseteq A)xyAA^{-1}$$

and

$$\begin{aligned} P_A(x) \wedge P_A(y) &= ((x \lrcorner A)A^{-1}) \wedge ((y \lrcorner A)A^{-1}) \\ &= ((x \subseteq A)xAA^{-1}) \wedge ((y \subseteq A)yAA^{-1}) \\ &= (x \subseteq A)(y \subseteq A)x \wedge y \\ &= (x \subseteq A)(y \subseteq A)(x \cap y = \emptyset)xy. \end{aligned}$$

Furthermore, we obviously have  $P_A(1) = 1$ , and (e.g. by again expanding in a basis) we also observe that  $P_A(V) \subseteq V$ , so  $P_A$  preserves grades.  $\square$

We shall now show that  $P_A$  is a projection. First, note the following

**Proposition 3.9.**  $P_A \circ P_A = P_A$

*Proof.* We would like to prove

$$(((x \lrcorner A)A^{-1}) \lrcorner A) A^{-1} = (x \lrcorner A)A^{-1} \quad \forall x \in \mathcal{G}.$$

As both sides are linear in  $x$ , we assume that  $x$  is a basis blade and obtain

$$(x \lrcorner A)A^{-1} = (x \subseteq A)xA A^{-1} = (x \subseteq A)x$$

and

$$\begin{aligned} (((x \lrcorner A)A^{-1}) \lrcorner A) A^{-1} &= (((x \subseteq A)x) \lrcorner A) A^{-1} \\ &= (x \subseteq A)(x \lrcorner A)A^{-1} = (x \subseteq A)(x \subseteq A)x = (x \subseteq A)x. \end{aligned}$$

□

Also note that, for all  $v \in V$ , we have by Proposition 2.10

$$v = vAA^{-1} = (vA)A^{-1} = (v \lrcorner A + v \wedge A)A^{-1}. \quad (3.20)$$

Now, define the *rejection*  $R_A$  of  $A$  through

$$R_A(x) := (x \wedge A)A^{-1}, \quad \text{for } x \in \mathcal{G}.$$

Then (3.20) becomes

$$v = P_A(v) + R_A(v) \quad \forall v \in V.$$

If  $v \in \bar{A} = \{u \in V : u \wedge A = 0\}$ , we obviously obtain

$$P_A(v) = v \quad \text{and} \quad R_A(v) = 0,$$

and if  $v \in \bar{A}^\perp = \{u \in V : u \lrcorner A = 0\}$ , we find

$$P_A(v) = 0 \quad \text{and} \quad R_A(v) = v.$$

We therefore see that  $P_A$  is the outermorphism of an orthogonal projection on  $\bar{A}$ , while  $R_A$  corresponds to an orthogonal projection on  $\bar{A}^\perp$ .

---

**Exercise 3.14.** Show that  $R_A$  is an outermorphism and that  $P_A$  and  $R_A$  are self-adjoint, i.e.  $P_A^* = P_A$  and  $R_A^* = R_A$ .

### 3.4 Some projective geometry

We finish this section by considering some applications in classical projective geometry.

Given a vector space  $V$ , the *projective geometry*  $P(V)$  on  $V$  is defined to be the set of all subspaces of  $V$ . The set of  $k$ -dimensional subspaces is denoted  $P_k(V)$ . Traditionally,  $P_1(V)$  is called the *projective space* and is usually denoted  $\mathbb{P}^{n-1}(\mathbb{F})$  if  $V$  is an  $n$ -dimensional vector space over the field  $\mathbb{F}$ .

$P(V)$  forms a natural so-called *lattice* with a *meet*  $\wedge$  and *join*  $\vee$  operation defined as follows: Let  $U$  and  $W$  be elements in  $P(V)$ , i.e. subspaces of  $V$ . The largest subspace of  $V$  which is contained in both  $U$  and  $W$  is denoted  $U \wedge W$ ,



and the smallest subspace of  $V$  which contains both  $U$  and  $W$  is denoted  $U \vee W$ . Hence,  $U \wedge W = U \cap W$  and  $U \vee W = U + W$ .

We saw earlier that when  $V$  is considered as embedded in a geometric algebra  $\mathcal{G}(V)$ , then to each nonzero  $k$ -blade  $A \in \mathcal{G}$  there is associated a unique  $k$ -dimensional subspace  $\bar{A} \in P_k(V)$ . Furthermore, we found that the outer and meet products of blades corresponded to the subspace operations<sup>9</sup>

$$\overline{A \wedge B} = \bar{A} \vee \bar{B} \quad \text{if} \quad \bar{A} \wedge \bar{B} = 0$$

and

$$\overline{A \vee B} = \bar{A} \wedge \bar{B} \quad \text{if} \quad \bar{A} \vee \bar{B} = V.$$

Let us now get acquainted with some concrete classical projective geometry in  $\mathbb{R}^n$ .

### 3.4.1 The cross ratio in $\mathbb{P}^1(\mathbb{R})$

Let  $V = \mathbb{R}^2$  and choose  $a, b, c, d \in V$  such that each pair of these vectors is linearly independent. Define the so-called *cross ratio of four points*

$$D(a, b, c, d) = \frac{(a \wedge b)(c \wedge d)}{(a \wedge c)(b \wedge d)}.$$

Note that each of the factors above is a nonzero multiple of the pseudoscalar, so that the above expression makes sense. The expression is a *projective invariant* in the following sense:

- i)  $D(a, b, c, d)$  only depends on  $\bar{a}, \bar{b}, \bar{c}, \bar{d}$ , i.e. on the four one-dimensional subspaces in  $\mathbb{P}^1(\mathbb{R})$  which correspond to the respective vectors.
- ii)  $D(Ta, Tb, Tc, Td) = D(a, b, c, d)$  for every linear bijection  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ .

We can reinterpret the projectively invariant quantity  $D(a, b, c, d)$  as a universal property of four points in  $\mathbb{R}^1$ . Namely, choose a line  $L$  away from the origin in  $V$ . By applying a projective transformation, i.e. a linear bijection, we can without loss of generality assume that the line or basis is such that  $L = \mathbb{R}e_1 + e_2$ . Then, representing a point  $A \in \mathbb{R}$  as  $a = Ae_1 + e_2 \in L$  etc., we find  $a \wedge b = (A - B)I$  and that the cross ratio is

$$D(a, b, c, d) = \frac{(A - B)(C - D)}{(A - C)(B - D)}. \quad (3.21)$$

Projective invariance, i.e. the possibility of mapping  $L$  to any other line away from the origin, then means that this quantity is invariant under both translations, rescalings and inversions in  $\mathbb{R}^1$ .

---

**Exercise 3.15.** Verify properties (i) and (ii) above. Derive the general form of the transformations of  $\mathbb{R}^1$  which leave the cross ratio invariant, and verify explicitly that the expression (3.21) is invariant under such a transformation.

<sup>9</sup>In this context the choice of notation is slightly unfortunate. However, both the notation for meet and join of subspaces (which matches the logical operations) and the outer and meet products of multivectors are strictly conventional.




Figure 3.1: Mapping a point  $\mathbf{x} \in \mathbb{R}^3$  to its projective representative  $X \in \mathbb{R}^4$ .

**Exercise 3.16.** Let us consider  $\mathcal{G}(\mathbb{R}^3 = \text{Span}\{e_1, e_2, e_3\})$  as a part of  $\mathcal{G}(\mathbb{R}^4 = \text{Span}\{e_1, e_2, e_3, e_4\})$  via the map

$$\begin{aligned} \mathbb{R}^3 &\rightarrow \mathbb{R}^4 \\ \mathbf{x} &\mapsto X := \mathbf{x} + e_4 \end{aligned}$$

which is represented in Figure 3.1. We denote the respective pseudoscalars by  $I = e_1e_2e_3$  and  $J = e_1e_2e_3e_4$ . Let  $\times$  denote the usual cross product in  $\mathbb{R}^3$  (as defined in (2.29)) and verify that

- a)  $\mathbf{x} \times \mathbf{y} = (e_4 \wedge X \wedge Y)J$
- b)  $\mathbf{y} - \mathbf{x} = e_4 \lrcorner (X \wedge Y)$

**Exercise 3.17.** Let  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  and  $\mathbf{d}$  be distinct vectors in  $\mathbb{R}^2$ . We are interested in determining the intersection of the lines through  $\mathbf{a}, \mathbf{b}$  and  $\mathbf{c}, \mathbf{d}$  respectively. As in the previous exercise, we consider a map (with a basis  $\{e_1, e_2, e_3\}$ )

$$\begin{aligned} \mathbb{R}^2 &\rightarrow \mathbb{R}^3 \\ \mathbf{x} &\mapsto X := \mathbf{x} + e_3 \end{aligned}$$

Introduce  $L = A \wedge B$  and  $M = C \wedge D$  and show that

$$L \vee M = [A, B, C]D - [A, B, D]C,$$

where  $[X, Y, Z] := (X \wedge Y \wedge Z) * (e_3 \wedge e_2 \wedge e_1)$  for  $X, Y, Z \in \mathbb{R}^3$ . Interpret this result geometrically.

### 3.4.2 Cross ratios and quadrics in $\mathbb{P}^2(\mathbb{R})$

Now, let  $V = \mathbb{R}^3$  (we could also choose  $\mathbb{R}^{s,t}$  with  $s+t=3$ ). Choose  $a, b, c, d, e \in V$ , pairwise linearly independent, such that  $a, b, c, d$  all lie in a two-dimensional subspace not containing  $e$ . In the projective jargon one then says that the projective points  $\bar{a}, \bar{b}, \bar{c}, \bar{d}$  lie on a line which does not contain  $\bar{e}$ . Let us think of the plane of this page as an affine<sup>10</sup> subspace of  $V$  which does not contain the origin, and which intersects the lines  $\bar{a}, \bar{b}, \bar{c}$ , and  $\bar{d}$  at points which we represent by the names of those lines. Let us imagine that we have one eye in the origin and that the line through the eye and the point in the paper denoted  $\bar{a}$  is the line  $\bar{a}$ . We then have a picture like in Figure 3.2.

Define

$$F(a, b, c, d, e) := \frac{(a \wedge b \wedge e)(c \wedge d \wedge e)}{(a \wedge c \wedge e)(b \wedge d \wedge e)}.$$

<sup>10</sup>i.e. constant plus linear




Figure 3.2: Projective points  $\bar{a}, \bar{b}, \bar{c},$  and  $\bar{d}$  on a common projective line which does not contain  $\bar{e}$ .

We immediately see that for all  $\alpha, \beta, \gamma, \delta, \epsilon \in \mathbb{R}^\times$ ,

$$F(\alpha a, \beta b, \gamma c, \delta d, \epsilon e) = F(a, b, c, d, e),$$

which shows that  $F$  only depends on  $\bar{a}, \bar{b}, \bar{c}, \bar{d}$  and  $\bar{e}$  in  $\mathbb{P}^2(\mathbb{R})$ . Furthermore, if  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is a linear bijection then

$$F(Ta, Tb, Tc, Td, Te) = \frac{(\det T)^2}{(\det T)^2} F(a, b, c, d, e),$$

and  $F(a, b, c, d, e)$  is a so-called projective invariant of five points.

So far we have not used that the projective points  $\bar{a}, \bar{b}, \bar{c},$  and  $\bar{d}$  lie on a common line. We shall now show, with our requirements on  $a, b, c, d, e$ , that  $F$  does not depend on  $e$ . Assume that  $e' = e + \alpha a + \beta b$ . Then  $a \wedge b \wedge e' = a \wedge b \wedge e$  and

$$c \wedge d \wedge e' = c \wedge d \wedge e + \alpha c \wedge d \wedge a + \beta c \wedge d \wedge b = c \wedge d \wedge e,$$

since  $c \wedge d \wedge a = c \wedge d \wedge b = 0$  (they all lie in the same plane in  $\mathbb{R}^3$ ). Similarly, we find that we could have added linear combinations of  $c$  and  $d$  as well, and that the factors in the denominator also do not depend on  $e$ . Hence, we can define the cross ratio

$$D(a, b, c, d) := F(a, b, c, d, e),$$

which then becomes a projective invariant of  $\bar{a}, \bar{b}, \bar{c}, \bar{d}$  with a similar geometric interpretation as before whenever these points lie on a common line.

Let us now have a look at quadrics in  $\mathbb{P}^2(\mathbb{R})$ . Let  $P(x_1, x_2, x_3)$  denote a homogeneous quadratic polynomial<sup>11</sup> in the (ordinary, commutative) polynomial ring  $\mathbb{R}[x_1, x_2, x_3]$ . From the homogeneity follows that if  $(\alpha, \beta, \gamma)$  is a zero of the polynomial  $P$ , then also  $t(\alpha, \beta, \gamma)$  is a zero, for all  $t \in \mathbb{R}$ . Hence, if we (as usual) interpret the triple  $(x_1, x_2, x_3)$  as the point  $x = x_1 e_1 + x_2 e_2 + x_3 e_3 \in \mathbb{R}^3$  then we see that zeros of  $P$  can be interpreted as points in  $\mathbb{P}^2(\mathbb{R})$ . The zero set  $Z(P)$  of a homogeneous quadratic polynomial  $P$  is called a *quadric*.

**Example 3.6.** Consider the homogeneous quadratic polynomial in three variables

$$P(x, y, z) = x^2 + y^2 - z^2$$

Note that the intersection of the quadric  $Z(P)$  with the plane  $\{(x, y, z) \in \mathbb{R}^3 : z = 1\}$  is a unit circle, but by choosing some other plane we may get an ellipse, a hyperbola, or even a parabola or a line.

<sup>11</sup>I.e. a quadratic form, but here we think of it as a polynomial rather than an extra structure of the vector space.

It is easy to see that five distinct points in  $\mathbb{P}^2(\mathbb{R})$  determine a homogeneous quadratic polynomial in three variables, up to a multiple, if we demand that the five points should be zeros of the polynomial. Assume that we are given five points  $a, b, c, d, e$  in  $\mathbb{R}^3$ , and form

$$P(x) = (a \wedge b \wedge e)(c \wedge d \wedge e)(a \wedge c \wedge x)(b \wedge d \wedge x) - (a \wedge b \wedge x)(c \wedge d \wedge x)(a \wedge c \wedge e)(b \wedge d \wedge e)$$

$P$  is then a homogeneous quadratic polynomial in  $(x_1, x_2, x_3)$ , where  $x = x_1 e_1 + x_2 e_2 + x_3 e_3$ . One verifies by direct substitution that

$$P(a) = P(b) = P(c) = P(d) = P(e) = 0,$$

and hence that  $Z(P)$  is a quadric containing  $\bar{a}, \bar{b}, \bar{c}, \bar{d}$ , and  $\bar{e}$ .

### 3.4.3 Pascal's theorem

There is an alternative way of finding a quadric through five points, which is equivalent to a classical theorem in plane projective geometry due to Pascal.

**Theorem 3.10** (Pascal). *Let  $a, b, c, d, e, f$  be points on a quadric in  $\mathbb{P}^2(\mathbb{R})$ . Then*

$$((a \wedge b) \vee (d \wedge e)) \wedge ((b \wedge c) \vee (e \wedge f)) \wedge ((c \wedge d) \vee (f \wedge a)) = 0 \quad (3.22)$$

*Proof.* Form the homogeneous quadratic polynomial  $P(x)$  which is obtained from the expression (3.22) when e.g.  $f$  is replaced with  $x$ . One easily verifies (e.g. using Exercise 3.17) that  $P(a) = P(b) = P(c) = P(d) = P(e) = 0$ , from which the theorem follows.  $\square$

The geometric interpretation of the expression (3.22) is that the three intersection points of the line  $a \wedge b$  with the line  $d \wedge e$ , the line  $b \wedge c$  with  $e \wedge f$ , resp.  $c \wedge d$  with  $f \wedge a$ , are all situated on one and the same line. See Figure 3.3.




Figure 3.3: A geometric interpretation of Pascal's theorem.

### 3.4.4 \*Polarizations in quadrics

Let  $Q$  be a quadric in  $\mathbb{P}^2$  and choose a point  $p \in \mathbb{P}^2$  according to Figure 3.4. Draw the tangent lines  $L_1$  and  $L_2$  to  $Q$  through  $p$  and form the line  $L$  through the tangent points  $t_1$  and  $t_2$ . Then  $L$  is called the *polar line* to  $p$  with respect to  $Q$ . In classical projective geometry there are a number of theorems related to this notion. We will now show how geometric algebra can be used to obtain simple proofs and more algebraic formulations of some of these theorems.




Figure 3.4: The polar line  $L$  to the point  $p$  with respect to the quadric  $Q$ .

Let  $\mathcal{G} = \mathcal{G}(\mathbb{R}^{s,t})$ , where  $n = s + t$ , and let  $T$  denote a self-adjoint outermorphism from  $\mathcal{G}$  to  $\mathcal{G}$ , i.e.  $T = T^*$ . Consider the map  $f : \mathcal{G} \rightarrow \mathbb{R}$ ,  $f(x) := x * T(x)$  and form<sup>12</sup>

$$Q := \{x \in \mathcal{G} : f(x) = 0\}, \quad \text{and} \quad Q^m := \mathcal{G}^m \cap Q.$$

We then have  $f(x+h) = f(x) + 2T(x) * h + h * T(h)$  so that  $f'(x)$ , the derivative of  $f$  at the point  $x \in \mathcal{G}$ , is given by the linear map

$$\begin{aligned} \mathcal{G} &\rightarrow \mathbb{R} \\ h &\mapsto 2T(x) * h \end{aligned}$$

The *tangent space*  $Q_x$  to the surface  $Q$  at the point  $x$  is then defined as the linear space

$$Q_x := \ker f'(x) \subseteq \mathcal{G}.$$

We also define  $Q_x^m := \mathcal{G}^m \cap Q_x$ , which is the tangent space of  $Q^m$  at  $x$ .

**Definition 3.4.** The *polar* to  $x$  with respect to  $Q$  is defined by

$$\text{Pol}_Q(x) := T(x)^c = T(x)I^{-1}.$$

We note in particular that  $\overline{\text{Pol}_Q(x)} = \left(\overline{T(x)}\right)^\perp$  if  $x \in \mathcal{B}^\bullet$  is a nonzero blade.

We have a number of results regarding the polar.

**Theorem 3.11.** *If  $x \in Q^1$  and  $h_1, \dots, h_k \in Q_x^1$  then  $x \wedge h_1 \wedge \dots \wedge h_k \in Q^{k+1}$ , i.e.*

$$T(x \wedge h_1 \wedge \dots \wedge h_k) * (x \wedge h_1 \wedge \dots \wedge h_k) = 0.$$

*Proof.* We have

$$(T(x) \wedge T(h_1) \wedge \dots \wedge T(h_k)) * (h_k \wedge \dots \wedge h_1 \wedge x) = \det \begin{bmatrix} T(x) * x & \dots \\ T(x) * h_1 & \dots \\ \vdots & \\ T(x) * h_k & \dots \end{bmatrix}$$

which is zero since  $f(x) = 0$  and  $2T(x) * h_j = f'(x)(h_j) = 0$  for  $j = 1, \dots, k$ .  $\square$

<sup>12</sup>Note that  $f$  is actually a homogeneous quadratic polynomial in  $2^n$  variables.

If  $x \in Q^m$  then its polar  $\text{Pol}_Q(x)$  is an element of  $Q^{n-m}$  because of the following

**Theorem 3.12.** *We have*

$$T(x)^c * T(T(x)^c) = \tau \det T \ x * T(x),$$

where  $\tau = (-1)^{mn-m} I^2$  and  $x \in \mathcal{G}^m$ .

*Proof.* Assume that  $x \in \mathcal{G}^m$  and recall that

$$T^* \circ T^c = (\det T) \text{id}$$

holds for all outermorphisms  $T$ . When  $T^* = T$  we then obtain

$$\begin{aligned} T(x)^c * T(T(x)^c) &= (T(x)I^{-1}) * T^*(T^c(xI^{-1})) \\ &= (T(x)I^{-1}) * ((\det T)xI^{-1}) = \det T \langle T(x)I^{-1}xI^{-1} \rangle_0 \\ &= (-1)^{m(n-m)} I^{-2} \det T \langle T(x)x \rangle_0, \end{aligned}$$

which proves the theorem.  $\square$

In other words,  $f(\text{Pol}_Q(x)) = \tau \det T f(x)$ . We also have a reciprocity theorem:

**Theorem 3.13.** *If  $x, y \in \mathcal{G}^m$  then*

$$x \wedge T(y)^c = y \wedge T(x)^c.$$

*Proof.* From Proposition 2.14 we have

$$x \wedge T(y)^c = (x \lrcorner T(y))^c = (x * T(y))^c = (y * T(x))^c = (y \lrcorner T(x))^c = y \wedge T(x)^c,$$

where we used that  $T(x), T(y) \in \mathcal{G}^m$ .  $\square$

Finally, we have the following duality theorem:

**Theorem 3.14.**  $(T(x \wedge y))^c = I^2 T(x)^c \vee T(y)^c$ .

*Proof.* This follows immediately from the definition of the meet product and the fact that  $T$  is an outermorphism.  $\square$

---

**Exercise 3.18.** Interpret the above theorems geometrically in the case  $\mathbb{P}^2$ .

## 4 Discrete geometry

In this section we consider applications of Clifford algebras in discrete geometry and combinatorics. We also extend the definition of Clifford algebra and geometric algebra to infinite sets and infinite-dimensional vector spaces.

### 4.1 Simplicial complexes

A *simplicial complex*  $K$  on a finite set  $V$  is a subset of  $\mathcal{P}(V)$  which is closed under taking subsets, i.e. such that for all  $A, B \in \mathcal{P}(V)$

$$A \subseteq B \in K \quad \Rightarrow \quad A \in K.$$

In the following, let  $\mathcal{C}l(V)$  denote the Clifford algebra over the set  $V$ , where  $v^2 = 1, \forall v \in V$ , and where we use  $R = \mathbb{Z}$  as scalars. Define the *chain group*  $C(K) := \bigoplus_K \mathbb{Z}$  and note that  $C(K)$  is an abelian subgroup of  $\mathcal{C}l(V)$  (w.r.t addition). The sets  $A \in K$  with signs are called (*oriented*) *simplices* and  $|A| - 1$  is the *dimension* of the simplex  $A$ . Hence, we decompose  $C(K)$  into simplices of varying dimension,

$$C(K) = \bigoplus_{d \geq -1} C_d(K), \quad \text{where} \quad C_d(K) := \bigoplus_{\substack{A \in K \\ |A| - 1 = d}} \mathbb{Z}.$$

An element of  $C_d(K) \subseteq \mathcal{C}l^{d+1}(V)$  is called a *d-chain*. Note that the Clifford algebra structure of  $\mathcal{C}l(V)$  handles the orientation of the simplices, so that e.g. a 1-simplex  $v_0 v_1$ , thought of as a line segment directed from the point  $v_0$  to  $v_1$ , expresses its orientation as  $v_0 v_1 = -v_1 v_0$ .

Let  $s_V := \sum_{v \in V} v \in \mathcal{C}l^1(V)$  and define the *boundary map*  $\partial_V$  by

$$\begin{aligned} \partial_V : \mathcal{C}l(V) &\rightarrow \mathcal{C}l(V), \\ x &\mapsto \partial_V(x) := s_V \lrcorner x. \end{aligned}$$

Then  $\partial_V^2 = 0$ , since

$$\partial_V \circ \partial_V(x) = s_V \lrcorner (s_V \lrcorner x) = (s_V \wedge s_V) \lrcorner x = 0, \quad (4.1)$$

and the action of  $\partial_V$  on a  $k$ -simplex  $v_0 v_1 \dots v_k$  is given by

$$\partial_V(v_0 v_1 \dots v_k) = \sum_{i=0}^k v_i \lrcorner (v_0 v_1 \dots v_k) = \sum_{i=0}^k (-1)^i v_0 v_1 \dots \check{v}_i \dots v_k.$$

Note that  $C(K)$  is invariant under the action of  $\partial_V$  by the definition of a simplicial complex. We also have, for  $x, y \in \mathcal{C}l$ , (see Exercise 2.19)

$$\begin{aligned} \partial_V(xy) &= \partial_V(x)y + x^* \partial_V(y), \\ \partial_V(x \wedge y) &= \partial_V(x) \wedge y + x^* \wedge \partial_V(y), \\ \partial_V : \mathcal{C}l^m(V) &\rightarrow \mathcal{C}l^{m-1}(V), \\ \partial_V : C_d(K) &\rightarrow C_{d-1}(K). \end{aligned}$$

**Example 4.1.** Consider a set of three vertices,  $V = \{v_1, v_2, v_3\}$ , and a simplicial complex  $K = \{\emptyset, v_1, v_2, v_3, v_1 v_2, v_2 v_3\}$ . An example of a 1-chain is  $x = v_1 v_2 + v_2 v_3$ , with boundary  $\partial_V x = v_2 - v_1 + v_3 - v_2 = v_3 - v_1$ . Applying  $\partial_V$  again, we explicitly obtain  $\partial_V^2 x = 1 - 1 = 0$ . See Figure 4.1.




Figure 4.1: A 1-chain with nontrivial boundary in a simplicial complex  $K$ .

#### 4.1.1 Chain maps and complex morphisms

Let  $K$  and  $L$  be simplicial complexes on the finite sets  $V$  and  $W$ , respectively. A grade preserving  $\mathbb{Z}$ -linear map  $F : C(K) \rightarrow C(L)$  is called a *chain map* if the following diagram commutes

$$\begin{array}{ccc} C(K) & \xrightarrow{F} & C(L) \\ \partial_V \downarrow & & \downarrow \partial_W \\ C(K) & \xrightarrow{F} & C(L) \end{array}$$

i.e. if  $\partial_W \circ F = F \circ \partial_V$ .

A natural way to obtain a chain map is through the following construction. Let  $f : V \rightarrow W$  be a map which takes simplices of  $K$  to simplices of  $L$ , i.e. such that

$$L \ni f(A) := \{f(a) \in W : a \in A\}$$

whenever  $A \in K$ . Then  $f$  is called a *complex morphism*.

We can extend a complex morphism  $f$  to an outermorphism  $f_\wedge : Cl(V) \rightarrow Cl(W)$  such that  $f_\wedge : C(K) \rightarrow C(L)$ .

**Lemma 4.1.**  $f_\wedge^*(s_W) = s_V$

*Proof.* For any  $v \in V$  we have

$$v * f_\wedge^*(s_W) = f_\wedge(v) * s_W = 1,$$

while  $x * f_\wedge^*(s_W) = 0$  for any  $x \in Cl^{d \neq 1}(V)$  because outermorphisms preserve grades. From this we conclude that  $f_\wedge^*(s_W) = s_V$ .  $\square$

**Corollary.**  $f_\wedge$  is a chain map.

*Proof.* Consider the diagram

$$\begin{array}{ccc} C(K) & \xrightarrow{f_\wedge} & C(L) \\ \partial_V \downarrow & & \downarrow \partial_W \\ C(K) & \xrightarrow{f_\wedge} & C(L) \end{array}$$

By Theorem 3.6, we have

$$\partial_W f_\wedge x = s_W \lrcorner (f_\wedge x) = f_\wedge((f_\wedge^* s_W) \lrcorner x) = f_\wedge(s_V \lrcorner x) = f_\wedge \partial_V x$$

for all  $x \in C(K)$ .  $\square$






Figure 4.2: A triangulation of the triangle  $ABC$ , with a labeling of the nodes of its sides. Positive orientation means with respect to the plane of the page, as in the triangle on the right.

#### 4.1.2 An index theorem and Sperner's lemma

Consider an arbitrary linear map  $F : Cl(V) \rightarrow Cl(W)$  such that  $F\partial_V = \partial_W F$  (typically the chain map  $f_\wedge$  induced by some complex morphism  $f$ ) and define two linear mappings, the *content* and the *index*, by

$$\begin{aligned} \text{Cont} : Cl(V) &\rightarrow \mathbb{Z} \\ x &\mapsto W * (Fx), \quad \text{and,} \\ \text{Ind} : Cl(V) &\rightarrow \mathbb{Z} \\ x &\mapsto (Wa) * (F\partial_V x), \end{aligned}$$

where  $a$  is an arbitrary but fixed element in  $W$ .

**Theorem 4.2.** *We have  $\text{Cont} = \text{Ind}$ , and in particular,  $\text{Ind}$  does not depend on the choice of  $a \in W$ .*

*Proof.* Since  $F$  is a chain map, we have for all  $x \in Cl(V)$

$$\begin{aligned} Wa * F\partial_V x &= Wa * \partial_W Fx = Wa * (s_W \lrcorner Fx) \\ &= (Wa \wedge s_W) * Fx = W * Fx, \end{aligned}$$

which proves the theorem.  $\square$

As an application we will prove Sperner's lemma (in the plane for simplicity).

**Theorem 4.3** (Sperner's lemma in the plane). *Consider a triangulation of a triangle  $ABC$  and label every node in the triangulation with one of the symbols  $a, b$  or  $c$  in such a manner that  $A, B, C$  is labeled  $a, b, c$  respectively and so that every node on the side  $AB$  is labeled with  $a$  or  $b$ , every node on side  $BC$  is labeled with  $b$  or  $c$ , and those on  $AC$  with  $a$  or  $c$ . A triangle in the triangulation is called complete if its nodes have distinct labels (i.e.  $a, b$  and  $c$ ). Sperner's lemma states that there exists an odd number of complete triangles in the triangulation.*

*Proof.* To prove this, let  $V$  denote the set of nodes in the triangulation. To each triangle in the triangulation we associate the simplex  $v \wedge v' \wedge v''$  in  $Cl(V)$ , where the order is chosen so that  $v, v', v''$  is positively oriented as in Figure 4.2. Let  $x$  denote the sum of all these simplices.




Figure 4.3: The nodes on the segment  $BC$  of a triangulation of  $ABC$ .

Now, we let  $f : V \rightarrow W := \{a, b, c\}$  be a map defined by the labeling, and extend it to an outermorphism (chain map)  $f_\wedge = F : \mathcal{Cl}(V) \rightarrow \mathcal{Cl}(W)$ . We now see that the triangulation has a complete triangle if  $\text{Cont } x \neq 0$ , i.e. if  $W * Fx \neq 0$ . But by the index theorem,  $\text{Cont} = \text{Ind}$ , so we only have to prove that  $\text{Ind } x = Wa * F(\partial_V x) \neq 0$ , or equivalently that  $abca * \partial_W Fx = bc * \partial_W Fx \neq 0$ . Note that  $\partial_V x$  is just the boundary chain of the triangulation. Hence, the only terms in  $\partial_W Fx$  we need to consider are those corresponding to the nodes on the segment  $BC$ . But since  $b \wedge b = c \wedge c = 0$  we conclude that (see Figure 4.3)

$$\text{Ind } x = bc * (b \wedge c + c \wedge b + b \wedge c + c \wedge b + \dots + b \wedge c) = bc * bc = -1,$$

so that  $W * Fx = -1$ . This proves that there is an odd number of complete triangles.  $\square$

By using induction on the dimension it is now straightforward to prove Sperner's lemma in higher dimensions.

---

**Exercise 4.1.** Formulate and prove Sperner's lemma in general.

### 4.1.3 Homology

Another example where chain maps are useful is in homology theory, which is an invaluable tool in e.g. topology.

Again, let  $K$  be a complex on  $V$ , and denote the boundary map on  $V$  by  $\partial_V$ . We define the *cycles*  $Z(K) = \ker \partial_V$  and the *boundaries*  $B(K) = \text{im } \partial_V$ .

Since  $\partial_V^2 = 0$  we have  $B(K) \subseteq Z(K)$ . Hence, we can introduce the *homology*

$$H(K) = Z(K)/B(K).$$

We also define  $d$ -cycles,  $d$ -boundaries, and the corresponding  $d$ th homology group,

$$\begin{aligned} Z_d(K) &= Z(K) \cap C_d(K), \\ B_d(K) &= B(K) \cap C_d(K), \\ H_d(K) &= Z_d(K)/B_d(K). \end{aligned}$$

The fundamental theorem of finitely generated abelian groups (see e.g. [25]) states that every finitely generated abelian group is a direct sum of cyclic subgroups. Thus,

$$H_d(K) = F_d \oplus T_d,$$

where  $F_d \cong \mathbb{Z}^r$  for some  $r \in \mathbb{N}$  called the *rank* of  $H_d(K)$  or the *Betti number* denoted  $\beta_d$ , and  $T_d$  is the *torsion part*: for all  $t \in T_d$  there is a positive integer  $n$  such that  $nt = 0$ .

**Theorem 4.4.** *Every chain map  $F : C(K) \rightarrow C(L)$  induces a homomorphism  $F_* : H(K) \rightarrow H(L)$  on the homology level. Moreover,  $F_* : H_d(K) \rightarrow H_d(L)$  for all  $d \geq 0$ .*

*Proof.* Consider the commutative diagram

$$\begin{array}{ccc} C(K) & \xrightarrow{\partial_V} & C(K) \\ F \downarrow & & \downarrow F \\ C(L) & \xrightarrow{\partial_W} & C(L) \end{array}$$

We have for any  $b \in B(K)$  that  $b = \partial_V x$ , for some  $x \in V$ , so that

$$Fb = F\partial_V x = \partial_W Fx \in B(L).$$

Also,  $z \in Z(K)$  implies  $\partial_W Fz = F\partial_V z = F0 = 0$ , and  $Fz \in Z(L)$ . Hence, the map

$$\begin{array}{ccc} H(K) & \rightarrow & H(L) \\ z + B(K) & \mapsto & Fz + B(L) \end{array}$$

is well defined. Lastly,  $F_* : H_d(K) \rightarrow H_d(L)$  follows since  $F$  is grade preserving.  $\square$

*Remark.* From Exercise 2.22 and (4.1) it follows that a more general boundary map  $\partial_p := s_p \lrcorner$ , with  $s_p := \sum_{A \in K, |A|=p} A$  and  $p$  odd, would also give rise to a homology on  $K$ .

## 4.2 The number of spanning trees in a graph

We consider a finite connected simple graph (i.e. no loops or multi-edges)  $G$  with a set of edges  $E$  and vertices  $V$ . A subgraph  $T$  of  $G$  is called a *spanning tree* if it is connected, has no cycles, and contains every vertex in  $V$ . This is equivalent with having no cycles and having  $|V| - 1$  edges. In this subsection we will use Clifford algebra to prove a well known result about the number of spanning trees in  $G$  (see e.g. [35] and references therein for recent generalizations).

Let us first choose a total ordering on  $V$ , denoted  $\prec$ . We introduce the Clifford algebras  $Cl(V)$  and  $Cl(E)$  with signatures 1. If  $e = \{v', v\} \in E$  with  $v' \prec v$ , then we define  $\delta(e) := v - v'$ , and extend  $\delta$  to an outermorphism

$$\delta : Cl(E) \rightarrow Cl(V).$$

We will need two simple lemmas.

**Lemma 4.5.** *If the subgraph of  $G$  induced by  $F \subseteq E$  contains a cycle  $v_0 v_1 \dots v_k v_0$ , then  $\delta F = 0$ .*

*Proof.* The identity

$$(v_0 - v_1) + (v_1 - v_2) + \dots + (v_{k-1} - v_k) + (v_k - v_0) = 0$$

implies  $(v_0 - v_1) \wedge (v_1 - v_2) \wedge \dots \wedge (v_k - v_0) = 0$ , and  $\delta F = 0$ .  $\square$

**Lemma 4.6.** *If  $F \subseteq E$ ,  $\delta F \neq 0$  and  $|F| = |V| - 1$ , then there exists an  $\epsilon = \pm 1$ , such that  $v \wedge \delta F = \epsilon V$  for every  $v$  in  $V$ .*

*Proof.* Note that, if  $e = \{v, v'\}$  is an edge in  $F$ , then we have  $(v - v') \wedge \delta F = \delta(e \wedge F) = 0$ . Then, since the conditions imply that the subgraph induced by  $F$  is connected and contains every vertex of  $V$ , we conclude that  $v \wedge \delta F$  does not depend on  $v \in V$ .

Fix a vertex  $v$  and let  $v_1, v_2, \dots, v_s$  be the neighbourhood of  $v$  in the subgraph induced by  $F$ , i.e. all the vertices in  $V$  such that  $e_1 := \{v, v_1\}, e_2 := \{v, v_2\}, \dots, e_s := \{v, v_s\}$  are edges in  $F$ . Denote the set of remaining edges in  $F$  by  $F'$ . Then

$$v \wedge (v_1 - v) \wedge (v_2 - v) \wedge \dots \wedge (v_s - v) = v \wedge v_1 \wedge v_2 \wedge \dots \wedge v_s,$$

so that, up to some signs  $\epsilon_k$ ,

$$v \wedge \delta F = \epsilon_1 v \wedge \delta(e_1 e_2 \dots e_s) \wedge \delta F' = \epsilon_1 \epsilon_2 v \wedge v_2 \wedge \dots \wedge v_s \wedge v_1 \wedge \delta F'.$$

By repeating this argument with  $v_1$  in place of  $v$ , and so on, we obviously obtain  $v \wedge \delta F = \epsilon V$ , for some  $\epsilon \in \{-1, 1\}$  independent of  $v$ . This proves the lemma.  $\square$

Before we state the theorem we define the outermorphism ‘‘Laplace of  $G$ ’’ by

$$\Delta = \delta \circ \delta^* : \mathcal{Cl}(V) \rightarrow \mathcal{Cl}(V),$$

where  $\delta^* : \mathcal{Cl}(V) \rightarrow \mathcal{Cl}(E)$  is defined by the property  $\delta^* x * y = x * \delta y$  for all  $x \in \mathcal{Cl}(V)$ ,  $y \in \mathcal{Cl}(E)$ . The adjugate  $\Delta^{\text{adj}}$  of  $\Delta$  was earlier defined by  $\Delta^{\text{adj}} = \Delta^{*\text{c}}$ , where  $\Delta^* = (\delta \circ \delta^*)^* = \delta \circ \delta^* = \Delta$  and  $\Delta^{\text{c}}(x) = \Delta(xV)V^{-1}$ . Hence,

$$\Delta^{\text{adj}}(x) = \Delta(xV)V^\dagger \quad \forall x \in \mathcal{Cl}(V).$$

**Theorem 4.7** (Kirchoff’s matrix tree theorem). *For all  $u, v \in V$  we have*

$$u * \Delta^{\text{adj}} v = N,$$

where  $N$  denotes the number of spanning trees in  $G$ .

*Proof.* In the proof we will use the following simple rules for  $v \in V$ ,  $x, y, z \in \mathcal{Cl}(V)$ :

- $\delta(F^\dagger) = (\delta F)^\dagger$
- $x * y = \langle xy \rangle_0 = \langle (xy)^\dagger \rangle_0 = \langle y^\dagger x^\dagger \rangle_0 = y^\dagger * x^\dagger = y * x$
- $x * (yz) = \langle xyz \rangle_0 = \langle zxy \rangle_0 = (xy) * z$
- $vx = v \lrcorner x + v \wedge x$

We have

$$u * \Delta^{\text{adj}} v = u * \delta \circ \delta^*(vV) V^\dagger = V^\dagger u * \delta \circ \delta^*(vV) = \delta^*(V^\dagger u) * \delta^*(vV).$$

Denote  $n = |V|$ . Now, since  $\delta^*$  is grade preserving, and  $V^\dagger u$  and  $vV$  are of grade  $n - 1$ , we conclude that  $\delta^*(V^\dagger u)$  and  $\delta^*(vV)$  also have grade  $n - 1$ . By expanding these in a basis of subsets of  $E$  we obtain

$$\begin{aligned} u * \Delta^{\text{adj}} v &= \left( \sum_{\substack{F \subseteq E \\ |F|=n-1}} (\delta^*(V^\dagger u) * F) F^\dagger \right) * \left( \sum_{\substack{F \subseteq E \\ |F|=n-1}} (\delta^*(vV) * F^\dagger) F \right) \\ &= \sum_{\substack{F \subseteq E \\ |F|=n-1}} ((V^\dagger u) * \delta F) ((vV) * \delta F^\dagger) \\ &= \sum_{\substack{F \subseteq E \\ |F|=n-1}} (V^\dagger * (u \delta F)) (V^\dagger * (v \delta F)) \\ &= \sum_{\substack{F \subseteq E \\ |F|=n-1}} \langle V^\dagger (u \wedge \delta F) \rangle_0 \langle V^\dagger (v \wedge \delta F) \rangle_0 \\ &= \sum_{\substack{F \text{ spanning tree}}} \langle V^\dagger \epsilon_F V \rangle_0 \langle V^\dagger \epsilon_F V \rangle_0 = N. \end{aligned}$$

□

*Remark.* Note that the boundary map  $\delta$  on edges is induced by the boundary map  $\partial_V$  defined previously. In general, we can consider the induced structure

$$\begin{array}{ccccc} \mathcal{Cl}(\binom{V}{k}) & \leftrightarrow & \mathcal{Cl}^1(\binom{V}{k}) & \leftrightarrow & \mathcal{Cl}^k(V) \\ \downarrow \delta & & \delta \downarrow & & \downarrow \partial_V \\ \mathcal{Cl}(\binom{V}{k-1}) & \leftrightarrow & \mathcal{Cl}^1(\binom{V}{k-1}) & \leftrightarrow & \mathcal{Cl}^{k-1}(V) \end{array}$$

(or similarly for  $\partial_p$ ), where  $\binom{V}{k}$  denotes the set of  $k$ -subsets of  $V$ .

### 4.3 Fermionic operators

Here we consider a finite set  $E = \{e_1, \dots, e_n\}$  and its corresponding Clifford algebra with signature 1 and scalars  $\mathbb{F}$ , which we denote by  $\mathcal{F} = \mathcal{Cl}(E, \mathbb{F}, 1)$ . Define the *fermion creation and annihilation operators* acting on  $\psi \in \mathcal{F}$

$$c_i^\dagger(\psi) := e_i \wedge \psi, \quad c_i(\psi) := e_i \lrcorner \psi. \quad (4.2)$$

Again, by the properties of the inner and outer products, we find, for any  $\psi \in \mathcal{F}$ ,

$$\begin{aligned} c_i c_j(\psi) &= (e_i \wedge e_j) \lrcorner \psi = -c_j c_i(\psi), \\ c_i^\dagger c_j^\dagger(\psi) &= e_i \wedge e_j \wedge \psi = -c_j^\dagger c_i^\dagger(\psi), \\ c_i c_j^\dagger(\psi) &= e_i \lrcorner (e_j \wedge \psi) = \delta_{ij} \psi - e_j \wedge (e_i \lrcorner \psi) = \delta_{ij} \psi - c_j^\dagger c_i(\psi). \end{aligned}$$

We summarize these properties by the *anticommutation relations*

$$\{c_i, c_j\} = 0, \quad \{c_i, c_j^\dagger\} = \delta_{ij}, \quad \{c_i^\dagger, c_j^\dagger\} = 0, \quad (4.3)$$

where  $\{A, B\} := AB + BA$  is called the *anticommutator*.

With  $\mathbb{F} = \mathbb{C}$ , the inner product  $\langle \phi, \psi \rangle_{\mathcal{F}} := \overline{\phi}^\dagger * \psi$  ( $\overline{\phantom{x}}$  denotes the complex conjugate acting on the scalar coefficients, and  $\psi^\dagger$  is the reverse as usual) makes  $\mathcal{F}$  into a complex  $2^n$ -dimensional Hilbert space. We also find that (see Exercise 2.15)

$$\langle \phi, c_i \psi \rangle_{\mathcal{F}} = \overline{\phi}^\dagger * (e_i \lrcorner \psi) = \overline{\phi^\dagger \wedge e_i} * \psi = \langle c_i^\dagger \phi, \psi \rangle_{\mathcal{F}},$$

so that  $c_i^\dagger$  is the adjoint of  $c_i$  with respect to this inner product. In quantum mechanics, this space  $\mathcal{F}$  is usually called the *fermionic Fock space of  $n$  particles* and the physical interpretation is that each of the operators  $c_i^\dagger$  and  $c_i$  creates resp. annihilates a particle of type/location  $i$ . Because  $(c_i^\dagger)^2 = 0$ , there cannot be more than one particle of the same type at the same location<sup>13</sup>. Any multivector  $\psi \in \mathcal{F}$ , here called a *state*, can be written as a linear combination

$$\psi = \left( \alpha_0 + \sum_i \alpha_i c_i^\dagger + \sum_{i < j} \alpha_{ij} c_i^\dagger c_j^\dagger + \dots + \alpha_{1\dots n} c_1^\dagger \dots c_n^\dagger \right) \psi_0 \quad (4.4)$$

of particle creations acting on an empty, zero-particle or *vacuum*, state  $\psi_0 := 1$  (note that  $c_i \psi_0 = 0$  for all  $i$ , so it is not possible to remove any more particles from this state). The scalar coefficients  $\alpha_i$  determine the probabilities  $p_i = |\alpha_i|^2 / \|\psi\|_{\mathcal{F}}^2$  of measuring the physical state  $\psi$  in a definite particle configuration  $e_i = c_{i_1}^\dagger c_{i_2}^\dagger \dots c_{i_k}^\dagger \psi_0$ . There is also a *number operator*,  $n_F := \sum_i c_i^\dagger c_i$ , which simply counts the number of particles present in a state.

---

**Exercise 4.2.** Show that  $n_F \psi = k \psi$  for  $\psi \in \mathcal{F}_k := \mathcal{Cl}^k(E, \mathbb{C}, 1)$ , by using only the anticommutation relations (4.3) and the representation (4.4), and in an alternative way by using only the properties of the inner and outer products.

## 4.4 Supersymmetric quantum mechanics on a lattice

To be written...

## 4.5 Infinite-dimensional Clifford algebra

This far we have only defined the Clifford algebra  $\mathcal{Cl}(X, R, r)$  of a *finite* set  $X$ , resulting in the *finite-dimensional* geometric algebra  $\mathcal{G}(V)$  whenever  $R$  is a field. In order for this combinatorial construction to qualify as a complete generalization of  $\mathcal{G}$ , we would like to be able to define the corresponding Clifford algebra of an infinite-dimensional vector space, something which was possible for  $\mathcal{G}$  in Definition 2.2.

The treatment of  $\mathcal{Cl}$  in the previous subsections is in a form which easily generalizes to an infinite  $X$ . Reconsidering Definition 2.4, we now have two possibilities: either we consider the set  $\mathcal{P}(X)$  of *all* subsets of  $X$ , or just the

---

<sup>13</sup>This property characterizes *fermionic* particles in contrast to *bosonic* particles, which can appear at the same location and which satisfy symmetry instead of antisymmetry, being represented by commutators instead of anticommutators.

set  $\mathcal{F}(X)$  of all *finite* subsets. With the tensor-algebraic definition in mind, we choose to first consider only  $\mathcal{F}(X)$ , and then the generalized case  $\mathcal{P}(X)$  in the next subsection. We therefore define, for an arbitrary set  $X$ , ring  $R$ , and signature  $r: X \rightarrow R$ ,

$$\mathcal{Cl}_{\mathcal{F}}(X, R, r) := \bigoplus_{\mathcal{F}(X)} R. \quad (4.5)$$

Elements in  $\mathcal{Cl}_{\mathcal{F}}$  are then finite linear combinations of finite subsets of  $X$ .

Our problem now is to define a Clifford product for  $\mathcal{Cl}_{\mathcal{F}}$ . This can be achieved just as in the finite case if only we can find a map  $\tau: \mathcal{F}(X) \times \mathcal{F}(X) \rightarrow R$  satisfying the conditions in Lemma 2.6. We show below that this is possible.

We call a map  $\tau: \mathcal{F}(X) \times \mathcal{F}(X) \rightarrow R$  *grassmannian on  $X$*  if it satisfies (i)-(v) in Lemma 2.6, with  $\mathcal{P}(X)$  replaced by  $\mathcal{F}(X)$ .

**Theorem 4.8.** *For any given  $X, R, r$  there exists a grassmannian map on  $\mathcal{F}(X)$ .*

*\*Proof.* We know that there exists such a map for any finite  $X$ . Let  $Y \subseteq X$  and assume  $\tau': \mathcal{F}(Y) \times \mathcal{F}(Y) \rightarrow R$  is grassmannian on  $Y$ . If there exists  $z \in X \setminus Y$  we can, by proceeding as in the proof of Lemma 2.6, extend  $\tau'$  to a grassmannian map  $\tau: \mathcal{F}(Y \cup \{z\}) \times \mathcal{F}(Y \cup \{z\}) \rightarrow R$  on  $Y \cup \{z\}$  such that  $\tau|_{\mathcal{F}(Y) \times \mathcal{F}(Y)} = \tau'$ .

We will now use *transfinite induction*, or the *Hausdorff maximality theorem*<sup>14</sup>, to prove that  $\tau$  can be extended to all of  $\mathcal{F}(X) \subseteq \mathcal{P}(X)$ . Note that if  $\tau$  is grassmannian on  $Y \subseteq X$  then  $\tau$  is also a relation  $\tau \subseteq \mathcal{P}(X) \times \mathcal{P}(X) \times R$ . Let

$$\mathcal{H} := \left\{ (Y, \tau) \in \mathcal{P}(X) \times \mathcal{P}(\mathcal{P}(X) \times \mathcal{P}(X) \times R) : \tau \text{ is grassmannian on } Y \right\}. \quad (4.6)$$

Then  $\mathcal{H}$  is partially ordered by

$$(Y, \tau) \leq (Y', \tau') \quad \text{iff} \quad Y \subseteq Y' \quad \text{and} \quad \tau'|_{\mathcal{F}(Y) \times \mathcal{F}(Y)} = \tau. \quad (4.7)$$

By the Hausdorff maximality theorem, there exists a maximal totally ordered “chain”  $\mathcal{K} \subseteq \mathcal{H}$ . Put  $Y^* := \bigcup_{(Y, \tau) \in \mathcal{K}} Y$ . We want to define a grassmannian map  $\tau^*$  on  $Y^*$ , for if we succeed in that, we find  $(Y^*, \tau^*) \in \mathcal{H} \cap \mathcal{K}$  and can conclude that  $Y^* = X$  by maximality and the former result.

Take finite subsets  $A$  and  $B$  of  $Y^*$ . Each of the finite elements in  $A \cup B$  lies in some  $Y$  such that  $(Y, \tau) \in \mathcal{K}$ . Therefore, by the total ordering of  $\mathcal{K}$ , there exists one such  $Y$  containing  $A \cup B$ . Put  $\tau^*(A, B) := \tau(A, B)$ , where  $(Y, \tau)$  is this chosen element in  $\mathcal{K}$ .  $\tau^*$  is well-defined since if  $A \cup B \subseteq Y$  and  $A \cup B \subseteq Y'$  where  $(Y, \tau), (Y', \tau') \in \mathcal{K}$  then  $Y \subseteq Y'$  or  $Y' \subseteq Y$  and  $\tau, \tau'$  agree on  $(A, B)$ . It is easy to verify that this  $\tau^*$  is grassmannian on  $Y^*$ , since for each  $A, B, C \in \mathcal{F}(X)$  there exists  $(Y, \tau) \in \mathcal{K}$  such that  $A \cup B \cup C \subseteq Y$ .  $\square$

We have shown that there exists a map  $\tau: \mathcal{F}(X) \times \mathcal{F}(X) \rightarrow R$  with the properties in Lemma 2.6. We can then define the Clifford product on  $\mathcal{Cl}_{\mathcal{F}}(X)$  as usual by  $AB := \tau(A, B)A \Delta B$  for  $A, B \in \mathcal{F}(X)$  and linear extension. Since only finite subsets are included, most of the previous constructions for finite-dimensional  $\mathcal{Cl}$  carry over to  $\mathcal{Cl}_{\mathcal{F}}$ . For example, the decomposition into graded subspaces remains but now extends towards infinity,

$$\mathcal{Cl}_{\mathcal{F}} = \bigoplus_{k=0}^{\infty} \mathcal{Cl}_{\mathcal{F}}^k. \quad (4.8)$$

<sup>14</sup>This theorem should actually be regarded as an axiom of set theory since it is equivalent to the Axiom of Choice.

Furthermore, Proposition 2.7 still holds, so the reverse and all other involutions behave as expected. On the other hand, note that we cannot talk about a pseudoscalar in this context.

Let us now see how  $\mathcal{Cl}_{\mathcal{F}}$  can be applied to the setting of an infinite-dimensional vector space  $V$  over a field  $\mathbb{F}$  and with a quadratic form  $q$ . By the Hausdorff maximality theorem one can actually find a (necessarily infinite) orthogonal basis  $E$  for this space in the sense that any vector in  $V$  can be written as a finite linear combination of elements in  $E$  and that  $\beta_q(e, e') = 0$  for any pair of disjoint elements  $e, e' \in E$ . We then have

$$\mathcal{G}(V, q) \cong \mathcal{Cl}_{\mathcal{F}}(E, \mathbb{F}, q|_E), \quad (4.9)$$

which is proved just like in the finite-dimensional case, using Theorem 2.1 and the proof and corollary of Theorem 2.8.

## 4.6 Embedded discrete geometry

Let us consider an application of the infinite-dimensional Clifford algebra  $\mathcal{Cl}_{\mathcal{F}}$ . Given a vector space  $V$ , define the *full simplicial complex algebra*

$$\mathcal{C}(V) := \mathcal{Cl}_{\mathcal{F}}(V, R, 1), \quad (4.10)$$

where we forget about the vector space structure of  $V$  and treat individual points  $\dot{v} \in V$  as orthogonal basis 1-vectors in  $\mathcal{Cl}_{\mathcal{F}}^1$  with  $\dot{v}^2 = 1$ . The dot indicates that we think of  $v$  as a point rather than a vector. A basis  $(k+1)$ -blade in  $\mathcal{C}(V)$  consists of a product  $\dot{v}_0 \dot{v}_1 \dots \dot{v}_k$  of individual points and represents a (possibly degenerate) oriented  $k$ -simplex embedded in  $V$ . This simplex is given by the convex hull

$$\text{Conv}\{v_0, v_1, \dots, v_k\} := \left\{ \sum_{i=0}^k \alpha_i v_i \in V : \alpha_i \geq 0, \sum_{i=0}^k \alpha_i = 1 \right\}. \quad (4.11)$$

Hence, an arbitrary element in  $\mathcal{C}(V)$  is a linear combination of simplices and can therefore represent a simplicial complex embedded in  $V$ . Restricting to a chosen embedded simplex  $K \subseteq \mathcal{F}(V)$ , we return to the context of the chain group  $C(K)$  introduced previously. However, the difference here is that the chain group of *any* simplicial complex properly embedded in  $V$  can be found as a subgroup of  $\mathcal{C}(V)$  – even infinite such complexes (composed of finite-dimensional simplices).

We can define a boundary map for the full simplicial complex algebra,

$$\begin{aligned} \partial_V: \mathcal{C}(V) &\rightarrow \mathcal{C}(V), \\ \partial_V(x) &:= \sum_{\dot{v} \in V} \dot{v} \lrcorner x. \end{aligned}$$

Note that this is well-defined since only a finite number of points  $\dot{v}$  can be present in any fixed  $x$ . Also,  $\partial_V^2 = 0$  follows by analogy with  $s_V$ , or simply

$$\partial_V^2(x) = \sum_{\dot{u} \in V} \dot{u} \lrcorner \left( \sum_{\dot{v} \in V} \dot{v} \lrcorner x \right) = \sum_{\dot{u}, \dot{v} \in V} \dot{u} \lrcorner (\dot{v} \lrcorner x) = \sum_{\dot{u}, \dot{v} \in V} (\dot{u} \wedge \dot{v}) \lrcorner x = 0. \quad (4.12)$$



We can also assign a *geometric measure*  $\sigma$  to embedded simplices, by mapping a  $k$ -simplex to a corresponding  $k$ -blade in  $\mathcal{G}(V)$  representing the directed volume of the simplex. Define  $\sigma: \mathcal{C}(V) \rightarrow \mathcal{G}(V)$  by

$$\begin{aligned}\sigma(1) &:= 0, \\ \sigma(\dot{v}) &:= 1, \\ \sigma(\dot{v}_0 \dot{v}_1 \dots \dot{v}_k) &:= \frac{1}{k!} (v_1 - v_0) \wedge (v_2 - v_0) \wedge \dots \wedge (v_k - v_0),\end{aligned}$$

and extending linearly. One can verify that this is well-defined and that the geometric measure of a boundary is zero, i.e.  $\sigma \circ \partial_V = 0$ . One can take this construction even further and arrive at “discrete” equivalents of differentials, integrals and Stokes’ theorem. See [43] or [33] for more on this.

## 4.7 \*Generalized infinite-dimensional Clifford algebra

We define, for an arbitrary set  $X$ , ring  $R$ , and signature  $r: X \rightarrow R$ ,

$$Cl(X, R, r) := \bigoplus_{\mathcal{P}(X)} R. \quad (4.13)$$

Hence, elements of  $Cl(X)$  are finite linear combinations of (possibly infinite) subsets of  $X$ . The following theorem asserts that it is possible to extend  $\tau$  all the way to  $\mathcal{P}(X)$  even in the infinite case. We therefore have a Clifford product also on  $Cl(X)$ .

**Theorem 4.9.** *For any set  $X$  there exists a map  $|\cdot|_2: \mathcal{P}(\mathcal{P}(X)) \rightarrow \mathbb{Z}_2$  such that*

- i)  $|\mathcal{A}|_2 \equiv |\mathcal{A}| \pmod{2}$  for finite  $\mathcal{A} \subseteq \mathcal{P}(X)$ ,*
- ii)  $|\mathcal{A} \triangle \mathcal{B}|_2 = |\mathcal{A}|_2 + |\mathcal{B}|_2 \pmod{2}$*

*Furthermore, for any commutative ring  $R$  with unit, and signature  $r: X \rightarrow R$  such that  $r(X)$  is contained in a finite and multiplicatively closed subset of  $R$ , there exists a map  $\tau: \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow R$  such that properties (i)-(v) in Lemma 2.6 hold, plus*

$$vi) \tau(A, B) = (-1)^{|\binom{A}{2}|_2 + |\binom{B}{2}|_2 + |\binom{A \triangle B}{2}|_2} \tau(B, A) \quad \forall A, B \in \mathcal{P}(X).$$

Here,  $\binom{A}{n}$  denotes the set of all subsets of  $A$  with  $n$  elements. Note that for a finite set  $A$ ,  $|\binom{A}{n}| = \binom{|A|}{n}$  so that for example  $|\binom{A}{1}| = |A|$  (in general,  $\text{card} \binom{A}{1} = \text{card} A$ ) and  $|\binom{A}{2}| = \frac{1}{2}|A|(|A| - 1)$ . This enables us to extend the basic involutions  $\star$ ,  $\dagger$  and  $\square$  to infinite sets as

$$\begin{aligned}A^\star &:= (-1)^{|\binom{A}{1}|_2} A, \\ A^\dagger &:= (-1)^{|\binom{A}{2}|_2} A,\end{aligned}$$

and because  $|\binom{A \triangle B}{1}|_2 = |\binom{A}{1}|_2 + |\binom{B}{1}|_2 \pmod{2}$  still holds, we find that they satisfy the fundamental properties (2.17)-(2.19) for all elements of  $Cl(X)$ . The extra requirement (vi) on  $\tau$  was necessary here since we cannot use Proposition 2.7 for infinite sets. Moreover, we can no longer write the decomposition (4.8)

since it goes beyond finite grades. We do have even and odd subspaces, though, defined by

$$\mathcal{Cl}^\pm := \{x \in \mathcal{Cl} : x^* = \pm x\}, \quad (4.14)$$

and  $\mathcal{Cl}^+$  and  $\mathcal{Cl}_{\mathcal{F}}$  (with this  $\tau$ ) are both subalgebras of  $\mathcal{Cl}$ .

It should be emphasized that  $\tau$  needs not be zero on intersecting infinite sets (a rather trivial solution), but if e.g.  $r: X \rightarrow \{\pm 1\}$  we can also demand that  $\tau: \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow \{\pm 1\}$ . A proof of the first part of Theorem 4.9 is given in Appendix A.4.

Although this construction provides a way to handle combinatorics of infinite sets, we should emphasize that its applicability in the context of infinite-dimensional vector spaces appears limited. Let us sketch an intuitive picture of why this is so.

With a countable basis  $E = \{e_i\}_{i=1}^\infty$ , an infinite basis blade in  $\mathcal{Cl}$  could be thought of as an infinite product  $A = e_{i_1}e_{i_2}e_{i_3}\dots = \prod_{k=1}^\infty e_{i_k}$ . A change of basis to  $E'$  would turn each  $e \in E$  into a finite linear combination of elements in  $E'$ , e.g.  $e_j = \sum_k \beta_{jk}e'_k$ . However, this would require  $A$  to be an *infinite sum* of basis blades in  $E'$ , which is not allowed. Note that this is no problem in  $\mathcal{Cl}_{\mathcal{F}}$  since a basis blade  $A = \prod_{k=1}^N e_{i_k}$  is a finite product and the change of basis therefore results in a finite sum.

This completes our excursion to infinite-dimensional Clifford algebras (we refer to [10, 36, 47] and references therein for more on this topic, however we are not aware of other constructions of the above generality). In the following sections we will always assume that  $X$  is finite and  $V$  finite-dimensional.

## 5 Classification of real and complex geometric algebras

In this section we establish an extensive set of relations and isomorphisms between real and complex geometric algebras of varying signature. This eventually leads to an identification of these algebras as matrix algebras over  $\mathbb{R}$ ,  $\mathbb{C}$ , or the quaternions  $\mathbb{H}$ . The complete listing of such identifications is usually called the *classification* of geometric algebras.

### 5.1 A prototype example

Let us for completeness introduce the notation

$$\mathcal{G}(\mathbb{F}^{0,0}) := Cl(\emptyset, \mathbb{F}, \emptyset) = \mathbb{F}.$$

We then have  $\mathcal{G}(\mathbb{R}^0) = \mathbb{R}$ , and also recall that for example  $\mathcal{G}(\mathbb{R}^{0,1}) \cong \mathbb{C}$  (Example 2.6),  $\mathcal{G}(\mathbb{R}^{0,2}) \cong \mathbb{H}$  (Example 2.7), and  $\mathcal{G}(\mathbb{R}^2) \cong \mathbb{R}^{2 \times 2}$  (Exercise 2.5), all of which are indeed matrix algebras (and very simple cases of such). On the other hand, we claim that the algebra of the euclidean line is  $\mathcal{G}(\mathbb{R}^1) \cong \mathbb{R} \oplus \mathbb{R}$ , a slightly more complicated algebra. This is however a prototype case which also deserves to be studied in some detail since it turns out that all higher-dimensional algebras share essentially the same structure as the already mentioned ones.

Indeed, first note that  $\mathbb{R} \oplus \mathbb{R} \not\cong \mathbb{R} \otimes \mathbb{R}$ , namely their dimensions are different as  $\mathbb{R} \otimes \mathbb{R} \cong \mathbb{R}$ , although they formally share the same multiplication law (see Section A.1.7)

$$(a, b)(c, d) = (ac, bd), \quad a, b, c, d \in \mathbb{R}.$$

The isomorphism  $\mathcal{G}(\mathbb{R}^1) \cong \mathbb{R} \oplus \mathbb{R}$ , despite matching dimensionality, is not immediately evident since taking two general elements  $x = \alpha + \beta e_1$  and  $y = \gamma + \delta e_1$  in  $\mathcal{G}(\mathbb{R}^1)$  we have

$$xy = (\alpha + \beta e_1)(\gamma + \delta e_1) = \alpha\gamma + \beta\delta + (\alpha\delta + \beta\gamma)e_1.$$

Instead, we form the two *projectors*

$$P_{\pm} := \frac{1}{2}(1 \pm e_1)$$

and note that they satisfy the relations

$$P_{\pm}^2 = P_{\pm}, \quad \text{and} \quad P_+ P_- = 0 = P_- P_+.$$

Mapping  $P_+ \mapsto (1, 0)$  and  $P_- \mapsto (0, 1)$ , we can therefore write for a general element

$$\begin{aligned} x &= \alpha + \beta e_1 = \alpha(P_+ + P_-) + \beta(P_+ - P_-) = (\alpha + \beta)P_+ + (\alpha - \beta)P_- \\ &\mapsto (\alpha + \beta, \alpha - \beta) \in \mathbb{R} \oplus \mathbb{R}, \end{aligned}$$

which one can verify completes the claimed isomorphism.

What we have done here is just to identify the two non-trivial two-sided ideals in  $\mathcal{G}(\mathbb{R}^1)$  and found canonical projections onto those. A similar procedure can be performed whenever we have an element in the algebra which commutes with everything and which squares to 1, which we know can happen for the pseudoscalar for certain dimensions and signatures (see (2.7) and Exercise 2.6).

## 5.2 Matrix algebra classification

We have already remarked that the even subspace  $\mathcal{G}^+$  of  $\mathcal{G}$  constitutes a subalgebra. The following proposition shows that this subalgebra actually is the geometric algebra of a space of one dimension lower.

**Proposition 5.1.** *We have the algebra isomorphisms*

$$\begin{aligned}\mathcal{G}^+(\mathbb{R}^{s,t}) &\cong \mathcal{G}(\mathbb{R}^{s,t-1}), \\ \mathcal{G}^+(\mathbb{R}^{s,t}) &\cong \mathcal{G}(\mathbb{R}^{t,s-1}), \\ \mathcal{G}^+(\mathbb{C}^n) &\cong \mathcal{G}(\mathbb{C}^{n-1}),\end{aligned}$$

for all  $s, t, n$  for which the expressions make sense.

*Proof.* Take an orthonormal basis  $\{e_1, \dots, e_s, \epsilon_1, \dots, \epsilon_t\}$  of  $\mathbb{R}^{s,t}$  such that  $e_i^2 = 1$ ,  $\epsilon_i^2 = -1$ , and a corresponding basis  $\{\underline{e}_1, \dots, \underline{e}_s, \underline{\epsilon}_1, \dots, \underline{\epsilon}_{t-1}\}$  of  $\mathbb{R}^{s,t-1}$ . Define  $f: \mathbb{R}^{s,t-1} \rightarrow \mathcal{G}^+(\mathbb{R}^{s,t})$  by mapping

$$\begin{aligned}\underline{e}_i &\mapsto e_i \epsilon_t, & i = 1, \dots, s, \\ \underline{\epsilon}_i &\mapsto \epsilon_i \epsilon_t, & i = 1, \dots, t-1,\end{aligned}$$

and extending linearly. We then have

$$\begin{aligned}f(\underline{e}_i)f(\underline{e}_j) &= -f(\underline{e}_j)f(\underline{e}_i), \\ f(\underline{\epsilon}_i)f(\underline{\epsilon}_j) &= -f(\underline{\epsilon}_j)f(\underline{\epsilon}_i)\end{aligned}$$

for  $i \neq j$ , and

$$\begin{aligned}f(\underline{e}_i)f(\underline{\epsilon}_j) &= -f(\underline{\epsilon}_j)f(\underline{e}_i), \\ f(\underline{e}_i)^2 &= 1, & f(\underline{\epsilon}_i)^2 &= -1\end{aligned}$$

for all reasonable  $i, j$ . By Theorem 2.1 or 2.8 (universality) we can extend  $f$  to a homomorphism  $F: \mathcal{G}(\mathbb{R}^{s,t-1}) \rightarrow \mathcal{G}^+(\mathbb{R}^{s,t})$ . Since  $\dim \mathcal{G}(\mathbb{R}^{s,t-1}) = 2^{s+t-1} = 2^{s+t}/2 = \dim \mathcal{G}^+(\mathbb{R}^{s,t})$  and  $F$  is easily seen to be surjective, we have that  $F$  is an isomorphism.

For the second statement, we take a corresponding basis  $\{\underline{e}_1, \dots, \underline{e}_t, \underline{\epsilon}_1, \dots, \underline{\epsilon}_{s-1}\}$  of  $\mathbb{R}^{t,s-1}$  and define  $f: \mathbb{R}^{t,s-1} \rightarrow \mathcal{G}^+(\mathbb{R}^{s,t})$  by

$$\begin{aligned}\underline{e}_i &\mapsto \epsilon_i e_s, & i = 1, \dots, t, \\ \underline{\epsilon}_i &\mapsto e_i e_s, & i = 1, \dots, s-1.\end{aligned}$$

Proceeding as above, we obtain the isomorphism.

The last isomorphism is left to the reader as an exercise.  $\square$

**Corollary.** *It follows immediately that*

$$\begin{aligned}\mathcal{G}(\mathbb{R}^{s,t}) &\cong \mathcal{G}(\mathbb{R}^{t+1,s-1}), \\ \mathcal{G}^+(\mathbb{R}^{s,t}) &\cong \mathcal{G}^+(\mathbb{R}^{t,s}).\end{aligned}$$

**Example 5.1.** We have already seen explicitly that  $\mathcal{G}^+(\mathbb{R}^2) \cong \mathbb{C} \cong \mathcal{G}(\mathbb{R}^{0,1})$  and that the even subalgebra of the space algebra is  $\mathcal{G}^+(\mathbb{R}^3) \cong \mathcal{G}(\mathbb{R}^{0,2})$  i.e. the quaternion algebra (see Example 2.9). Moreover, since the pseudoscalar  $k$  in  $\mathbb{H}$  is an imaginary unit, we also see that  $\mathcal{G}^+(\mathbb{R}^{0,2}) \cong \mathbb{C} \cong \mathcal{G}^+(\mathbb{R}^{2,0})$ .

The property of geometric algebras that leads to their eventual classification as matrix algebras is that they can be split up into tensor products of geometric algebras of lower dimension.

**Proposition 5.2.** *We have the algebra isomorphisms*

$$\begin{aligned}\mathcal{G}(\mathbb{R}^{n+2,0}) &\cong \mathcal{G}(\mathbb{R}^{0,n}) \otimes \mathcal{G}(\mathbb{R}^{2,0}), \\ \mathcal{G}(\mathbb{R}^{0,n+2}) &\cong \mathcal{G}(\mathbb{R}^{n,0}) \otimes \mathcal{G}(\mathbb{R}^{0,2}), \\ \mathcal{G}(\mathbb{R}^{s+1,t+1}) &\cong \mathcal{G}(\mathbb{R}^{s,t}) \otimes \mathcal{G}(\mathbb{R}^{1,1}),\end{aligned}$$

for all  $n, s$  and  $t$  for which the expressions make sense.

*Proof.* For the first expression, take orthonormal bases  $\{e_i\}$  of  $\mathbb{R}^{n+2}$ ,  $\{\underline{e}_i\}$  of  $\mathbb{R}^{0,n}$  and  $\{\bar{e}_i\}$  of  $\mathbb{R}^2$ . Define a mapping  $f: \mathbb{R}^{n+2} \rightarrow \mathcal{G}(\mathbb{R}^{0,n}) \otimes \mathcal{G}(\mathbb{R}^2)$  by

$$\begin{aligned}e_j &\mapsto \underline{e}_j \otimes \bar{e}_1 \bar{e}_2, & j = 1, \dots, n, \\ e_j &\mapsto 1 \otimes \bar{e}_{j-n}, & j = n+1, n+2,\end{aligned}$$

and extend to an algebra homomorphism  $F$  using the universal property. Since  $F$  maps onto a set of generators for  $\mathcal{G}(\mathbb{R}^{0,n}) \otimes \mathcal{G}(\mathbb{R}^2)$  it is clearly surjective. Furthermore,  $\dim \mathcal{G}(\mathbb{R}^{n+2}) = 2^{n+2} = \dim \mathcal{G}(\mathbb{R}^{0,n}) \otimes \mathcal{G}(\mathbb{R}^2)$ , so  $F$  is an isomorphism.

The second expression is proved similarly. For the third expression, take orthonormal bases  $\{e_1, \dots, e_{s+1}, \epsilon_1, \dots, \epsilon_{t+1}\}$  of  $\mathbb{R}^{s+1,t+1}$ ,  $\{\underline{e}_1, \dots, \underline{e}_s, \underline{\epsilon}_1, \dots, \underline{\epsilon}_t\}$  of  $\mathbb{R}^{s,t}$  and  $\{\bar{e}, \bar{\epsilon}\}$  of  $\mathbb{R}^{1,1}$ , where  $e_i^2 = 1, \epsilon_i^2 = -1$  etc. Define  $f: \mathbb{R}^{s+1,t+1} \rightarrow \mathcal{G}(\mathbb{R}^{s,t}) \otimes \mathcal{G}(\mathbb{R}^{1,1})$  by

$$\begin{aligned}e_j &\mapsto \underline{e}_j \otimes \bar{e} \bar{\epsilon}, & j = 1, \dots, s, \\ \epsilon_j &\mapsto \underline{\epsilon}_j \otimes \bar{e} \bar{\epsilon}, & j = 1, \dots, t, \\ e_{s+1} &\mapsto 1 \otimes \bar{e}, \\ \epsilon_{t+1} &\mapsto 1 \otimes \bar{\epsilon}.\end{aligned}$$

Proceeding as above, we can extend  $f$  to an algebra isomorphism. □

**Corollary.** *It follows that*

$$\mathcal{G}(\mathbb{R}^{4,0}) \cong \mathcal{G}(\mathbb{R}^{0,4}) \cong \mathbb{H}^{2 \times 2}.$$

*Proof.* The proof is left to the reader as Exercise 5.3. □

We can also relate certain real geometric algebras to complex equivalents.

**Proposition 5.3.** *If  $s+t$  is odd and  $I^2 = -1$  then*

$$\mathcal{G}(\mathbb{R}^{s,t}) \cong \mathcal{G}^+(\mathbb{R}^{s,t}) \otimes \mathbb{C} \cong \mathcal{G}(\mathbb{C}^{s+t-1}).$$

*Proof.* Since  $s+t$  is odd, the pseudoscalar  $I$  commutes with all elements. This, together with the property  $I^2 = -1$ , makes it a good candidate for a scalar imaginary. Define  $F: \mathcal{G}^+(\mathbb{R}^{s,t}) \otimes \mathbb{C} \rightarrow \mathcal{G}(\mathbb{R}^{s,t})$  by linear extension of

$$\begin{aligned}E \otimes 1 &\mapsto E & \in \mathcal{G}^+, \\ E \otimes i &\mapsto EI & \in \mathcal{G}^-, \end{aligned}$$

for even basis blades  $E$ .  $F$  is easily seen to be a surjective algebra homomorphism. Using that the dimensions of these algebras are equal, we have an isomorphism.

For the second isomorphism, note that Proposition 5.1 gives  $\mathcal{G}^+(\mathbb{R}^{s,t}) \otimes \mathbb{C} \cong \mathcal{G}(\mathbb{R}^{s,t-1}) \otimes \mathbb{C}$ . Finally, the order of complexification is unimportant since all nondegenerate complex quadratic forms are equivalent.  $\square$

**Corollary.** *It follows immediately that, for the conditions of Proposition 5.3,*

$$\mathcal{G}(\mathbb{R}^{s,t}) \cong \mathcal{G}(\mathbb{R}^{p,q-1}) \otimes \mathbb{C}$$

for any  $p \geq 0$ ,  $q \geq 1$  such that  $p + q = s + t$ .

One important consequence of the tensor algebra isomorphisms in Proposition 5.2 is that geometric algebras experience a kind of periodicity over 8 real dimensions in the underlying vector space.

**Proposition 5.4.** *For all  $n \geq 0$ , there are periodicity isomorphisms*

$$\begin{aligned}\mathcal{G}(\mathbb{R}^{n+8,0}) &\cong \mathcal{G}(\mathbb{R}^{n,0}) \otimes \mathcal{G}(\mathbb{R}^{8,0}), \\ \mathcal{G}(\mathbb{R}^{0,n+8}) &\cong \mathcal{G}(\mathbb{R}^{0,n}) \otimes \mathcal{G}(\mathbb{R}^{0,8}), \\ \mathcal{G}(\mathbb{C}^{n+2}) &\cong \mathcal{G}(\mathbb{C}^n) \otimes_{\mathbb{C}} \mathcal{G}(\mathbb{C}^2).\end{aligned}$$

*Proof.* Using Proposition 5.2 repeatedly, we obtain

$$\begin{aligned}\mathcal{G}(\mathbb{R}^{n+8,0}) &\cong \mathcal{G}(\mathbb{R}^{0,n+6}) \otimes \mathcal{G}(\mathbb{R}^{2,0}) \\ &\cong \mathcal{G}(\mathbb{R}^{n,0}) \otimes \mathcal{G}(\mathbb{R}^{0,2}) \otimes \mathcal{G}(\mathbb{R}^{2,0}) \otimes \mathcal{G}(\mathbb{R}^{0,2}) \otimes \mathcal{G}(\mathbb{R}^{2,0}) \\ &\cong \mathcal{G}(\mathbb{R}^{n,0}) \otimes \mathcal{G}(\mathbb{R}^{8,0}),\end{aligned}$$

and analogously for the second statement.

For the last statement we take orthonormal bases  $\{e_i\}$  of  $\mathbb{C}^{n+2}$ ,  $\{\underline{e}_i\}$  of  $\mathbb{C}^n$  and  $\{\bar{e}_i\}$  of  $\mathbb{C}^2$ . Define a mapping  $f: \mathbb{C}^{n+2} \rightarrow \mathcal{G}(\mathbb{C}^n) \otimes_{\mathbb{C}} \mathcal{G}(\mathbb{C}^2)$  by

$$\begin{aligned}e_j &\mapsto i \underline{e}_j \otimes_{\mathbb{C}} \bar{e}_1 \bar{e}_2, & j = 1, \dots, n, \\ e_j &\mapsto 1 \otimes_{\mathbb{C}} \bar{e}_{j-n}, & j = n+1, n+2,\end{aligned}$$

and extend to an algebra isomorphism as usual.  $\square$

**Theorem 5.5** (Real classification). *We obtain the classification of real geometric algebras as matrix algebras, given by Table 5.1 together with the periodicity*

$$\mathcal{G}(\mathbb{R}^{s+8,t}) \cong \mathcal{G}(\mathbb{R}^{s,t+8}) \cong \mathcal{G}(\mathbb{R}^{s,t}) \otimes \mathbb{R}^{16 \times 16}.$$

*Proof.* Start with the following already discussed isomorphisms (see Section 5.1, Examples 2.6, 2.7 and Exercise 2.5):

$$\begin{aligned}\mathcal{G}(\mathbb{R}^{1,0}) &\cong \mathbb{R} \oplus \mathbb{R}, \\ \mathcal{G}(\mathbb{R}^{0,1}) &\cong \mathbb{C}, \\ \mathcal{G}(\mathbb{R}^{2,0}) &\cong \mathbb{R}^{2 \times 2}, \\ \mathcal{G}(\mathbb{R}^{0,2}) &\cong \mathbb{H}.\end{aligned}$$

8	$\mathbb{R}[16]$	$\mathbb{R}[16] \oplus \mathbb{R}[16]$	$\mathbb{R}[32]$	$\mathbb{C}[32]$	$\mathbb{H}[32]$	$\mathbb{H}[32] \oplus \mathbb{H}[32]$	$\mathbb{H}[64]$	$\mathbb{C}[128]$	$\mathbb{R}[256]$
7	$\mathbb{R}[8] \oplus \mathbb{R}[8]$	$\mathbb{R}[16]$	$\mathbb{C}[16]$	$\mathbb{H}[16]$	$\mathbb{H}[16] \oplus \mathbb{H}[16]$	$\mathbb{H}[32]$	$\mathbb{C}[64]$	$\mathbb{R}[128]$	$\mathbb{R}[128] \oplus \mathbb{R}[128]$
6	$\mathbb{R}[8]$	$\mathbb{C}[8]$	$\mathbb{H}[8]$	$\mathbb{H}[8] \oplus \mathbb{H}[8]$	$\mathbb{H}[16]$	$\mathbb{C}[32]$	$\mathbb{R}[64]$	$\mathbb{R}[64] \oplus \mathbb{R}[64]$	$\mathbb{R}[128]$
5	$\mathbb{C}[4]$	$\mathbb{H}[4]$	$\mathbb{H}[4] \oplus \mathbb{H}[4]$	$\mathbb{H}[8]$	$\mathbb{C}[16]$	$\mathbb{R}[32]$	$\mathbb{R}[32] \oplus \mathbb{R}[32]$	$\mathbb{R}[64]$	$\mathbb{C}[64]$
4	$\mathbb{H}[2]$	$\mathbb{H}[2] \oplus \mathbb{H}[2]$	$\mathbb{H}[4]$	$\mathbb{C}[8]$	$\mathbb{R}[16]$	$\mathbb{R}[16] \oplus \mathbb{R}[16]$	$\mathbb{R}[32]$	$\mathbb{C}[32]$	$\mathbb{H}[32]$
3	$\mathbb{H} \oplus \mathbb{H}$	$\mathbb{H}[2]$	$\mathbb{C}[4]$	$\mathbb{R}[8]$	$\mathbb{R}[8] \oplus \mathbb{R}[8]$	$\mathbb{R}[16]$	$\mathbb{C}[16]$	$\mathbb{H}[16]$	$\mathbb{H}[16] \oplus \mathbb{H}[16]$
2	$\mathbb{H}$	$\mathbb{C}[2]$	$\mathbb{R}[4]$	$\mathbb{R}[4] \oplus \mathbb{R}[4]$	$\mathbb{R}[8]$	$\mathbb{C}[8]$	$\mathbb{H}[8]$	$\mathbb{H}[8] \oplus \mathbb{H}[8]$	$\mathbb{H}[16]$
1	$\mathbb{C}$	$\mathbb{R}[2]$	$\mathbb{R}[2] \oplus \mathbb{R}[2]$	$\mathbb{R}[4]$	$\mathbb{C}[4]$	$\mathbb{H}[4]$	$\mathbb{H}[4] \oplus \mathbb{H}[4]$	$\mathbb{H}[8]$	$\mathbb{C}[16]$
0	$\mathbb{R}$	$\mathbb{R} \oplus \mathbb{R}$	$\mathbb{R}[2]$	$\mathbb{C}[2]$	$\mathbb{H}[2]$	$\mathbb{H}[2] \oplus \mathbb{H}[2]$	$\mathbb{H}[4]$	$\mathbb{C}[8]$	$\mathbb{R}[16]$
	0	1	2	3	4	5	6	7	8

Table 5.1: The algebra  $\mathcal{G}(\mathbb{R}^{s,t})$  in the box  $(s,t)$ , where  $\mathbb{F}[N] = \mathbb{F}^{N \times N}$ .

We can now work out the cases  $(n, 0)$  and  $(0, n)$  for  $n = 0, 1, \dots, 7$  in a criss-cross fashion using Proposition 5.2 and the tensor isomorphisms (see Exercises 5.6-5.8)

$$\begin{aligned}\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} &\cong \mathbb{C} \oplus_{\mathbb{R}} \mathbb{C}, \\ \mathbb{C} \otimes_{\mathbb{R}} \mathbb{H} &\cong \mathbb{C}^{2 \times 2}, \\ \mathbb{H} \otimes_{\mathbb{R}} \mathbb{H} &\cong \mathbb{R}^{4 \times 4},\end{aligned}$$

and (see A.1.7)

$$\begin{aligned}\mathbb{R}^{n \times n} \otimes_{\mathbb{R}} \mathbb{R}^{m \times m} &\cong \mathbb{R}^{nm \times nm}, \\ \mathbb{R}^{n \times n} \otimes_{\mathbb{R}} \mathbb{K} &\cong \mathbb{K}^{n \times n},\end{aligned}$$

for all  $n, m$  and  $\mathbb{K} = \mathbb{C}$  or  $\mathbb{H}$ . With  $\mathcal{G}(\mathbb{R}^{1,1}) \cong \mathcal{G}(\mathbb{R}^{2,0})$  and Proposition 5.2 we can then work our way through the whole table diagonally. The periodicity follows from Proposition 5.4 and

$$\mathcal{G}(\mathbb{R}^{8,0}) \cong \mathbb{H} \otimes \mathbb{R}^{2 \times 2} \otimes \mathbb{H} \otimes \mathbb{R}^{2 \times 2} \cong \mathbb{R}^{16 \times 16}. \quad \square$$

Because all nondegenerate complex quadratic forms on  $\mathbb{C}^n$  are equivalent, the complex version of the above theorem turns out to be much simpler.

**Theorem 5.6** (Complex classification). *We obtain the classification of complex geometric algebras as matrix algebras, given by*

$$\begin{aligned}\mathcal{G}(\mathbb{C}^0) &\cong \mathbb{C}, \\ \mathcal{G}(\mathbb{C}^1) &\cong \mathbb{C} \oplus \mathbb{C},\end{aligned}$$

together with the periodicity

$$\mathcal{G}(\mathbb{C}^{n+2}) \cong \mathcal{G}(\mathbb{C}^n) \otimes_{\mathbb{C}} \mathbb{C}^{2 \times 2}.$$

In other words,

$$\begin{aligned}\mathcal{G}(\mathbb{C}^{2k}) &\cong \mathbb{C}^{2^k \times 2^k}, \\ \mathcal{G}(\mathbb{C}^{2k+1}) &\cong \mathbb{C}^{2^k \times 2^k} \oplus \mathbb{C}^{2^k \times 2^k},\end{aligned}$$

for  $k = 0, 1, 2, \dots$

*Proof.* The isomorphism  $\mathcal{G}(\mathbb{C}^n) \cong \mathcal{G}(\mathbb{R}^n) \otimes \mathbb{C}$  gives

$$\begin{aligned}\mathcal{G}(\mathbb{C}^0) &\cong \mathbb{C} \\ \mathcal{G}(\mathbb{C}^1) &\cong (\mathbb{R} \oplus \mathbb{R}) \otimes \mathbb{C} \cong \mathbb{C} \oplus \mathbb{C} \\ \mathcal{G}(\mathbb{C}^2) &\cong \mathbb{R}^{2 \times 2} \otimes \mathbb{C} \cong \mathbb{C}^{2 \times 2}.\end{aligned}$$

Then use Proposition 5.4 for periodicity. □

**Example 5.2.** A concrete representation of the space algebra  $\mathcal{G}(\mathbb{R}^3)$  as a matrix algebra is obtained by considering the so-called *Pauli matrices*

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (5.1)$$

These satisfy  $\sigma_1^2 = \sigma_2^2 = \sigma_3^2 = 1$ ,  $\sigma_j \sigma_k = -\sigma_k \sigma_j$ ,  $j \neq k$ , and  $\sigma_1 \sigma_2 \sigma_3 = i$ , and hence identifies the space algebra with the *Pauli algebra*  $\mathbb{C}^{2 \times 2}$  through the



isomorphism  $\rho : \mathcal{G}(\mathbb{R}^3) \rightarrow \mathbb{C}^{2 \times 2}$ , defined by  $\rho(e_k) := \sigma_k$ . An arbitrary element  $x = \alpha + \mathbf{a} + \mathbf{b}I + \beta I \in \mathcal{G}$  is then represented as the matrix

$$\rho(x) = \begin{bmatrix} \alpha + a_3 + (\beta + b_3)i & a_1 + b_2 + (b_1 - a_2)i \\ a_1 - b_2 + (b_1 + a_2)i & \alpha - a_3 + (\beta - b_3)i \end{bmatrix}. \quad (5.2)$$

The periodicity of geometric algebras actually has a number of far-reaching consequences. One example is *Bott periodicity*, which, simply put, gives a periodicity in the homotopy groups  $\pi_k$  of the unitary, orthogonal and symplectic groups. An example is the following

**Theorem 5.7** (Bott periodicity). *For  $n \geq k + 2$  and all  $k \geq 1$  we have*

$$\pi_k(\mathrm{O}(n)) \cong \pi_k(\mathrm{SO}(n)) \cong \begin{cases} \{0\} & \text{if } k \equiv 2, 4, 5, 6 \pmod{8} \\ \mathbb{Z}_2 & \text{if } k \equiv 0, 1 \pmod{8} \\ \mathbb{Z} & \text{if } k \equiv 3, 7 \pmod{8} \end{cases}$$

See [27] for a proof using K-theory, or [34] and references therein for more examples.

**Exercise 5.1.** Complete the proof of Proposition 5.1.

**Exercise 5.2.** Prove the second expression of Proposition 5.2.

**Exercise 5.3.** Prove the corollary to Proposition 5.2.

**Exercise 5.4.** Complete the proof of the second isomorphism in Proposition 5.3.

**Exercise 5.5.** Complete the proof of Proposition 5.4.

**Exercise 5.6.** Prove that  $\mathbb{C} \oplus_{\mathbb{R}} \mathbb{C} \cong \mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ .

*Hint:* Consider the map

$$\begin{aligned} (1, 0) &\mapsto \frac{1}{2}(1 \otimes 1 + i \otimes i), \\ (0, 1) &\mapsto \frac{1}{2}(1 \otimes 1 - i \otimes i). \end{aligned}$$

**Exercise 5.7.** Prove that  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{H} \cong \mathbb{C}^{2 \times 2}$ .

*Hint:* Consider  $\mathbb{H}$  as a  $\mathbb{C}$ -module under left scalar multiplication, and define an  $\mathbb{R}$ -bilinear map  $\Phi : \mathbb{C} \times \mathbb{H} \rightarrow \mathrm{Hom}_{\mathbb{C}}(\mathbb{H}, \mathbb{H})$  by setting  $\Phi_{z,q}(x) := zx\bar{q}$ . This extends (by the universal property of the tensor product; see Theorem A.1) to an  $\mathbb{R}$ -linear map  $\Phi : \mathbb{C} \otimes_{\mathbb{R}} \mathbb{H} \rightarrow \mathrm{Hom}_{\mathbb{C}}(\mathbb{H}, \mathbb{H}) \cong \mathbb{C}^{2 \times 2}$ .

**Exercise 5.8.** Prove that  $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{H} \cong \mathbb{R}^{4 \times 4}$ .

*Hint:* Consider the  $\mathbb{R}$ -bilinear map  $\Psi : \mathbb{H} \times \mathbb{H} \rightarrow \mathrm{Hom}_{\mathbb{R}}(\mathbb{H}, \mathbb{H})$  given by setting  $\Psi_{q_1, q_2}(x) := q_1 x \bar{q}_2$ .

### 5.3 Graded tensor products and the mother algebra

Let us also consider a different characterization of geometric algebras.

Let  $\mathcal{G}(V_1, q_1)$  and  $\mathcal{G}(V_2, q_2)$  be two geometric algebras, and form their tensor product (considered not as an algebra, but as an  $\mathbb{F}$ -module)

$$T = \mathcal{G}(V_1, q_1) \otimes \mathcal{G}(V_2, q_2).$$

We shall now introduce a new multiplication on  $T$  through

$$\begin{aligned} T \times T &\rightarrow T \\ (x \otimes y, x' \otimes y') &\mapsto (-1)^{\delta(y)\delta(x')} (xx') \otimes (yy') \end{aligned}$$

for elements such that

$$\begin{aligned} x, x' &\in \mathcal{G}^+(V_1, q_1) \cup \mathcal{G}^-(V_1, q_1), \\ y, y' &\in \mathcal{G}^+(V_2, q_2) \cup \mathcal{G}^-(V_2, q_2), \end{aligned}$$

and where  $\delta(z) = 0$  if  $z \in \mathcal{G}^+$  and  $\delta(z) = 1$  if  $z \in \mathcal{G}^-$ . This multiplication map is then extended to the other elements through bilinearity.

We immediately note that

$$\begin{aligned} (v_1 \otimes 1 + 1 \otimes v_2)^2 &= (v_1 \otimes 1)(v_1 \otimes 1) + (v_1 \otimes 1)(1 \otimes v_2) \\ &\quad + (1 \otimes v_2)(v_1 \otimes 1) + (1 \otimes v_2)(1 \otimes v_2) \\ &= v_1^2 \otimes 1 + v_1 \otimes v_2 - v_1 \otimes v_2 + 1 \otimes v_2^2 \\ &= (v_1^2 + v_2^2)1 \otimes 1, \end{aligned}$$

so that, if we as usual identify  $1 \otimes 1$  with  $1 \in \mathbb{F}$ , we obtain

$$(v_1 \otimes 1 + 1 \otimes v_2)^2 = v_1^2 + v_2^2 = q(v_1) + q(v_2).$$

Hence, if we introduce the vector space

$$V = \{v_1 \otimes 1 + 1 \otimes v_2 : v_1 \in V_1, v_2 \in V_2\} \cong V_1 \oplus V_2,$$

and the quadratic form  $q : V \rightarrow \mathbb{F}$ ,

$$q(v_1 \otimes 1 + 1 \otimes v_2) := q_1(v_1) + q_2(v_2),$$

then we find that  $\mathcal{G}(V, q)$  becomes a Clifford algebra which is isomorphic to  $T = \mathcal{G}(V_1, q_1) \hat{\otimes} \mathcal{G}(V_2, q_2)$ , where the symbol  $\hat{\otimes}$  signals that we have defined a special so-called  $\mathbb{Z}_2$ -graded multiplication on the usual tensor product space.

As a consequence, we have the following

**Proposition 5.8.** *For all  $s, t, p, q \geq 0$ , there is a graded tensor algebra isomorphism*

$$\mathcal{G}(\mathbb{R}^{s+p, t+q}) \cong \mathcal{G}(\mathbb{R}^{s, t}) \hat{\otimes} \mathcal{G}(\mathbb{R}^{p, q}).$$

**Corollary.** *It follows immediately that*

$$\mathcal{G}(\mathbb{R}^{s, t}) \cong \underbrace{\mathcal{G}(\mathbb{R}^{1,0}) \hat{\otimes} \dots \hat{\otimes} \mathcal{G}(\mathbb{R}^{1,0})}_{s \text{ factors}} \hat{\otimes} \underbrace{\mathcal{G}(\mathbb{R}^{0,1}) \hat{\otimes} \dots \hat{\otimes} \mathcal{G}(\mathbb{R}^{0,1})}_{t \text{ factors}}.$$

When working with a geometric algebra over a mixed-signature space, where the quadratic form is neither positive nor negative definite, it can often be a good idea to embed the algebra in a larger one.

Consider  $\mathcal{G}(\mathbb{R}^{s,t,u})$  generated by an orthonormal basis

$$\{e_1^+, \dots, e_s^+, e_1^-, \dots, e_t^-, e_1^0, \dots, e_u^0\}$$

with  $(e_i^+)^2 = 1$ ,  $(e_j^-)^2 = -1$ , and  $(e_k^0)^2 = 0$ . Introduce the *mother algebra*  $\mathcal{G}(\mathbb{R}^{n,n})$ , with  $n = s + t + u$ , and an orthonormal basis

$$\{f_1^+, \dots, f_n^+, f_1^-, \dots, f_n^-\}$$

with  $(f_i^+)^2 = 1$  and  $(f_j^-)^2 = -1$ .

We now define  $\Phi : \mathcal{G}(\mathbb{R}^{s,t,u}) \rightarrow \mathcal{G}(\mathbb{R}^{n,n})$  on  $\mathbb{R}^{s,t,u}$  by setting

$$\Phi(e_i^+) := f_i^+, \quad \Phi(e_j^-) := f_j^-, \quad \Phi(e_k^0) := f_{s+t+k}^+ - f_{s+t+k}^-,$$

and extending linearly. Note that

$$\Phi(e_i^+)^2 = (e_i^+)^2, \quad \Phi(e_j^-)^2 = (e_j^-)^2, \quad \Phi(e_k^0)^2 = (e_k^0)^2.$$

By universality,  $\Phi$  extends to all of  $\mathcal{G}(\mathbb{R}^{s,t,u})$  and since  $\Phi$  is injective  $\mathbb{R}^{s,t,u} \rightarrow \mathbb{R}^{n,n}$ , it follows that  $\Phi$  is an injective homomorphism of geometric algebras.

Similarly, it also follows that every finite-dimensional geometric algebra  $\mathcal{G}(\mathbb{R}^{s,t,u})$  is embedded in the *infinite-dimensional mother algebra*  $\mathcal{Cl}_{(\mathcal{F})}(\mathbb{Z}, \mathbb{R}, r)$ , where  $r(k) := (k \geq 0) - (k < 0)$ ,  $k \in \mathbb{Z}$ .

---

**Exercise 5.9.** Verify that the graded multiplication on  $T$  introduced above is associative.

## 6 Groups

One of the reasons that geometric algebras appear naturally in many areas of mathematics and physics is the fact that they contain a number of important groups. These are groups under the geometric product and thus lie embedded within the multiplicative group of invertible elements in  $\mathcal{G}$ . In this section we will discuss the properties of various embedded groups and their relation to other familiar transformation groups such as the orthogonal and Lorentz groups. The introduced notion of a rotor will be seen to be essential for the description of the orthogonal groups, and will later play a fundamental role for understanding the concept of spinors.

Although we start out in full generality, we shall for the most part of this section assume that our scalars are real numbers unless otherwise stated. This is reasonable both from a geometric viewpoint and from the fact that e.g. many common complex groups can be represented by groups embedded in real geometric algebras. Furthermore, we assume that  $\mathcal{G}$  is nondegenerate so that we are working with a vector space of type  $\mathbb{R}^{s,t}$ . The corresponding groups associated to this space will be denoted  $SO(s,t)$  etc.

### 6.1 Groups in $\mathcal{G}$ and their actions on $\mathcal{G}$

An obvious group contained in  $\mathcal{G}$  is of course the group of all invertible elements of  $\mathcal{G}$ , which we denote by  $\mathcal{G}^\times$ . All other groups we will discuss in this section are actually subgroups of  $\mathcal{G}^\times$ , or directly related to such subgroups. Let us introduce the subgroups we will be discussing through the following definition:

**Definition 6.1.** We identify the following groups embedded in  $\mathcal{G}$ :

$\mathcal{G}^\times$	$:= \{x \in \mathcal{G} : \exists y \in \mathcal{G} : xy = yx = 1\}$	<i>the group of all invertible elements</i>
$\tilde{\Gamma}$	$:= \{x \in \mathcal{G}^\times : x^*Vx^{-1} \subseteq V\}$	<i>the Lipschitz group</i>
$\Gamma$	$:= \{v_1v_2 \dots v_k \in \mathcal{G} : v_i \in V^\times\}$	<i>the versor group</i>
$\text{Pin}$	$:= \{x \in \Gamma : xx^\dagger \in \{-1, 1\}\}$	<i>the group of unit versors</i>
$\text{Spin}$	$:= \text{Pin} \cap \mathcal{G}^+$	<i>the group of even unit versors</i>
$\text{Spin}^+$	$:= \{x \in \text{Spin} : xx^\dagger = 1\}$	<i>the rotor group</i>

In the above,  $V^\times := \{v \in V : v^2 \neq 0\}$  denotes the set of all invertible vectors.

One of the central, and highly non-trivial, results of this section is that the versor group  $\Gamma$  and Lipschitz group  $\tilde{\Gamma}$  actually are equal. Therefore,  $\Gamma$  is also called the Lipschitz group in honor of its creator. Sometimes it is also given the name *Clifford group*, but we will, in accordance with other conventions, use that name to denote the finite group generated by a fixed orthonormal basis under Clifford multiplication.

The versor (Lipschitz) group  $\Gamma$  is the smallest group which contains  $V^\times$ . Its elements are finite products of invertible vectors and are called *versors*. As seen from Definition 6.1, the  $\text{Pin}$ ,  $\text{Spin}$ , and rotor groups are all subgroups of this group. These subgroups are generated by unit vectors, and in the case of  $\text{Spin}$  and  $\text{Spin}^+$ , only an *even* number of such vector factors can be present. The elements of  $\text{Spin}^+$  are called *rotors* and, as we will see, these groups are intimately connected to orthogonal groups and rotations.

**Example 6.1.** Consider a rotation by an angle  $\varphi$  in a plane. The relevant algebra is the plane algebra  $\mathcal{G}(\mathbb{R}^2)$ , and we have seen in the introduction that the rotation can be represented as the map  $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,

$$\mathbf{v} \mapsto \mathbf{v}e^{\varphi I} = e^{-\frac{\varphi}{2}I}\mathbf{v}e^{\frac{\varphi}{2}I}, \quad I = e_1e_2. \quad (6.1)$$

The invertible element  $R_\varphi := e^{\frac{\varphi}{2}I} = \cos \frac{\varphi}{2} + \sin \frac{\varphi}{2}I = (R_{-\varphi})^{-1}$  is clearly in the even subalgebra, so  $R_\varphi^* = R_\varphi$ , and we also see from (6.1) that  $R_\varphi\mathbb{R}^2R_\varphi^{-1} \subseteq \mathbb{R}^2$ , hence by definition  $R_\varphi \in \tilde{\Gamma}$ . The set of elements on this form, i.e. the even unit multivectors  $\{e^{\varphi I} : \varphi \in \mathbb{R}\}$ , is obviously identical to the group  $U(1)$  of unit complex numbers. Furthermore, we can always write e.g.

$$R_\varphi = e_1 \left( \cos \frac{\varphi}{2} e_1 + \sin \frac{\varphi}{2} e_2 \right),$$

i.e.  $R_\varphi$  as a product of two unit vectors (a unit versor), and conversely any product of two unit vectors can be put on this form by a choice of basis. Hence we have found the group of rotors in two dimensions:  $\text{Spin}(2) = \text{Spin}^+(2) \cong U(1)$ . However, note that because of the factor  $\frac{1}{2}$  multiplying the angle  $\varphi$  in  $R_\varphi$ , the map from the rotors to the actual rotations (6.1) is a *two-to-one* map  $U(1) \rightarrow \text{SO}(2) \cong U(1)$ . In a sense (which will be made more precise in this section), a rotor can be thought of as representing (plus/minus) the square root of a rotation.

In order to understand how groups embedded in a geometric algebra are related to more familiar groups of linear transformations, it is necessary to study how groups in  $\mathcal{G}$  can act on the vector space  $\mathcal{G}$  itself and on the embedded underlying vector space  $V$ . The following are natural candidates for such actions.

**Definition 6.2.** Derived from the geometric product  $(x, y) \mapsto xy$ , we have the following canonical actions:<sup>15</sup>

$$\begin{aligned} L: \mathcal{G} &\rightarrow \text{End } \mathcal{G} && \textit{left action} \\ x &\mapsto L_x: y \mapsto xy \\ R: \mathcal{G} &\rightarrow \text{End } \mathcal{G} && \textit{right action} \\ x &\mapsto R_x: y \mapsto yx \\ \text{Ad}: \mathcal{G}^\times &\rightarrow \text{Aut } \mathcal{G} \subseteq \text{End } \mathcal{G} && \textit{adjoint action} \\ x &\mapsto \text{Ad}_x: y \mapsto xyx^{-1} \\ \widetilde{\text{Ad}}: \mathcal{G}^\times &\rightarrow \text{Aut } \mathcal{G} \subseteq \text{End } \mathcal{G} && \textit{twisted adjoint action} \\ x &\mapsto \widetilde{\text{Ad}}_x: y \mapsto x^*yx^{-1} \end{aligned}$$

where  $\text{End } \mathcal{G}$  resp.  $\text{Aut } \mathcal{G}$  are the (vector space) endomorphisms resp. automorphisms of  $\mathcal{G}$ .

Note that the left and right actions  $L$  and  $R$  are algebra homomorphisms while  $\text{Ad}$  and  $\widetilde{\text{Ad}}$  are group homomorphisms (Exercise 6.2). These actions give rise to canonical representations of the groups embedded in  $\mathcal{G}$ . Although, using e.g. the expansion (3.1), one can verify that  $\text{Ad}_x$  is always an outermorphism while

<sup>15</sup>Note the convention on the adjoint actions, which will imply an orientation convention on rotors that differs from (6.1).

in general  $\widetilde{\text{Ad}}_x$  is not (Exercise 6.2), the twisted adjoint action  $\widetilde{\text{Ad}}$  takes the graded structure of  $\mathcal{G}$  properly into account and will be seen to actually play a more important role than the normal adjoint action in geometric algebra. Note, however, that these actions agree on the subgroup of even invertible elements  $\mathcal{G}^\times \cap \mathcal{G}^+$ , and therefore in particular on the Spin/rotor group.

**Example 6.2.** As an application, let us recall that, because the algebra  $\mathcal{G}$  is assumed to be finite-dimensional, left inverses in  $\mathcal{G}$  are always right inverses and *vice versa*. This can be seen as follows. First note that the left and right actions are injective. Namely, assume that  $L_x = 0$ . Then  $L_x(y) = 0 \forall y$  and in particular  $L_x(1) = x = 0$  (and similarly for  $R$ ). Suppose now that  $xy = 1$  for some  $x, y \in \mathcal{G}$ . But then  $L_x L_y = L_1 = \text{id}$ , so that  $L_y$  is a right inverse to  $L_x$ . Now, using the dimension theorem for finite-dimensional linear maps/matrices

$$\dim \ker L_y + \dim \text{im } L_y = \dim \mathcal{G}$$

with  $\ker L_y = 0$ , we can conclude that  $L_y$  is also a left inverse to  $L_x$ . Hence,  $L_y L_x = \text{id}$ , so that  $L_{yx^{-1}} = 0$ , and  $yx = 1$ .

---

**Exercise 6.1.** Verify that the groups defined in Definition 6.1 really are groups under the geometric product (i.e. satisfy the group axioms, cp. A.1.1).

**Exercise 6.2.** Verify that the actions  $L$  and  $R$  are  $\mathbb{R}$ -algebra homomorphisms, while  $\text{Ad}$  and  $\widetilde{\text{Ad}} : \mathcal{G}^\times \rightarrow \text{GL}(\mathcal{G}) = \text{Aut } \mathcal{G}$  are group homomorphisms. Also, show that  $\text{Ad}_x$  is an outermorphism for fixed  $x \in \mathcal{G}^\times$ , while

$$\left( \widetilde{\text{Ad}}_x \Big|_V \right)_\wedge (y) = \text{Ad}_x(y^*)$$

for  $x \in \mathcal{G}^\times \cap \mathcal{G}^-$  and  $y \in \mathcal{G}$  (hence,  $\widetilde{\text{Ad}}_x \Big|_{\mathcal{G}}$  is *not* an outermorphism).

## 6.2 The Cartan–Dieudonné Theorem

Let us begin with studying the properties of the twisted adjoint action. For  $v \in V^\times$  we obtain by definition

$$\widetilde{\text{Ad}}_v(v) = v^* v v^{-1} = -v, \quad (6.2)$$

and, if  $w \in V$  is orthogonal to  $v$ ,

$$\widetilde{\text{Ad}}_v(w) = v^* w v^{-1} = -v w v^{-1} = w v v^{-1} = w. \quad (6.3)$$

Hence,  $\widetilde{\text{Ad}}_v$  acts on  $V$  as a reflection along  $v$ , or in other words in the hyperplane orthogonal to  $v$ . Such a transformation is a linear isometry, i.e. an element of the group of *orthogonal transformations* of  $(V, q)$ ,

$$\text{O}(V, q) := \{f: V \rightarrow V : f \text{ linear bijection s.t. } q \circ f = q\}. \quad (6.4)$$

Obviously,

$$q(\widetilde{\text{Ad}}_v(u)) = \widetilde{\text{Ad}}_v(u)^2 = v u v^{-1} v u v^{-1} = u^2 v v^{-1} = q(u) \quad (6.5)$$

for all  $u \in V$ .

For a general versor  $x = u_1 u_2 \dots u_k \in \Gamma$  we have

$$\begin{aligned} \widetilde{\text{Ad}}_x(v) &= (u_1 \dots u_k)^* v (u_1 \dots u_k)^{-1} = u_1^* \dots u_k^* v u_k^{-1} \dots u_1^{-1} \\ &= \widetilde{\text{Ad}}_{u_1} \circ \dots \circ \widetilde{\text{Ad}}_{u_k}(v), \end{aligned} \quad (6.6)$$

i.e. the map  $\widetilde{\text{Ad}}_x$  is a product of reflections in hyperplanes. It follows that the twisted adjoint action (restricted to act only on  $V$  which is clearly invariant) gives a homomorphism from the versor group into the orthogonal group  $O(V, q)$ :

$$\begin{array}{ccc} \widetilde{\text{Ad}} : & \mathcal{G}^\times & \rightarrow & \text{Aut } V \\ & \cup & & \cup \\ & \Gamma & \rightarrow & O(V, q) \end{array}$$

In Example 6.1 above we saw that any rotation in two dimensions can be written by means of an action of a rotor  $R_\varphi$  which is a product of two unit vectors, i.e. as a product of two reflections. In general, we have the following fundamental theorem regarding the orthogonal group:

**Theorem 6.1** (Cartan–Dieudonné). *Every orthogonal transformation on a non-degenerate space  $(V, q)$  is a product of reflections in hyperplanes. The number of reflections required is at most equal to the dimension of  $V$ .*

We shall first give a simple and constructive proof, where the bound on the number of reflections is  $n = \dim V$  for definite signatures, and  $2n$  for mixed-signature spaces. We then proceed with the general case concerning the optimal number of reflections, with a proof adopted from [4] and streamlined to our presentation. We only demand that  $\text{char } \mathbb{F} \neq 2$ , so that orthogonal bases exist.

**Corollary.** *The homomorphism  $\widetilde{\text{Ad}} : \Gamma \rightarrow O(V, q)$  is surjective.*

*Proof.* We know that any  $f \in O(V, q)$  can be written  $f = \widetilde{\text{Ad}}_{v_1} \circ \dots \circ \widetilde{\text{Ad}}_{v_k}$  for some invertible vectors  $v_1, \dots, v_k$ ,  $k \leq n$ . But then  $f = \widetilde{\text{Ad}}_{v_1 \dots v_k}$ , where  $v_1 v_2 \dots v_k \in \Gamma$ .  $\square$

The idea of our proof of the Cartan–Dieudonné theorem is to find an explicit map which takes a given vector into another given vector using only one or two reflections, and then apply this map consecutively to construct a given orthogonal map explicitly in terms of reflections, by mapping a prescribed set of vectors into another prescribed set of vectors (and these can eventually be taken to be orthogonal bases).

We will in the following find it convenient to introduce the following shorthand notation for the twisted adjoint action:

$$\underline{U} := \widetilde{\text{Ad}}_U, \quad \text{i.e.} \quad \underline{U}(x) = U^* x U^{-1}.$$

**Lemma 6.2.** *If  $x, y \in V^\times$  and  $x^2 = y^2$  then either  $x + y$  or  $x - y$  is in  $V^\times$ .*

*Proof.* Assume to the contrary that

$$\begin{aligned} 0 &= (x + y)^2 = x^2 + y^2 + xy + yx & \text{and} \\ 0 &= (x - y)^2 = x^2 + y^2 - xy - yx. \end{aligned}$$

Thus  $2(x^2 + y^2) = 0$ , but  $2 \neq 0$  in  $\mathbb{F}$  then implies  $2x^2 = x^2 + y^2 = 0$ , which is a contradiction.  $\square$

Note that if the quadratic form  $q$  is either positive or negative definite then, under the same conditions, either  $x = y$ ,  $x = -y$ , or *both*  $x \pm y \in V^\times$ . In fact, the only way that the invertibility of either  $x + y$  or  $x - y$  can fail is if  $x, y \in V$  are either linearly dependent or span a *degenerate* subspace (see Exercise 6.6).

**Lemma 6.3.** *With the conditions in Lemma 6.2 it holds that*

$$\begin{aligned} x + y \in V^\times &\Rightarrow \underline{(x + y)}(x) = -y \quad \text{and} \\ x - y \in V^\times &\Rightarrow \underline{(x - y)}(x) = y. \end{aligned}$$

*Proof.* If  $x + y \in V^\times$  we have

$$\begin{aligned} (x + y)^*x(x + y)^{-1} &= -(x^2 + yx)(x + y)^{-1} = -(y^2 + yx)(x + y)^{-1} \\ &= -y(y + x)(x + y)^{-1} = -y. \end{aligned}$$

The other implication is obtained by replacing  $y$  with  $-y$ . □

Finally we also note that when  $x^2 = y^2 \neq 0$  and  $x + y \in V^\times$  then

$$\underline{y(x + y)}(x) = \underline{y}(-y) = y.$$

We can now prove a general lemma:

**Lemma 6.4.** *Let  $\{e_1, \dots, e_n\}$  and  $\{f_1, \dots, f_n\}$  be two sequences of orthogonal invertible vectors in  $V$ , such that  $e_j^2 = f_j^2$ ,  $j = 1, \dots, n \leq \dim V$ . Then there exist explicit versors  $\theta_1, \dots, \theta_n \in \Gamma$ , where each  $\theta_j$  is either a 0-versor ( $\theta_j = 1$ ), 1-versor ( $\theta_j = v \in V^\times$ ), or 2-versor ( $\theta_j = v_1v_2$ ,  $v_1, v_2 \in V^\times$ ), such that*

$$\underline{\theta_n \dots \theta_1}(e_j) = f_j, \quad j = 1, \dots, n.$$

*In the case of definite signatures, only 0- or 1-versors are necessary above.*

*Proof.* Consider the first vectors of the two sequences, satisfying  $e_1^2 = f_1^2 \neq 0$ . By the above observations, we may map  $\underline{\theta_1}(e_1) = f_1$ , by taking either  $\theta_1 = 1$ ,  $\theta_1 = e_1 - f_1$  or  $\theta_1 = f_1(e_1 + f_1)$ , according to whether  $e_1 = f_1$ ,  $e_1 - f_1 \in V^\times$  or  $e_1 + f_1 \in V^\times$ , respectively (and the last case is avoided for definite  $q$ ). Denote the images of the remaining vectors by  $e_{j,1} := \underline{\theta_1}(e_j)$ ,  $j = 2, \dots, n$ . Note that, because of the orthogonality of the map  $\underline{\theta_1}$ , we have  $e_{j,1}^2 = e_j^2$  and

$$f_1 * e_{j,1} = \underline{\theta_1}(e_1) * \underline{\theta_1}(e_j) = e_1 * e_j = 0, \quad \text{for } j = 2, \dots, n,$$

and also

$$e_{j,1} * e_{k,1} = \underline{\theta_1}(e_j) * \underline{\theta_1}(e_k) = e_j * e_k = 0, \quad j \neq k,$$

which implies that the new sequence  $\{f_1, e_{2,1}, \dots, e_{n,1}\}$  is again orthogonal, satisfying the same assumptions as the original sequence  $\{e_j\}_j$ .

In the same way then, for  $e_{2,1}^2 = e_2^2 = f_2^2$ , we may find  $\theta_2 \in \Gamma$  involving only  $e_{2,1}$  and  $f_2$ , and s.t.  $\underline{\theta_2}(e_{2,1}) = f_2$ . Moreover, by the orthogonality of  $f_1$  to both these vectors, we have  $\underline{\theta_2}(f_1) = f_1$ . We then denote  $e_{j,2} := \underline{\theta_2}(e_{j,1})$ ,  $j = 3, \dots, n$ , and observe that  $\{f_1, f_2, e_{3,2}, \dots, e_{n,2}\}$  is again a suitable orthogonal sequence.



Proceeding like this, we obtain in the  $k$ :th step a versor  $\theta_k$  involving only  $f_k$  and  $e_{k,k-1}$  such that  $\theta_k(e_{k,k-1}) = f_k$  and  $\theta_k(f_j) = f_j$  for all  $j = 1, \dots, k-1$ . The procedure can be summarized in the following diagram:

$$\begin{array}{cccccccc}
& e_1 & e_2 & e_3 & \dots & e_k & \dots & e_n \\
\theta_1 & \downarrow & \downarrow & \downarrow & & \downarrow & & \downarrow \\
& f_1 & e_{2,1} & e_{3,1} & \dots & e_{k,1} & \dots & e_{n,1} \\
\theta_2 & \downarrow & \downarrow & \downarrow & & \downarrow & & \downarrow \\
& f_1 & f_2 & e_{3,2} & \dots & e_{k,2} & \dots & e_{n,2} \\
\theta_3 & \downarrow & \downarrow & \downarrow & & \downarrow & & \downarrow \\
\vdots & \vdots & \vdots & \vdots & & \vdots & & \vdots \\
\theta_n & \downarrow & \downarrow & \downarrow & & \downarrow & & \downarrow \\
& f_1 & f_2 & f_3 & \dots & f_k & \dots & f_n
\end{array}$$

In the final row we have arrived at the desired orthogonal sequence  $\{f_j\}_j$ .  $\square$

The above lemma proves the Cartan–Dieudonné theorem with the optimal number of reflections in the case of definite signatures, and with up to  $2 \dim V$  reflections for arbitrary non-degenerate spaces  $(V, q)$  over fields with  $\text{char } \mathbb{F} \neq 2$ . Namely, given any  $f \in O(V, q)$  we may take an arbitrary orthogonal basis  $\{e_j\}_j$  of  $V$  and then form the orthogonal sequence  $\{f_j := f(e_j)\}_j$  to which the lemma may be applied, proving that  $f = \theta_n \dots \theta_1$ ,  $n = \dim V$ .

We now turn to the slightly more involved proof of the “optimal” version of the theorem which is also valid for arbitrary non-degenerate spaces over fields of characteristic  $\neq 2$ . We start with some general and simple observations.

**Proposition 6.5.** *Let  $(V, q)$  be an arbitrary space,  $\text{char } \mathbb{F} \neq 2$ , and  $W \subseteq V$  a subspace. Then*

$$(OBS1) \quad W \subseteq W^\perp \Leftrightarrow W \cap V^\times = \emptyset \Leftrightarrow \forall x \in W \quad x^2 = 0,$$

$$(OBS2) \quad \text{If } a, b \text{ and } c \text{ are linearly dependent in } V, \text{ then } abc \in V.$$

Furthermore, if  $V$  is finite-dimensional and non-degenerate,

$$(OBS3) \quad \dim W + \dim W^\perp = \dim V,$$

$$(OBS4) \quad W \subseteq W^\perp \Rightarrow \dim W \leq \frac{1}{2} \dim V.$$

*Proof.* Assume  $W \cap V^\times = \emptyset$  and take  $x, y \in W$ . Then  $0 = (x+y)^2 - x^2 - y^2 = 2x * y$ , hence  $x * y = 0$ . This shows that  $W \subseteq W^\perp$ . The other implications in (OBS1) are trivial.

To verify (OBS2), assume for instance that  $c = \alpha a + \beta b$ ,  $\alpha, \beta \in \mathbb{F}$ . Then

$$abc = ab(\alpha a + \beta b) = \alpha aba + \beta b^2 a = \alpha(2a * b a - a^2 b) + \beta b^2 a \in V.$$

The other cases are treated similarly.

To prove (OBS3), note that every linear functional  $\mu: W \rightarrow \mathbb{F}$  extends to a linear functional  $\mu: V \rightarrow \mathbb{F}$  since  $V$  is finite-dimensional (just extend a basis for  $W$  to  $V$ ), i.e.  $L(V, \mathbb{F}) \rightarrow L(W, \mathbb{F})$ . Since moreover  $V$  is non-degenerate, every  $\mu \in L(V, \mathbb{F})$  can be uniquely represented by some  $v \in V$  as

$$\mu(x) = v * x, \quad \forall x \in V.$$

Hence we have a linear surjection

$$\tau: V \longleftrightarrow L(V, \mathbb{F}) \twoheadrightarrow L(W, \mathbb{F}),$$

with  $\ker \tau = W^\perp$ . By linear algebra we know that  $\dim L(W, \mathbb{F}) = \dim W$ , and therefore from the dimension theorem we can conclude (OBS3).

The last observation (OBS4) follows directly from (OBS3).  $\square$

We also note that if  $f \in \mathcal{O}(V)$  and if  $W$  is a non-degenerate invariant subspace for  $f$ , then  $W^\perp$  is also non-degenerate and invariant under  $f$  (Exercise 6.8). Thus if  $W \cap W^\perp = 0$  and  $f(W) \subseteq W$  then

$$f|_W \in \mathcal{O}(W, q|_W), \quad f|_{W^\perp} \in \mathcal{O}(W^\perp, q|_{W^\perp}), \quad \text{and } V = W \oplus W^\perp.$$

This simple fact will be used in the proof below.

**Definition 6.3.** A *simple reflection* is a map of the form  $\widetilde{\text{Ad}}_v = \underline{v}$  for some  $v \in V^\times$ . We say that a map  $f \in \mathcal{O}(V)$  is of *type  $k$*  if it can be written as a composition of  $k$  simple reflections and  $k$  is the smallest such number.

**Theorem 6.6.** *Suppose  $f \in \mathcal{O}(V)$ , where  $\mathcal{G}(V, q, \mathbb{F})$  is non-degenerate and  $\text{char } \mathbb{F} \neq 2$ . Then  $f$  is of type at most  $d = \dim V$ .*

*Proof.* The case  $d = 1$  is trivial, since  $f$  must be the identity map  $\text{id}$  or  $-\text{id}$ , and  $-\text{id}$  is a simple reflection.

Next assume that  $d = 2$ . Then we know by the earlier version of the theorem, Lemma 6.4, that  $f$  is of type at most 4, i.e. either  $f$  is of type at most 2, or  $f = \widetilde{\text{Ad}}_{abc}$ , or  $f = \widetilde{\text{Ad}}_{abcd}$  with any three of  $a, b, c, d \in V$  linearly dependent. Hence  $f$  is of type at most 2 by (OBS2).

So from now on we will assume that  $d \geq 3$ , and we proceed by induction on  $d$ . Introduce  $g := \text{id} - f$ , and consider each of the following three possibilities.

**Lemma 6.7.** *If  $V^\times \cap \ker g \neq \emptyset$  then  $f$  is of type  $\leq d - 1$ .*

*Proof of Lemma 6.7.* In this case there exists  $x \in V^\times$  such that  $f(x) = x$ . Then  $V$  splits into non-degenerate invariant subspaces,  $V = \mathbb{F}x \oplus x^\perp$ , where  $\dim x^\perp = d - 1$ , and so  $f$  is of type at most  $d - 1$  (note that the corresponding versor in  $\mathcal{G}(x^\perp)$  does not affect  $\mathbb{F}x$ ).  $\square$

**Lemma 6.8.** *If  $V^\times \cap g(V^\times) \neq \emptyset$  then  $f$  is of type  $\leq d$ .*

*Proof of Lemma 6.8.* In this case there exists  $x \in V^\times$  such that  $(x - f(x))^2 \neq 0$ . Taking  $h = \underline{x - f(x)} \circ f \in \mathcal{O}(V)$  we find  $h(x) = x$ , so by Lemma 6.7 we conclude that  $h$  is of type at most  $d - 1$ , and thus  $f = \underline{x - f(x)} \circ h$  is of type at most  $d$ .  $\square$

**Lemma 6.9.** *If both  $V^\times \cap \ker g = \emptyset$  and  $V^\times \cap g(V^\times) = \emptyset$  then  $f$  has type at most  $d$ .*

*Proof of Lemma 6.9.* First note that  $\ker g \subseteq (\ker g)^\perp$ , so that  $\dim \ker g \leq d/2$  by (OBS4).

We claim that also  $\text{im } g \subseteq (\text{im } g)^\perp$ , that is  $V^\times \cap g(V) = \emptyset$ . By the assumptions we need only prove that  $V^\times \cap g(V \setminus V^\times) = \emptyset$ , that is  $g(y)^2 = 0$  whenever  $y^2 = 0$ ,  $y \in V$ . Take then  $y \in V$  with  $y^2 = 0$ . Since  $d \geq 3$ , we have by (OBS3)

that  $\dim y^\perp = d - 1 > d/2$ . Then, by (OBS4) there exists some  $z \in y^\perp \cap V^\times$ . But then  $z^2 \neq 0$  and

$$(y \pm z)^2 = y^2 + z^2 \pm 2y * z = z^2 \neq 0,$$

and therefore

$$g(z)^2 = g(y - z)^2 = g(y + z)^2 = 0,$$

by the assumption on  $g$ . Using the parallelogram identity we then have

$$\begin{aligned} 0 &= g(y - z)^2 + g(y + z)^2 = (g(y) - g(z))^2 + (g(y) + g(z))^2 \\ &= 2g(y)^2 + 2g(z)^2 = 2g(y)^2, \end{aligned}$$

hence also  $g(y)^2 = 0$ .

Thus we have proved  $\operatorname{im} g \subseteq (\operatorname{im} g)^\perp$ , and it follows that  $\dim \operatorname{im} g \leq d/2$ . But  $\dim \ker g + \dim \operatorname{im} g = d$ , so we obtain

$$\dim \ker g = \dim \operatorname{im} g = d/2.$$

From this we also conclude that  $d$  must be even.

Our next step is to show that  $\ker g$  and  $\operatorname{im} g$  are orthogonal. Take  $z \in \ker g$  and  $y \in \operatorname{im} g$ . Then  $f(z) = z$  and  $y = x - f(x)$  for some  $x \in V$ , and thus

$$z * y = z * (x - f(x)) = z * x - z * f(x) = z * x - f(z) * f(x) = 0.$$

Therefore  $\ker g \perp \operatorname{im} g$ , and we can conclude that

$$\ker g \subseteq (\ker g)^\perp \cap (\operatorname{im} g)^\perp \quad \text{and} \quad \operatorname{im} g \subseteq (\operatorname{im} g)^\perp \cap (\ker g)^\perp.$$

We also claim that  $\operatorname{im} g \subseteq \ker g$ , so that  $g \circ g = 0$ . But this follows again from a dimensionality argument:

$$\dim \ker g + \dim(\ker g)^\perp = d = \dim \operatorname{im} g + \dim(\operatorname{im} g)^\perp,$$

implying  $\dim(\ker g)^\perp = d/2$ , hence  $\ker g = (\ker g)^\perp$ , and similarly for  $\operatorname{im} g$ , so that what we actually have is the null-space

$$\ker g = \operatorname{im} g = (\ker g)^\perp = (\operatorname{im} g)^\perp.$$

We have shown that  $d$  is even and  $g \circ g = 0$ . Now let  $\{e_1, \dots, e_{d/2}\}$  be any basis for  $\ker g$  and extend it to a basis  $\{e_1, \dots, e_d\}$  for  $V$ . Then  $g(e_i) = 0$ , i.e.  $f(e_i) = e_i$ , for  $1 \leq i \leq d/2$ , and  $g(e_j) \in \ker g = \mathbb{F}e_1 + \dots + \mathbb{F}e_{d/2}$ , for  $d/2 < j \leq d$ . We can also write  $f(e_j) = e_j - f_j$ , where  $f_j := g(e_j) \in \ker g$ ,  $d/2 < j \leq d$ . Then  $f_j \wedge e_1 \wedge \dots \wedge e_{d/2} = 0$ , so

$$\begin{aligned} &f \wedge (e_1 \wedge \dots \wedge e_{d/2} \wedge e_{d/2+1} \wedge \dots \wedge e_d) \\ &= e_1 \wedge \dots \wedge e_{d/2} \wedge (e_{d/2+1} - f_{d/2+1}) \wedge \dots \wedge (e_d - f_d) = e_1 \wedge \dots \wedge e_d, \end{aligned}$$

and hence  $\det f = 1$ .

Finally, let  $h = \underline{v} \circ f$  for some  $v \in V^\times$ . Then  $\det h = -1$ , so  $h$  must satisfy the conditions of Lemma 6.7 or Lemma 6.8, and furthermore  $h$  has odd type. But since  $d$  is even,  $h$  must then have type at most  $d - 1$ . Then  $f = \underline{v} \circ h$  is of type  $\leq d$ .  $\square$

We have above covered all possible cases, and thus Theorem 6.6 is proved.  $\square$

A couple of general observations follow only from this optimal version of the Cartan–Dieudonné theorem:

**Corollary.** *If  $f \in O(V)$ ,  $\dim V$  is odd, and  $\det f = 1$ , then  $f$  has a fixed point:*

$$\exists x \in V \setminus \{0\} \text{ such that } f(x) = x.$$

*Proof.* The map  $f$  must have even type, so  $f = \underline{x_1} \circ \underline{x_2} \circ \dots \circ \underline{x_{2m}}$ , where  $x_j \in V^\times$ ,  $j \leq 2m < \dim V$ . Now choose  $0 \neq x \in x_1^\perp \cap \dots \cap x_{2m}^\perp$ . Then obviously  $f(x) = x$ .  $\square$

**Corollary.** *If  $f \in O(V)$ ,  $\dim V$  is odd, and  $\det f = -1$ , then*

$$\exists x \in V \setminus \{0\} \text{ such that } f(x) = -x.$$

*Proof.* Look at  $-f$  and use the previous corollary.  $\square$

**Exercise 6.3.** Show that for  $f \in O(V, q)$  (as defined in (6.4)),  $f_\wedge(x) * f_\wedge(y) = x * y \forall x, y \in \mathcal{G}$ ,  $f^* = f^{-1}$ , and  $\det f \in \{1, -1\}$ .

**Exercise 6.4.** Show that if  $f \in O(V, q)$  then also  $f_\wedge \in O(\mathcal{G}, q_\mathcal{G})$ , where  $q_\mathcal{G}$  is the quadratic form on  $\mathcal{G}$  inherited from  $(V, q)$  and defined by either  $q_\mathcal{G}(x) := x * x$ ,  $x \in \mathcal{G}$ , or  $q_\mathcal{G}(x) := \langle x, x \rangle_\mathcal{G}$ , as defined in Exercise 2.15.

**Exercise 6.5.** Show that  $\widetilde{\text{Ad}}_v$  is a reflection along  $v$ , in an alternative way, by using the formula for a projection in Section 3.3. Also, verify that  $\det \widetilde{\text{Ad}}_v = -1$ .

**Exercise 6.6.** Show that  $(x + y)^2(x - y)^2 = -4(x \wedge y)^2$  for all  $x, y \in V$  s.t.  $x^2 = y^2$ , and construct an example of two linearly independent vectors  $x, y$  in a nondegenerate vector space  $V$  such that  $x^2 = y^2 \neq 0$  but  $x - y \notin V^\times$ .

**Exercise 6.7.** Use Lemma 6.4 to find  $v_1, v_2, v_3 \in (\mathbb{R}^3)^\times$  such that

$$\widetilde{\text{Ad}}_{v_3 v_2 v_1} = f$$

where  $f \in O(3)$  is defined by  $f(e_1) = e_3$ ,  $f(e_2) = \frac{4}{5}e_1 + \frac{3}{5}e_2$  and  $f(e_3) = \frac{3}{5}e_1 - \frac{4}{5}e_2$ .

**Exercise 6.8.** Show that if  $f \in O(V)$  and if  $W \subseteq V$  is a non-degenerate invariant subspace for  $f$ , then  $W^\perp$  is also non-degenerate and invariant under  $f$ .

### 6.3 The Lipschitz group

We saw above that the twisted adjoint action  $\widetilde{\text{Ad}}$  maps the versor group onto the full group of orthogonal transformations of  $V$ . The largest group in  $\mathcal{G}$  for which  $\widetilde{\text{Ad}}$  forms a representation on  $V$ , i.e. leaves  $V$  invariant, is by Definition 6.1 the Lipschitz group  $\tilde{\Gamma}$ . Hence, we see that  $\Gamma \subseteq \tilde{\Gamma}$  (this can also be seen explicitly from (6.6)). Our objective is to prove that actually  $\Gamma = \tilde{\Gamma}$ .

We will first introduce an important function on  $\mathcal{G}$ , conventionally called the *norm function*,

$$\begin{aligned} N: \mathcal{G} &\rightarrow \mathbb{F}, \\ N(x) &:= x^\square x. \end{aligned} \tag{6.7}$$

The name could be a bit misleading since  $N$  is not even guaranteed to take values in  $\mathbb{F}$ . However, for some special cases of algebras it does act as a natural norm (squared) and we will also find that it can be extended in many lower-dimensional algebras to a function which acts as a kind of determinant. Our first main result for this function is that it has properties in common with a determinant on  $\tilde{\Gamma}$ . This will help us prove that  $\Gamma = \tilde{\Gamma}$ .

**Lemma 6.10.** *Assume that  $\mathcal{G}$  is nondegenerate. If  $x \in \mathcal{G}$  and  $x^*v = vx$  for all  $v \in V$  then  $x$  must be a scalar.*

*Proof.* From Proposition 2.10 we have that  $v \lrcorner x = 0$  for all  $v \in V$ . This means that, for a  $k$ -blade,  $(v_1 \wedge \cdots \wedge v_{k-1} \wedge v_k) * x = (v_1 \wedge \cdots \wedge v_{k-1}) * (v_k \lrcorner x) = 0$  whenever  $k \geq 1$ . The nondegeneracy of the scalar product implies that  $x$  must have grade 0.  $\square$

**Theorem 6.11.** *The norm function is a group homomorphism  $N: \tilde{\Gamma} \rightarrow \mathbb{F}^\times$ .*

*Proof.* First note that if  $xx^{-1} = 1$  then also  $x^*(x^{-1})^* = 1$  and  $(x^{-1})^\dagger x^\dagger = 1$ , hence  $(x^*)^{-1} = (x^{-1})^*$  and  $(x^\dagger)^{-1} = (x^{-1})^\dagger$ .

Now take  $x \in \tilde{\Gamma}$ . Then  $x^*vx^{-1} \in V$  for all  $v \in V$  and therefore

$$x^*vx^{-1} = (x^*vx^{-1})^\dagger = (x^{-1})^\dagger vx^\square. \tag{6.8}$$

This means that  $x^\dagger x^*v = vx^\square x$ , or  $N(x)^*v = vN(x)$ . By Lemma 6.10 we find that  $N(x) \in \mathbb{F}$ . The homomorphism property now follows easily, since for  $x, y \in \tilde{\Gamma}$ ,

$$N(xy) = (xy)^\square xy = y^\square x^\square xy = y^\square N(x)y = N(x)N(y). \tag{6.9}$$

Finally, because  $1 = N(1) = N(xx^{-1}) = N(x)N(x^{-1})$ , we must have that  $N(x)$  is nonzero.  $\square$

**Lemma 6.12.** *We have a homomorphism  $\widetilde{\text{Ad}}: \tilde{\Gamma} \rightarrow \text{O}(V, q)$  with kernel  $\mathbb{F}^\times$ .*

*Proof.* We first prove that  $\widetilde{\text{Ad}}_x$  is an orthogonal map for all  $x \in \tilde{\Gamma}$ . Note that, for  $v \in V$ ,

$$\begin{aligned} N(\widetilde{\text{Ad}}_x(v)) &= N(x^*vx^{-1}) = (x^*vx^{-1})^\square x^*vx^{-1} \\ &= (x^{-1})^\square v^\square x^\dagger x^*vx^{-1} = (x^{-1})^\square v^\square N(x)^*vx^{-1} \\ &= (x^{-1})^\square v^\square vx^{-1}N(x)^* = N(v)N(x^{-1})N(x) = N(v). \end{aligned} \tag{6.10}$$

Then, since  $N(v) = v^\square v = -v^2$ , we have that  $\widetilde{\text{Ad}}_x(v)^2 = v^2$ .

Now, if  $\widetilde{\text{Ad}}_x = \text{id}$  then  $x^*v = vx$  for all  $v \in V$  and by Lemma 6.10 we must have  $x \in \mathbb{F} \cap \tilde{\Gamma} = \mathbb{F}^\times$ .  $\square$

We finally obtain

**Theorem 6.13.** *The Lipschitz group and the versor group are equal,  $\tilde{\Gamma} = \Gamma$ .*

*Proof.* We saw earlier that  $\Gamma \subseteq \tilde{\Gamma}$ . Take  $x \in \tilde{\Gamma}$ . By the above lemma we have  $\tilde{\text{Ad}}_x \in \text{O}(V, q)$ . Using the corollary to Theorem 6.1 we then find that  $\tilde{\text{Ad}}_x = \tilde{\text{Ad}}_y$  for some  $y \in \Gamma$ . Then  $\tilde{\text{Ad}}_{xy^{-1}} = \text{id}$ , and  $xy^{-1} = \lambda \in \mathbb{F}^\times$ . Hence,  $x = \lambda y \in \Gamma$ .  $\square$

Combining this theorem with the Cartan–Dieudonné theorem also leads to the following simplification of the length of any versor.

**Corollary.** *In any non-degenerate Clifford algebra  $\mathcal{G}(V, q, \mathbb{F})$ , with  $\dim V = n$  and  $\text{char } \mathbb{F} \neq 2$ , any versor  $x \in \Gamma$  can be written using at most  $n$  vector factors, i.e.  $x = v_1 v_2 \dots v_k$ , where  $v_j \in V^\times$ ,  $1 \leq j \leq k \leq n$ , or  $x \in \mathbb{F}^\times$ .*

*Proof.* Given  $x \in \Gamma$  we have  $\tilde{\text{Ad}}_x = \tilde{\text{Ad}}_{v_1 v_2 \dots v_k}$  for at most  $k \leq n$  vectors  $v_j \in V^\times$  by the optimal formulation of the Cartan–Dieudonné theorem 6.1, and hence  $x = (\lambda v_1) v_2 \dots v_k$ ,  $\lambda \in \mathbb{F}^\times$ , by Lemma 6.12 and Theorem 6.13.  $\square$

---

**Exercise 6.9.** Let  $\mathcal{G}^{\times\pm} := \mathcal{G}^\times \cap \mathcal{G}^\pm$  and show that the Lipschitz group  $\Gamma$  also can be defined through

$$\Gamma = \{x \in \mathcal{G}^{\times+} \cup \mathcal{G}^{\times-} : \text{Ad}_x(V) \subseteq V\}.$$

## 6.4 Properties of the Pin and Spin groups

We shall from now on restrict our attention to spaces over  $\mathbb{R}$ . From the previous subsections followed that  $\tilde{\text{Ad}}$  gives a surjective homomorphism from the versor/Lipschitz group  $\Gamma$  to the orthogonal group. The kernel of this homomorphism is the set of invertible scalars. Because the Pin and Spin groups consist of *normalized* versors (i.e.  $N(x) = \pm 1$ ) we then obtain the following

**Theorem 6.14.** *The homomorphisms*

$$\begin{aligned} \tilde{\text{Ad}}: \text{Pin}(s, t) &\rightarrow \text{O}(s, t) \\ \tilde{\text{Ad}}: \text{Spin}(s, t) &\rightarrow \text{SO}(s, t) \\ \tilde{\text{Ad}}: \text{Spin}^+(s, t) &\rightarrow \text{SO}^+(s, t) \end{aligned}$$

*are continuous and surjective with kernel  $\{\pm 1\}$ .*

The homomorphism onto the *special orthogonal group*,

$$\text{SO}(V, q) := \{f \in \text{O}(V, q) : \det f = 1\}, \quad (6.11)$$

follows since it is generated by an *even* number of reflections (by the parity of the determinant; cf. Example 3.5). Furthermore, we have denoted the image of the rotor group under  $\tilde{\text{Ad}}$  by  $\text{SO}^+$  and, as will be explained shortly, this subgroup is actually the connected component of  $\text{SO}$  containing the identity.

Note that the continuity of the map  $\tilde{\text{Ad}}$  follows from the continuous (actually smooth) operation of taking products and inverses in  $\mathcal{G}^\times$ , and that both  $x$  and  $-x$  in (S)Pin are mapped to the same element in (S)O. In other words, Theorem 6.14 shows that the Pin and Spin groups are two-sheeted coverings of

the orthogonal groups. Furthermore, we have the following relations between these groups.

Take a unit versor  $\psi = u_1 u_2 \dots u_k \in \text{Pin}(s, t)$ . Note that we can always interchange its parity, namely if  $\psi$  is odd then we can multiply by a unit vector  $e$  so that  $\psi = \psi' e$  with  $\psi' = \pm \psi e \in \text{Spin}(s, t)$ . Furthermore, when the signature is euclidean we have  $\psi \psi^\dagger = u_1^2 u_2^2 \dots u_k^2 = 1$  for all unit versors. The same holds for *even* unit versors in anti-euclidean spaces since the signs cancel out. Hence,  $\text{Spin} = \text{Spin}^+$  unless there is mixed signature. But in that case we can find two orthogonal unit versors  $e_+, e_-$  such that  $e_+^2 = 1$  and  $e_-^2 = -1$ . Since  $e_+ e_- (e_+ e_-)^\dagger = -1$  we can then write  $\psi = \psi'' e_+ e_-$ , where  $\psi'' \psi''^\dagger = \psi e_+ e_- (\psi e_+ e_-)^\dagger = 1$  if  $\psi \psi^\dagger = -1$ .

Summing up, we have that, for mixed signature  $s, t \geq 1$  and any pair of orthogonal vectors  $e_+, e_-$  such that  $e_+^2 = 1, e_-^2 = -1$ ,

$$\begin{aligned}\text{Pin}(s, t) &= \text{Spin}^+(s, t) \cdot \{1, e_+, e_-, e_+ e_-\}, \\ \text{Spin}(s, t) &= \text{Spin}^+(s, t) \cdot \{1, e_+ e_-\},\end{aligned}$$

while for euclidean and anti-euclidean signatures,

$$\text{Spin} = \text{Spin}^+ \quad \text{and} \quad \text{Pin}(s, t) = \text{Spin}^{(+)}(s, t) \cdot \{1, e\},$$

for any  $e \in V$  such that  $e^2 = \pm 1$ . From the isomorphism  $\mathcal{G}^+(\mathbb{R}^{s, t}) \cong \mathcal{G}^+(\mathbb{R}^{t, s})$  we also have the signature symmetry

$$\text{Spin}(s, t) \cong \text{Spin}(t, s), \quad \text{Spin}^+(s, t) \cong \text{Spin}^+(t, s).$$

And in all cases,

$$\Gamma(s, t) = \mathbb{R}^\times \cdot \text{Pin}(s, t).$$

From these considerations it turns out to be sufficient to study the properties of the rotor groups in order to understand the Pin, Spin and orthogonal groups. Fortunately, it turns out that the rotor groups have very convenient topological features.

**Theorem 6.15.** *The groups  $\text{Spin}^+(s, t)$  are pathwise connected to either 1 or -1 for arbitrary signatures, and to 1 (hence connected) for  $s \geq 2$  or  $t \geq 2$ .*

*Proof.* Pick a rotor  $R \in \text{Spin}^+(s, t)$ , where  $s$  or  $t$  is greater than one. Then  $R = v_1 v_2 \dots v_{2k}$  with an even number of  $v_i \in V$  such that  $v_i^2 = 1$  and an even number such that  $v_i^2 = -1$ . Note that for any two invertible vectors  $a, b$  we have  $ab = aba^{-1}a = b'a$ , where  $b' = -a(b)$  is a vector with  $b'^2 = b^2$ . Hence, we can rearrange the vectors so that those with positive square come first, i.e.

$$R = a_1 b_1 \dots a_p b_p a'_1 b'_1 \dots a'_q b'_q = R_1 \dots R_p R'_1 \dots R'_q, \quad (6.12)$$

where  $a_i^2 = b_i^2 = 1$  and  $R_i = a_i b_i = a_i * b_i + a_i \wedge b_i$  (similarly for  $a_i'^2 = b_i'^2 = -1$ ) are so called *simple rotors* which are connected to either 1 or -1. This holds because  $1 = R_i R_i^\dagger = (a_i * b_i)^2 - (a_i \wedge b_i)^2$ , so we can (see Exercise 6.10) write  $R_i = \pm e^{\phi_i a_i \wedge b_i}$  for some  $\phi_i \in \mathbb{R}$ . Depending on the signature of the plane associated to  $a_i \wedge b_i$ , i.e. on the sign of  $(a_i \wedge b_i)^2 \in \mathbb{R}$ , the set  $e^{\mathbb{R} a_i \wedge b_i} \subseteq \text{Spin}^+$  forms either a circle, a line or a hyperbola. In any case, it goes through the unit element. Finally, for  $s > 1$  or  $t > 1$  we can connect -1 to 1 using for example the circle  $e^{\mathbb{R} e_1 e_2}$ , where  $e_1, e_2$  are two orthonormal basis elements with the same signature.  $\square$

Continuity of the map  $\widetilde{\text{Ad}}$  now implies that the set of rotations represented by rotors, i.e.  $\text{SO}^+$ , forms a connected subgroup containing the identity map (unless the dimension is very low). For euclidean and lorentzian signatures, we have an even simpler situation:

**Theorem 6.16.** *The groups  $\text{Spin}^+(s, t)$  are simply connected for  $(s, t) = (n, 0)$ ,  $(0, n)$ ,  $(1, n)$  or  $(n, 1)$ , where  $n \geq 3$ . Hence, these are the universal covering groups of  $\text{SO}^+(s, t)$ .*

*Proof.* This follows from the fact that, for these signatures,

$$\pi_1(\text{SO}^+(s, t)) = \mathbb{Z}_2 = \ker \widetilde{\text{Ad}}|_{\text{Spin}^+(s, t)}$$

(in general  $\pi_1(\text{SO}^+(s, t)) = \pi_1(\text{SO}(s)) \times \pi_1(\text{SO}(t))$ ; cp. e.g. Theorem 5.7 for the euclidean case), together with the fact that the rotor groups are connected, and the properties of the universal covering groups (see e.g. Theorem VII.6.4 in [41] and references therein).  $\square$

This sums up the situation nicely for higher-dimensional euclidean and lorentzian spaces: The Pin group, which is a double-cover of the orthogonal group, consists of two (euclidean case) or four (lorentzian case) simply connected components. These components are copies of the rotor group, whose structure will be considered in the next subsection. In lower dimensions the structure of all these groups can be worked out in detail; see Section 6.6.

---

**Exercise 6.10.** Verify that if  $R = ab = a * b + a \wedge b$ , with  $a, b \in \mathcal{G}^1$ , and  $RR^\dagger = 1$ , then  $R = \sigma e^{\phi a \wedge b}$  for some  $\phi \in \mathbb{R}$  and  $\sigma \in \{\pm 1\}$ .

## 6.5 The bivector Lie algebra $\mathcal{G}^2$

We have reduced the study of the orthogonal groups to a study of the rotor groups, which in all higher dimensions are connected, smooth Lie groups. An understanding of such groups is provided by their local structure, i.e. their corresponding tangent space Lie algebra, which in this case is simply the space  $\mathcal{G}^2$  of bivectors with a Lie product given by the commutator bracket  $[\cdot, \cdot]$  (recall Exercise 2.21). To see this, take any smooth curve  $R : \mathbb{R} \rightarrow \text{Spin}^+$ ,  $t \rightarrow R(t)$  through the unit element;  $R(0) = 1$ . We claim that the tangent  $R'(0)$  is a bivector. Namely, differentiating the identity  $R(t)R(t)^\dagger = 1$  at  $t = 0$  gives

$$R'(0)^\dagger = -R'(0). \tag{6.13}$$

Furthermore, since  $R(t) \in \mathcal{G}^+$  for all  $t$ , we must have  $R'(0) \in \bigoplus_{k=0,1,\dots} \mathcal{G}^{2+4k}$ . We write  $R'(0) = B + Z$ , where  $B \in \mathcal{G}^2$  and  $Z$  contains no grades lower than 6. Also, because  $R(t) \in \tilde{\Gamma}$  along the curve, we have for any  $v \in V$

$$w(t) := R(t)vR(t)^\dagger \in V \quad \forall t.$$

Again, differentiating this expression and using (6.13) yields

$$V \ni w'(0) = R'(0)v - vR'(0) = -2v \lrcorner R'(0) = -2v \lrcorner B - 2v \lrcorner Z.$$



Inspecting the grades involved in this equation, it follows that  $v \lrcorner Z = 0$  for all  $v \in V$ , and by Lemma 6.10, that  $Z = 0$ .

The commutator product of  $\mathcal{G}^2$  is inherited from the geometric product of  $\text{Spin}^+$ , which through the twisted adjoint action corresponds to taking products of orthogonal transformations. In fact, there is a canonical Lie algebra isomorphism between the bivector space  $\mathcal{G}^2$  and the algebra  $\mathfrak{so}(V, q)$  of antisymmetric transformations of  $V$  (which is the Lie algebra of  $\text{SO}(V, q)$ ), given by (cp. Exercise 3.12)

$$\begin{aligned} \mathfrak{so} &\cong \mathfrak{spin} = \mathcal{G}^2 \\ f = \text{ad}_B &\leftrightarrow B = \frac{1}{2} \sum_{i,j} e^i * f(e^j) e_i \wedge e_j, \end{aligned}$$

where  $\{e_i\}_i$  is any general basis of  $V$ . One verifies, by expanding in the geometric product, that (Exercise 6.11)

$$\frac{1}{2}[e_i \wedge e_j, e_k \wedge e_l] = (e_j * e_k) e_i \wedge e_l - (e_j * e_l) e_i \wedge e_k + (e_i * e_l) e_j \wedge e_k - (e_i * e_k) e_j \wedge e_l. \quad (6.14)$$

*Remark.* Actually, this bivector Lie algebra is more general than it might first seem. Namely, one can show (see e.g. [11] or [12]) that the Lie algebra  $\mathfrak{gl}$  of the general linear group can be represented as a bivector algebra. From the fact that any finite-dimensional Lie algebra has a faithful finite-dimensional representation (Ado's Theorem for characteristic zero, Iwasawa's Theorem for nonzero characteristic, see e.g. [24]) it then follows that *any* finite-dimensional real or complex Lie algebra can be represented as a bivector algebra.

Recall that (see Exercise 2.7), for any choice of norm on  $\mathcal{G}$ , the exponential defined by  $e^x := \sum_{k=0}^{\infty} \frac{x^k}{k!}$  converges and satisfies

$$e^x e^{-x} = e^{-x} e^x = 1$$

and  $x e^x = e^x x$  for any  $x \in \mathcal{G}$ . The following holds for any signature.

**Theorem 6.17.** *For any bivector  $B \in \mathcal{G}^2$  we have that  $\pm e^B \in \text{Spin}^+$ .*

*Proof.* It is obvious that  $\pm e^B$  is an even multivector and that  $e^B (e^B)^\dagger = e^B e^{-B} = 1$ . Hence, it is sufficient to prove that  $\pm e^B \in \Gamma$ , or by Theorem 6.13, that  $e^B V e^{-B} \subseteq V$ .

Let  $W := \{y \in \mathcal{G} : v * y = 0 \ \forall v \in V\} = \bigoplus_{k \neq 1} \mathcal{G}^k$  and choose  $x \in V$  and  $y \in W$ . Define a map  $f : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(t) := (e^{tB} x e^{-tB}) * y.$$

$f$  is obviously a real analytic function with derivatives

$$f'(t) = (B e^{tB} x e^{-tB} - e^{tB} x e^{-tB} B) * y = [B, e^{tB} x e^{-tB}] * y,$$

$$f''(t) = [B, B e^{tB} x e^{-tB} - e^{tB} x e^{-tB} B] * y = [B, [B, e^{tB} x e^{-tB}]] * y,$$

etc. It follows that  $f^{(k)}(0) = \text{ad}_B^k(x) * y = 0$  for all  $k \geq 0$ , since  $\text{ad}_B : V \rightarrow V$  (see Exercise 3.11). Hence,  $f(t) = 0 \ \forall t \in \mathbb{R}$  and  $e^{tB} x e^{-tB} \in V$ .  $\square$

In other words,  $\pm e^{\mathcal{G}^2} \subseteq \text{Spin}^+$ . Actually, the converse inclusion holds for (anti-)euclidean and lorentzian spaces:

**Theorem 6.18.** *For  $(s, t) = (n, 0), (0, n), (1, n)$  or  $(n, 1)$ , we have*

$$\text{Spin}^+(s, t) = \pm e^{\mathcal{G}^2(\mathbb{R}^{s,t})},$$

*i.e. any rotor can be written as (minus) the exponential of a bivector. The minus sign is only required in the lower-dimensional cases  $(0, 0), (1, 0), (0, 1), (1, 1), (1, 2), (2, 1), (1, 3)$  and  $(3, 1)$ .*

The proof can be found in [38]. Essentially, it relies on the fact that any isometry of an euclidean or lorentzian space can be generated by a single infinitesimal transformation. This holds for these spaces *only*, so that for example  $\text{Spin}^+(2, 2) \neq \pm e^{\mathcal{G}^2(\mathbb{R}^{2,2})}$ , where for instance the rotor

$$\pm e_1 e_2 e_3 e_4 e^{\beta(e_1 e_2 + 2e_1 e_4 + e_3 e_4)}, \quad \beta > 0,$$

cannot be reduced to a single exponential; see [28] and [38] pp. 150-152.

Similarly, we see below that even though one can always decompose a bivector into a sum of at most  $\frac{1}{2} \cdot \dim V$  basis blades, the decomposition is particularly nice only in the euclidean and lorentzian cases.

**Theorem 6.19.** *Given any bivector  $B \in \mathcal{G}^2(V)$ , there exists a basis  $\{f_1, \dots, f_n\}$  of  $V$  such that*

$$B = \beta_{12} f_1 \wedge f_2 + \beta_{34} f_3 \wedge f_4 + \dots + \beta_{2k-1, 2k} f_{2k-1} \wedge f_{2k}, \quad (6.15)$$

*where  $2k \leq n$ . If  $V$  is (anti-)euclidean then the basis can be chosen orthonormal, while if  $V$  is (anti-)lorentzian then either the  $\{f_i\}_i$  can be chosen to form an orthonormal basis, or we need to choose  $f_1 = e_0 + e_1$ , where  $\{e_0, e_1, f_2, \dots, f_n\}$  is an orthonormal basis of  $\mathbb{R}^{1,n}$ . For other signatures, such an orthogonal decomposition is not always possible.*

*Proof.* We will first prove the orthogonal decomposition for the euclidean case, and then observe that this also gives the general (weaker) statement. For the lorentzian case and the last statement of the theorem we refer to [38].

We start with a general observation: If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is linear then there exists a subspace  $M \subseteq \mathbb{R}^n$  such that  $T(M) \subseteq M$  and  $1 \leq \dim M \leq 2$ . To see this we start by extending  $T : \mathbb{C}^n \rightarrow \mathbb{C}^n$  and therefore realize that there exists  $\lambda \in \mathbb{C}$  and  $z \in \mathbb{C}^n \setminus \{0\}$  such that  $Tz = \lambda z$  (this follows from the fact that the polynomial  $\det(\lambda \text{id} - T)$  has a zero in  $\mathbb{C}$ , by the fundamental theorem of algebra). Now, write  $\lambda = \sigma + i\tau$  and  $z = x + iy$ , where  $\sigma, \tau \in \mathbb{R}$  and  $x, y \in \mathbb{R}^n$ . Let  $M = \text{Span}_{\mathbb{R}}\{x, y\}$ . We now have that

$$Tx + iTy = T(x + iy) = (\sigma + i\tau)(x + iy) = (\sigma x - \tau y) + i(\tau x + \sigma y),$$

which gives  $Tx = \sigma x - \tau y$  and  $Ty = \tau x + \sigma y$ . Hence,  $T(M) \subseteq M$  and  $\dim M \leq 2$ .

Let us now assume that the signature is euclidean, i.e.  $V = \mathbb{R}^n$ , and prove that, given any bivector  $B \in \mathcal{G}^2$ , there exists an orthonormal basis  $\{f_j\}_{j=1}^n$  such that the decomposition (6.15) holds. Introduce the *rank* of  $B$ ,  $\text{rank}(B)$ , as the least integer  $m$  such that there exists a basis  $\{e_1, \dots, e_n\}$  in which  $B$  can be written

$$B = \sum_{1 \leq i < j \leq m} \beta_{ij} e_i \wedge e_j. \quad (6.16)$$

We prove the theorem by induction on  $\text{rank}(B)$ , i.e. for each fixed  $m \geq 2$  we show that for every  $B \in \mathcal{G}^2$  with  $\text{rank}(B) = m$  there exists an orthonormal basis  $\{f_1, \dots, f_n\}$  such that (6.15) holds.

For  $m = 2$  the statement is trivial (choose an orthonormal basis for the plane  $\overline{e_1 \wedge e_2}$ ), so assume that  $m > 2$  and put  $W = \mathbb{R}e_1 + \dots + \mathbb{R}e_m = \overline{e_1 \wedge \dots \wedge e_m}$  (where  $\{e_i\}_i$  is the basis in (6.16)). But then the map  $W \ni x \mapsto x \lrcorner B \in W$  has a nontrivial invariant subspace  $M$  of dimension  $\leq 2$ .

First assume that  $\dim M = 1$ . Then there exists an orthonormal basis  $\{f_1, \dots, f_m, e_{m+1}, \dots, e_n\}$  of  $V$  such that  $\overline{f_1 \wedge \dots \wedge f_m} = \overline{e_1 \wedge \dots \wedge e_m}$ ,  $M = \mathbb{R}f_1$ , and  $B = \sum_{1 \leq i < j \leq m} \beta'_{ij} f_i f_j$ . But then

$$\lambda f_1 = f_1 \lrcorner B = \beta'_{12} f_2 + \beta'_{13} f_3 + \dots + \beta'_{1m} f_m,$$

implying  $\beta'_{1j} = 0 = \lambda$  for  $1 < j \leq m$ . This means that at most  $m - 1$  basis vectors are required in the expression for  $B$ , contradicting the assumption that  $\text{rank}(B) = m$ .

Hence,  $\dim M = 2$  and we can assume that  $M = \mathbb{R}f_1 + \mathbb{R}f_2$ , where once again  $\{f_1, \dots, f_m, e_{m+1}, \dots, e_n\}$  denotes an orthonormal basis of  $V$  such that  $B = \sum_{1 \leq i < j \leq m} \beta'_{ij} f_i f_j$ . We can write

$$\alpha f_1 + \beta f_2 = f_1 \lrcorner B = \sum_{j=2}^m \beta'_{1j} f_j,$$

so that  $\alpha = 0$  and  $\beta = \beta'_{12}$ . Furthermore, if  $f_2 \lrcorner B = \gamma f_1 + \delta f_2$  then  $\delta = f_2 \lrcorner (f_2 \lrcorner B) = (f_2 \wedge f_2) \lrcorner B = 0$  and

$$\gamma = f_1 \lrcorner (f_2 \lrcorner B) = (f_1 \wedge f_2) \lrcorner B = -(f_2 \wedge f_1) \lrcorner B = -f_2 \lrcorner (f_1 \lrcorner B) = -\beta'_{12}.$$

Thus,  $f_1 \lrcorner B = \beta'_{12} f_2$  and  $f_2 \lrcorner B = -\beta'_{12} f_1$ . If we let  $B' := B - \beta'_{12} f_1 f_2$  we therefore have  $f_1 \lrcorner B' = f_2 \lrcorner B' = 0$  and, writing  $B' = \sum_{1 \leq i < j \leq m} \beta''_{ij} f_i f_j$ , we find

$$0 = f_1 \lrcorner B' = \sum_{1 < j} \beta''_{1j} f_j \quad \Rightarrow \quad \beta''_{1j} = 0 \quad \forall 1 < j \leq m,$$

$$0 = f_2 \lrcorner B' = \sum_{2 < j} \beta''_{2j} f_j \quad \Rightarrow \quad \beta''_{2j} = 0 \quad \forall 2 < j \leq m.$$

Hence,  $B'$  can be expressed solely in terms of the  $m - 2$  basis vectors  $f_3, \dots, f_m$ , i.e.  $\text{rank}(B') \leq m - 2$ . The induction assumption now implies existence of an orthonormal basis on the form  $\{f_1, f_2, g_3, \dots, g_m, e_{m+1}, \dots, e_n\}$  such that  $B' = \sum_{j=3,5,7,\dots,k < m} \gamma_{j,j+1} g_j g_{j+1}$ , and hence that

$$B = \beta'_{12} f_1 f_2 + \gamma_{34} g_3 g_4 + \gamma_{56} g_5 g_6 + \dots + \gamma_{k,k+1} g_k g_{k+1},$$

which proves the theorem in the euclidean case.

For the weaker but signature-independent statement, note that a decomposition of the form (6.15) in terms of a general basis  $\{f_j\}_{j=1}^n$  only concerns the exterior algebra associated to the outer product. Hence, given a bivector  $B = \sum_{1 \leq i < j \leq n} \beta_{ij} e_i \wedge e_j$ , there is no loss in generality in temporarily switching

to euclidean signature (where e.g.  $e_i * e_j = \delta_{ij}$ ), with respect to which one can find an orthonormal basis  $\{f_j = \sum_k \alpha_{jk} e_k\}_{j=1}^n$  such that (6.15) holds. Once we have found such a basis we can switch back to the original signature, and realize that  $\{f_j\}_{j=1}^n$  is of course still a basis (not necessarily orthogonal) and that the expression (6.15) reduces to the original expression for  $B$  upon expressing  $f_j = \sum_k \alpha_{jk} e_k$ .  $\square$

*Remark.* Let us see why the additional requirement is necessary in the lorentzian case. Namely, assume that  $\{e_0, e_1, \dots, e_n\}$  is an orthonormal basis of  $\mathbb{R}^{1,n}$  with  $e_0^2 = 1$ . Consider  $B := (e_0 + e_1) \wedge e_2$ , and assume that  $B$  can also be written

$$B = \beta_1 f_1 f_2 + \beta_3 f_3 f_4 + \dots + \beta_{2p-1} f_{2p-1} f_{2p}, \quad 2p \leq n+1$$

for some other orthonormal basis  $\{f_1, f_2, \dots, f_{n+1}\}$ . Since

$$\begin{aligned} 0 = B \wedge B &= 2\beta_1 \beta_3 f_1 f_2 f_3 f_4 + 2\beta_1 \beta_5 f_1 f_2 f_5 f_6 + \dots \\ &\dots + 2\beta_{2p-3} \beta_{2p-1} f_{2p-3} f_{2p-2} f_{2p-1} f_{2p}, \end{aligned}$$

we must have  $\beta_{2s+1} \beta_{2t+1} = 0$  for all  $s \neq t$ . But then only one term in the sum can survive, and we can assume, e.g.

$$(e_0 + e_1) \wedge e_2 = \beta f_1 f_2, \quad \beta \neq 0.$$

Squaring both sides, we find

$$0 = -(e_0 + e_1)^2 e_2^2 = -\beta^2 f_1^2 f_2^2,$$

so that either  $f_1^2 = 0$  or  $f_2^2 = 0$ , but this contradicts the signature of the space.  $\square$

Another way to state Theorem 6.19 in the euclidean and lorentzian case is that every bivector  $B$  can be written as a sum of commuting 2-blades:

$$B = B_1 + B_2 + \dots + B_k, \quad (6.17)$$

such that  $B_i \in \mathcal{B}_2$ ,  $B_i B_j = B_j B_i \forall i, j$ ,  $\dim \bar{B}_i = 2$ ,  $\bar{B}_i \perp \bar{B}_j \forall i \neq j$ , and  $k \leq n/2$ . It then follows that every rotor  $e^B$  can be written  $e^{B_1} e^{B_2} \dots e^{B_k}$ , where each factor is a simple rotor (as discussed in the proof of Theorem 6.15). The decomposition (6.17) is unique unless  $B_i^2 = B_j^2$  for some  $i \neq j$ .

---

**Exercise 6.11.** Verify (6.14) and observe that, with  $L_{ij} := \frac{1}{2} e_i \wedge e_j$  and the metric tensor  $g_{ij} := e_i * e_j$  (recall that  $\{e_i\}_i$  is any general basis), this leads to the conventional commutation relations

$$[L_{ij}, L_{kl}] = g_{jk} L_{il} - g_{jl} L_{ik} + g_{il} L_{jk} - g_{ik} L_{jl}$$

for the (anti-hermitian) generators of  $\mathfrak{so}(s, t)$ .

**Exercise 6.12.** Use Theorem 6.19 to prove that a bivector  $B \in \mathcal{G}^2(V)$  is a blade if and only if  $B^2$  is a scalar (regardless of the signature of  $V$ ).

**Exercise 6.13.** Construct an example of a homogeneous multivector which squares to a nonzero scalar even though it is not a blade.

**Exercise 6.14.** Show that the *Killing form*,

$$K(A, B) := \text{tr}_{\mathcal{G}^2}(\text{ad}_A \text{ad}_B), \quad A, B \in \mathcal{G}^2,$$

of the Lie algebra  $\mathfrak{spin} = \mathcal{G}^2$  simplifies to

$$K(A, B) = c A * B,$$

and determine the constant  $c$ . (Note that for (anti-)euclidean signatures,  $K \leq 0$ , reflecting the fact that the corresponding Spin groups are compact.)

*Hint:* Use reciprocal bases and Exercise 3.6.

## 6.6 Examples in lower dimensions

We finish this section by considering various examples in lower dimensions, and work out the respective Pin, Spin and rotor groups in each case. We also find that we can use various involutions and extended versions of the norm function (6.7) to determine the corresponding full group of invertible elements.

In the sequel it will be convenient to use the following characterization of the rotor group in lower dimensions as simply being the group of *even unit multivectors*.

**Proposition 6.20.** *If (and only if)  $\dim V \leq 5$  then*

$$\text{Spin}^+ = \{R \in \mathcal{G}^+ : RR^\dagger = 1\}.$$

*Proof.* By definition, the rotor group is a subset of the set on the right hand side in all dimensions, so let us consider the converse inclusion and pick any even-grade multivector  $R$  such that  $RR^\dagger = 1$ . Then  $R$  is invertible and  $R^{-1} = R^\dagger \in \mathcal{G}^+$ . We claim that  $RV R^\dagger \subseteq V$ , so  $R \in \tilde{\Gamma}$  and hence is also a rotor. The claim follows because for any  $v \in V$  the expression  $RvR^\dagger$  is both odd and self-reversing, and hence in  $\mathcal{G}^1 \oplus \mathcal{G}^5$ . This proves the claim and hence the proposition for  $\dim V < 5$ . In the case  $\dim V = 5$ , assume to the contrary that

$$RvR^\dagger = w + \alpha I,$$

with  $w \in V$  and  $\alpha \neq 0$ . Then, since the pseudoscalar commutes with everything,

$$\alpha = \langle RvR^\dagger I^{-1} \rangle_0 = \langle RvI^{-1}R^\dagger \rangle_0 = \langle vI^{-1}R^\dagger R \rangle_0 = \langle vI^{-1} \rangle_0 = 0,$$

which is a contradiction.

The necessity on the dimension is left to the reader as Exercise 6.15.  $\square$

We will also find it useful to extend our set of grade-based involutions with the following generalized ones.

**Definition 6.4.** We define

$$\begin{aligned} [A] &:= (-1)^{|A \neq \emptyset|} A, \\ [A]_{i_1, i_2, \dots, i_k} &:= (-1)^{(|A| \in \{i_1, i_2, \dots, i_k\})} A, \end{aligned}$$

for  $A \in \mathcal{P}(X)$  and extend linearly to  $\mathcal{Cl}(X, R, r)$ .

---

**Exercise 6.15.** Show that Proposition 6.20 cannot be extended to  $\dim V \geq 6$ .

### 6.6.1 The euclidean line

Let us first consider the algebra of the euclidean line,  $\mathcal{G}(\mathbb{R}^1) = \text{Span}_{\mathbb{R}}\{1, e\}$ , where  $e$  is the basis element,  $e^2 = 1$ . This is a commutative algebra with pseudoscalar  $e$ . Because the only unit vectors are  $\{\pm e\}$ , the unit versors resp. rotors form the discrete groups

$$\begin{aligned}\text{Pin}(1, 0) &= \{1, -1, e, -e\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2, \\ \text{Spin}^{(+)}(1, 0) &= \{1, -1\} \cong \mathbb{Z}_2.\end{aligned}$$

Note that  $\underline{\pm 1}$  is the identity map, while  $\underline{\pm e}$  is the unique reflection of the line.

One easily finds the multiplicative group  $\mathcal{G}^\times(\mathbb{R}^1)$  by considering the norm function, which with a one-dimensional vector space is given by

$$N_1(x) := x^\square x = x^* x.$$

For an arbitrary element  $x = \alpha + \beta e$  then

$$N_1(x) = (\alpha - \beta e)(\alpha + \beta e) = \alpha^2 - \beta^2 \in \mathbb{R}.$$

When  $N_1(x) \neq 0$  we find that  $x$  has an inverse  $x^{-1} = \frac{1}{N_1(x)} x^* = \frac{1}{\alpha^2 - \beta^2}(\alpha - \beta e)$ . Hence,

$$\mathcal{G}^\times(\mathbb{R}^1) = \{x \in \mathcal{G} : N_1(x) \neq 0\} = \{\alpha + \beta e \in \mathcal{G} : \alpha^2 \neq \beta^2\}.$$

Note also that  $N_1(xy) = x^* y^* x y = N_1(x) N_1(y)$  for all  $x, y \in \mathcal{G}$  since the algebra is commutative.

### 6.6.2 The anti-euclidean line

The algebra of the anti-euclidean line,  $\mathcal{G}(\mathbb{R}^{0,1}) = \text{Span}_{\mathbb{R}}\{1, i\} \cong \mathbb{C}$ , is of course also a commutative algebra, but, unlike the previous example, this is even a *field* since every nonzero element is invertible. The norm function is an actual norm (squared) in this case,

$$N_1(\alpha + \beta i) = (\alpha - \beta i)(\alpha + \beta i) = \alpha^2 + \beta^2 \in \mathbb{R}^+,$$

namely the modulus of the complex number. We have already noted that the grade involution represents the complex conjugate and, as above,  $x^{-1} = \frac{1}{N_1(x)} x^*$ . The relevant groups are

$$\begin{aligned}\mathcal{G}^\times(\mathbb{R}^{0,1}) &= \mathcal{G} \setminus \{0\} \cong \mathbb{C}^\times, \\ \text{Pin}(0, 1) &= \{1, -1, i, -i\} \cong \mathbb{Z}_4, \\ \text{Spin}^{(+)}(0, 1) &= \{1, -1\} \cong \mathbb{Z}_2.\end{aligned}$$

### 6.6.3 The degenerate line

We also take a brief look at the simplest example of a degenerate algebra,  $\mathcal{G}(\mathbb{R}^{0,0,1})$ , just to see what happens in such a situation. Let the vector  $n$  span a one-dimensional space with quadratic form  $q = 0$ . Then

$$\mathcal{G}(\mathbb{R}^{0,0,1}) = \text{Span}_{\mathbb{R}}\{1, n\} \cong \bigwedge^* \mathbb{R}^1,$$

and  $n^2 = 0$ . The norm function depends only on the scalar part in this case,

$$N_1(\alpha + \beta n) = (\alpha - \beta n)(\alpha + \beta n) = \alpha^2 \in \mathbb{R}^+.$$

Hence, an element is invertible if and only if the scalar part is nonzero;

$$\mathcal{G}^\times(\mathbb{R}^{0,0,1}) = \{\alpha + \beta n \in \mathcal{G} : \alpha \neq 0\}.$$

Since no vectors are invertible, we are left with only the empty product in the versor group, i.e.  $\Gamma = \{1\}$ . Note, however, that for  $\alpha \neq 0$

$$(\alpha + \beta n)^* n (\alpha + \beta n)^{-1} = (\alpha - \beta n) n \frac{1}{\alpha^2} (\alpha - \beta n) = n,$$

so the Lipschitz group in this case is

$$\tilde{\Gamma} = \mathcal{G}^\times \neq \Gamma.$$

This shows that the assumption on nondegeneracy was necessary in the discussion about the Lipschitz group in Section 6.3.

#### 6.6.4 The euclidean plane

Let  $\{e_1, e_2\}$  be an orthonormal basis of  $\mathbb{R}^2$  and consider the plane algebra

$$\mathcal{G}(\mathbb{R}^2) = \text{Span}_{\mathbb{R}}\{1, e_1, e_2, I = e_1 e_2\}.$$

We recall that the even subalgebra is  $\mathcal{G}^+(\mathbb{R}^2) \cong \mathcal{G}(\mathbb{R}^{0,1}) \cong \mathbb{C}$  and that the rotor group corresponds to the group of unit complex numbers,

$$\text{Spin}^{(+)}(2, 0) = e^{\mathbb{R}I} \cong \text{U}(1).$$

Furthermore,

$$\underline{e^{\varphi I}}(e_1) = e^{\varphi I} e_1 e^{-\varphi I} = e_1 e^{-2\varphi I} = e_1 (\cos 2\varphi - I \sin 2\varphi) = \cos 2\varphi e_1 - \sin 2\varphi e_2,$$

so a rotor on the form  $\pm e^{-\frac{\varphi}{2}I}$  represents a counter-clockwise<sup>16</sup> rotation in the plane by an angle  $\varphi$ . The Pin group is found by picking any unit vector  $e \in S^1$ :

$$\text{Pin}(2, 0) = e^{\mathbb{R}I} \bigsqcup e^{\mathbb{R}I} e \cong \text{U}(1) \bigsqcup S^1,$$

i.e. two copies of a unit circle, where we write  $\bigsqcup$  to emphasize a *disjoint* union. An element  $\pm e^{-\frac{\varphi}{2}I} e \in S^1$  of the second circle corresponds to a reflection along  $e$  (which here serves as a reference axis) followed by a rotation by  $\varphi$  in the plane.

Let us next determine the full multiplicative group of the plane algebra. For algebras over two-dimensional spaces we use the original norm function

$$N_2(x) := x^\square x$$

since it satisfies  $N_2(x)^\square = N_2(x)$  for all  $x \in \mathcal{G}$ . The properties of the involutions in Table 2.1 then require this to be a scalar, so we have a map

$$N_2: \mathcal{G} \rightarrow \mathcal{G}^0 = \mathbb{R}.$$

<sup>16</sup>Assuming, of course, that  $e_1$  points at 3 o'clock and  $e_2$  at 12 o'clock.

For an arbitrary element  $x = \alpha + a_1e_1 + a_2e_2 + \beta I \in \mathcal{G}$ , we explicitly find

$$\begin{aligned} N_2(x) &= (\alpha - a_1e_1 - a_2e_2 - \beta I)(\alpha + a_1e_1 + a_2e_2 + \beta I) \\ &= \alpha^2 - a_1^2 - a_2^2 + \beta^2. \end{aligned}$$

Furthermore,  $N_2(x^\square) = N_2(x)$  and  $N_2(xy) = y^\square N_2(x)y = N_2(x)N_2(y)$  for all  $x, y \in \mathcal{G}$ . Proceeding as in the one-dimensional case, we find that  $x$  has an inverse  $x^{-1} = \frac{1}{N_2(x)}x^\square$  if and only if  $N_2(x) \neq 0$ , i.e.

$$\begin{aligned} \mathcal{G}^\times(\mathbb{R}^2) &= \{x \in \mathcal{G} : N_2(x) \neq 0\} \\ &= \{\alpha + a_1e_1 + a_2e_2 + \beta I \in \mathcal{G} : \alpha^2 + \beta^2 \neq a_1^2 + a_2^2\}. \end{aligned}$$

### 6.6.5 The anti-euclidean plane

In the case of the anti-euclidean plane,  $\mathcal{G}(\mathbb{R}^{0,2}) \cong \mathbb{H}$ , the norm function  $N_2$  has similar properties as in the euclidean algebra, except that it now once again represents the square of an actual norm, namely the *quaternion norm*,

$$N_2(x) = \alpha^2 + a_1^2 + a_2^2 + \beta^2.$$

Just as for the complex numbers then, all nonzero elements are invertible,

$$\mathcal{G}^\times(\mathbb{R}^{0,2}) = \{x \in \mathcal{G} : N_2(x) \neq 0\} = \mathcal{G} \setminus \{0\}.$$

The even subalgebra is also in this case isomorphic to the complex numbers, and the Pin, Spin and rotor groups are analogous to the euclidean case.

### 6.6.6 The lorentzian plane

With the standard basis  $\{e_0, e_1\}$  of  $\mathbb{R}^{1,1}$ , the two-dimensional lorentzian algebra is given by

$$\mathcal{G}(\mathbb{R}^{1,1}) = \text{Span}_{\mathbb{R}}\{1, e_0, e_1, I = e_1e_0\},$$

where  $e_0$  is interpreted as a time direction in physical applications, while  $e_1$  defines a spatial direction. In general, a vector (or blade)  $v$  is called *timelike* if  $v^2 > 0$ , *spacelike* if  $v^2 < 0$ , and *lightlike* or *null* if  $v^2 = 0$ .

The multiplicative group is like previously given by

$$\begin{aligned} \mathcal{G}^\times(\mathbb{R}^2) &= \{x \in \mathcal{G} : N_2(x) \neq 0\} \\ &= \{\alpha + a_0e_0 + a_1e_1 + \beta I \in \mathcal{G} : \alpha^2 - a_0^2 + a_1^2 - \beta^2 \neq 0\}. \end{aligned}$$

However, the pseudoscalar<sup>17</sup>  $I = e_1e_0$  squares to the identity in this case and the even subalgebra is therefore  $\mathcal{G}^+(\mathbb{R}^{1,1}) \cong \mathcal{G}(\mathbb{R}^1)$ . This has as an important consequence that the rotor group is fundamentally different from the (anti-)euclidean case,

$$\text{Spin}^+(1,1) = \{R = \alpha + \beta I \in \mathcal{G}^+ : \alpha^2 - \beta^2 = 1\} = \pm e^{\mathbb{R}I},$$

(recall Exercise 6.10, or Theorem 6.18). Thus, the set of rotors can be understood as a pair of disjoint hyperbolas passing through the points 1 and -1,

<sup>17</sup>The choice of orientation of  $I$  here corresponds to the fact that space-time diagrams are conventionally drawn with the time axis vertical.






Figure 6.1: A Lorentz-boosted vector  $\underline{R}(e_0)$ , where  $R = \pm e^{\alpha I/2}$  and  $\alpha > 0$ .

respectively. The rotations that are represented by rotors of this form are called *Lorentz boosts*. Note the hyperbolic nature of these rotations,

$$\begin{aligned} e^{\alpha I} e_0 e^{-\alpha I} &= e_0 e^{-2\alpha I} = e_0 (\cosh 2\alpha - I \sinh 2\alpha) \\ &= \cosh 2\alpha e_0 + \sinh 2\alpha e_1, \\ e^{\alpha I} e_1 e^{-\alpha I} &= \cosh 2\alpha e_1 + \sinh 2\alpha e_0. \end{aligned}$$

Hence, a rotor  $\pm e^{\frac{\alpha}{2} I}$  transforms, or *boosts*, timelike vectors by a hyperbolic angle  $\alpha$  in the positive spacelike direction (see Figure 6.1).

The Spin group consists of four hyperbolas and the Pin group of eight,

$$\begin{aligned} \text{Spin}(1, 1) &= \pm e^{\mathbb{R}I} \sqcup \pm e^{\mathbb{R}I} I, \\ \text{Pin}(1, 1) &= \pm e^{\mathbb{R}I} \sqcup \pm e^{\mathbb{R}I} e_0 \sqcup \pm e^{\mathbb{R}I} e_1 \sqcup \pm e^{\mathbb{R}I} I, \end{aligned}$$

Note that the action of the rotor  $I = e_1 e_0$  in the second pair of components of the Spin group is a reflection both in the time and the space direction;

$$\underline{I}(x) = IxI^{-1} = x^* = (-\text{id})_{\wedge}(x), \quad x \in \mathcal{G}.$$

Hence, it preserves the total orientation of the space, but it cannot be smoothly transformed into to an identity transformation because of the hyperbolic nature of the space, or intuitively, because spacelike and timelike vectors cannot be boosted across the light-cone, reflecting the physical interpretation in relativity that nothing can travel faster than light. The two additional pairs of components of the Pin group correspond to a reflection in the time resp. space direction, followed by a boost.

### 6.6.7 The space algebra

We write an arbitrary element  $x \in \mathcal{G}(\mathbb{R}^3)$  of the space algebra as

$$x = \alpha + \mathbf{a} + \mathbf{b}I + \beta I, \quad (6.18)$$

where  $\alpha, \beta \in \mathbb{R}$  and  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ . The norm function  $N_2$  does not take values in  $\mathbb{R}$  in this algebra, but due to the properties of the Clifford conjugate we have  $N_2(x) = N_2(x)^{\square} \in \mathcal{G}^0 \oplus \mathcal{G}^3$ . This subalgebra is isomorphic to  $\mathbb{C}$ , and its corresponding complex conjugate is given by  $[x]_3$  (or  $[x]$ ). Using these properties, we can construct a real-valued map  $N_3: \mathcal{G} \rightarrow \mathbb{R}^+$  by taking the complex modulus,

$$N_3(x) := [N_2(x)]_3 N_2(x) = [x^{\square} x] x^{\square} x. \quad (6.19)$$

Plugging in (6.18) we obtain

$$N_2(x) = \alpha^2 - \mathbf{a}^2 + \mathbf{b}^2 - \beta^2 + 2(\alpha\beta - \mathbf{a} * \mathbf{b})I,$$

and

$$N_3(x) = (\alpha^2 - \mathbf{a}^2 + \mathbf{b}^2 - \beta^2)^2 + 4(\alpha\beta - \mathbf{a} * \mathbf{b})^2. \quad (6.20)$$

Although  $N_3$  takes values in  $\mathbb{R}^+$ , it is not a norm<sup>18</sup> on  $\mathcal{G}$  since there are nonzero elements with  $N_3(x) = 0$ . It does however have the multiplicative property

$$\begin{aligned} N_3(xy) &= [N_2(xy)]_3 N_2(xy) = [N_2(x)]_3 [N_2(y)]_3 N_2(x) N_2(y) \\ &= N_3(x) N_3(y), \end{aligned} \quad (6.21)$$

for all  $x, y \in \mathcal{G}$ , since  $N_2(x)$  commutes with all of  $\mathcal{G}$ , and therefore

$$N_2(xy) = (xy)^\square xy = y^\square N_2(x)y = N_2(x)N_2(y).$$

We also observe from (6.20) that  $N_3(x^\square) = N_3(x)$ . The expression (6.19) singles out the invertible elements as those elements (6.18) for which  $N_3(x) \neq 0$ , i.e.

$$\mathcal{G}^\times(\mathbb{R}^3) = \{x \in \mathcal{G} : (\alpha^2 - \mathbf{a}^2 + \mathbf{b}^2 - \beta^2)^2 + 4(\alpha\beta - \mathbf{a} * \mathbf{b})^2 \neq 0\},$$

and then  $x^{-1} = \frac{1}{N_3(x)} [x^\square x] x^\square$ .

The rotor group of the space algebra is, according to the isomorphism  $\mathcal{G}^+(\mathbb{R}^{3,0}) \cong \mathbb{H}$  and Proposition 6.20, the group of unit quaternions (note that the reverse here acts as the quaternion conjugate),

$$\text{Spin}^{(+)}(3, 0) = \{\alpha + \mathbf{b}I \in \mathcal{G}^+ : \alpha^2 + \mathbf{b}^2 = 1\} = e^{\mathcal{G}^2(\mathbb{R}^3)}. \quad (6.22)$$

The exponentiation of the bivector algebra in the second identity followed from Theorem 6.18. An arbitrary rotor  $R$  can according to (6.22) be written in the polar form  $R = e^{\varphi \mathbf{n}I}$ , where  $\mathbf{n} \in S^2$  is a unit vector, and represents (cp. e.g. (1.2)-(1.3)) a rotation by an angle  $-2\varphi$  in the plane  $\mathbf{n}I = -\mathbf{n}^c$ , i.e.  $2\varphi$  clockwise around the axis  $\mathbf{n}$ .

The Pin group consists of two copies of the rotor group,

$$\text{Pin}(3, 0) = e^{\mathcal{G}^2(\mathbb{R}^3)} \bigsqcup e^{\mathcal{G}^2(\mathbb{R}^3)} \mathbf{n},$$

for any unit vector  $\mathbf{n} \in S^2$ . The Pin group can be understood topologically as a pair of unit 3-spheres  $S^3$  lying in the even and odd subspaces, respectively. The odd one includes an additional action of a reflection and corresponds to the non-orientation-preserving part of  $O(3)$ .

By the isomorphism  $\rho$  in Example 5.2 we can also represent the spatial rotor group in terms of the complex matrices

$$\rho(R) = \begin{bmatrix} \alpha + b_3 i & b_2 + b_1 i \\ -b_2 + b_1 i & \alpha - b_3 i \end{bmatrix}.$$

We find that the hermitian conjugate of such a matrix is  $\rho(R)^\dagger = \rho(R)^{-1}$  and  $\det_{\mathbb{C}} \rho(R) = 1$ , and hence that these form the *special unitary group* of  $\mathbb{C}^2$ , i.e.

$$\text{Spin}^{(+)}(3, 0) \cong \text{SU}(2).$$

This is the representation of the spatial rotor group which is most often encountered in physics, in conjunction with spin in quantum mechanics.

<sup>18</sup>Actually, it is not possible to find a norm  $|\cdot|$  on  $\mathcal{G}(\mathbb{R}^{s,t})$ ,  $s+t \geq 3$ , satisfying  $|xy| = |x||y|$ , as will be clear from Theorem 8.5 and the remark following it.

**Exercise 6.16.** Show that if  $x \in \mathcal{G}^1(\mathbb{R}^3) \oplus \mathcal{G}^2(\mathbb{R}^3)$  is noninvertible, then it is on the form

$$x = \alpha e(1 + \mathbf{f}),$$

where  $\alpha \in \mathbb{R}$  and  $e, \mathbf{f} \in S^2$  are orthogonal unit vectors.

### 6.6.8 The spacetime algebra

As a four-dimensional example we consider the *spacetime algebra* (STA), which is the geometric algebra of *Minkowski spacetime*,  $\mathbb{R}^{1,3}$ . By convention<sup>19</sup>, we denote an orthonormal basis of the Minkowski space by  $\{\gamma_0, \gamma_1, \gamma_2, \gamma_3\}$ , where  $\gamma_0$  is timelike and the other  $\gamma_i$  are spacelike. The STA expressed in this basis is

$$\begin{aligned} \mathcal{G}(\mathbb{R}^{1,3}) = \\ \text{Span}_{\mathbb{R}}\{1, \gamma_0, \gamma_1, \gamma_2, \gamma_3, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 I, \mathbf{e}_2 I, \mathbf{e}_3 I, \gamma_0 I, \gamma_1 I, \gamma_2 I, \gamma_3 I, I\}, \end{aligned}$$

where the pseudoscalar is  $I := \gamma_0 \gamma_1 \gamma_2 \gamma_3$  and we set  $\mathbf{e}_i := \gamma_i \gamma_0$ ,  $i = 1, 2, 3$ . The form of the STA basis chosen above emphasizes the duality which exists between the graded subspaces. It also hints that the even subalgebra of the STA is the space algebra. This is actually the case since there is an isomorphism  $\mathcal{G}^+(\mathbb{R}^{1,3}) \cong \mathcal{G}(\mathbb{R}^{3,0})$ , but we can also verify this explicitly by noting that  $\mathbf{e}_i^2 = 1$  and  $\mathbf{e}_i \mathbf{e}_j = -\mathbf{e}_j \mathbf{e}_i$ ,  $i \neq j$ , as well as  $\mathbf{e}_1 \mathbf{e}_2 \mathbf{e}_3 = I$ . Hence, the timelike (positive square) blades  $\{\mathbf{e}_i\}$  form a basis of a 3-dimensional euclidean space called the *relative space* to  $\gamma_0$ . Given any timelike vector  $a$  we can find a similar relative space spanned by the bivectors  $\{b \wedge a\}$  for  $b \in \mathbb{R}^{1,3}$ . These spaces all generate the *relative space algebra*  $\mathcal{G}^+$ , and only the precise split between the vector and bivector part of this relative algebra depends on the chosen reference vector  $a$ .

Using boldface to denote relative space elements, an arbitrary multivector  $x \in \mathcal{G}$  can be written

$$x = \alpha + a + \mathbf{a} + \mathbf{b}I + bI + \beta I, \quad (6.23)$$

where  $\alpha, \beta \in \mathbb{R}$ ,  $a, b \in \mathbb{R}^{1,3}$  and  $\mathbf{a}, \mathbf{b}$  in relative space  $\mathbb{R}^3$ . As in previous examples, we would like to find the invertible elements. Looking at the norm function  $N_2: \mathcal{G} \rightarrow \mathcal{G}^0 \oplus \mathcal{G}^3 \oplus \mathcal{G}^4$ , it is not obvious that we can extend this to a real-valued function on  $\mathcal{G}$ . Fortunately, we have for  $X = \alpha + bI + \beta I \in \mathcal{G}^0 \oplus \mathcal{G}^3 \oplus \mathcal{G}^4$  that

$$X[X]_{3,4} = [X]_{3,4}X = (\alpha - bI - \beta I)(\alpha + bI + \beta I) = \alpha^2 - b^2 + \beta^2 \in \mathbb{R}. \quad (6.24)$$

Hence, we can define a map  $N_4: \mathcal{G} \rightarrow \mathbb{R}$  by

$$N_4(x) := [N_2(x)]_{3,4}N_2(x) = [x^\square x]x^\square x. \quad (6.25)$$

Plugging in (6.23) into  $N_2$ , we obtain after some simplifications

$$\begin{aligned} N_2(x) = & \alpha^2 - a^2 - \mathbf{a}^2 + \mathbf{b}^2 + b^2 - \beta^2 \\ & + 2(\alpha b - \beta a - a \lrcorner \mathbf{b} + b \lrcorner \mathbf{a} - a \lrcorner \mathbf{a}^c - b \lrcorner \mathbf{b}^c)I \\ & + 2(\alpha\beta - a * b - \mathbf{a} * \mathbf{b})I \end{aligned} \quad (6.26)$$

<sup>19</sup>This choice of notation is motivated by the Dirac representation of the STA in terms of so called gamma matrices which is conventionally used by physicists.

and hence, by (6.24),

$$\begin{aligned}
N_4(x) &= (\alpha^2 - a^2 - \mathbf{a}^2 + \mathbf{b}^2 + b^2 - \beta^2)^2 \\
&\quad - 4(\alpha b - \beta a - a \lrcorner \mathbf{b} + b \lrcorner \mathbf{a} - a \lrcorner \mathbf{a}^c - b \lrcorner \mathbf{b}^c)^2 \\
&\quad + 4(\alpha\beta - a * b - \mathbf{a} * \mathbf{b})^2.
\end{aligned} \tag{6.27}$$

We will prove some rather non-trivial statements about this norm function where we need that  $[xy]x = x[yx]$  for all  $x, y \in \mathcal{G}$ . This is a quite general property of this involution.

**Lemma 6.21.** *In any Clifford algebra  $Cl(X, R, r)$  (even when  $X$  is infinite), we have*

$$[xy]x = x[yx] \quad \forall x, y \in Cl.$$

*Proof.* By linearity, we can take  $y = A \in \mathcal{P}(X)$  and expand  $x$  in coordinates  $x_B \in R$  as  $x = \sum_{B \in \mathcal{P}(X)} x_B B$ . We obtain

$$\begin{aligned}
x[Ax] &= \sum_{B,C} x_B x_C B[AC] \\
&= \sum_{B,C} x_B x_C ((A \Delta C = \emptyset) - (A \Delta C \neq \emptyset)) BAC \\
&= \sum_{B,C} x_B x_C ((A = C) - (A \neq C)) BAC \\
&= \sum_B x_B x_A BAA - \sum_{C \neq A} \sum_B x_B x_C BAC
\end{aligned}$$

and

$$\begin{aligned}
[xA]x &= \sum_{B,C} x_B x_C [BA]C \\
&= \sum_{B,C} x_B x_C ((B = A) - (B \neq A)) BAC \\
&= \sum_C x_A x_C AAC - \sum_{B \neq A} \sum_C x_B x_C BAC \\
&= x_A^2 AAA + \sum_{C \neq A} x_A x_C AAC - \sum_{B \neq A} x_B x_A \underbrace{BAA}_{AAB} \\
&\quad - \sum_{B \neq A} \sum_{C \neq A} x_B x_C BAC \\
&= x_A^2 AAA - \sum_{B \neq A} \sum_{C \neq A} x_B x_C BAC \\
&= x[Ax].
\end{aligned}$$

□

We now have the following

**Lemma 6.22.**  $N_4(x^\square) = N_4(x)$  for all  $x \in \mathcal{G}(\mathbb{R}^{1,3})$ .

(Note that this identity is not at all obvious from the expression (6.27).)

*Proof.* Using Lemma 6.21 we have that

$$N_4(x^\square) = [xx^\square]x^\square = x[x^\square x]x^\square.$$

Since  $N_4$  takes values in  $\mathbb{R}$ , this must be a scalar, so that

$$N_4(x^\square) = \langle x[x^\square x]x^\square \rangle_0 = \langle [x^\square x]x^\square x \rangle_0 = \langle N_4(x) \rangle_0 = N_4(x),$$

where we used the symmetry of the scalar product. □

**Lemma 6.23.** For all  $X, Y \in \mathcal{G}^0 \oplus \mathcal{G}^3 \oplus \mathcal{G}^4(\mathbb{R}^{1,3})$  we have

$$[XY] = [Y][X].$$

*Proof.* Take arbitrary elements  $X = \alpha + bI + \beta I$  and  $Y = \alpha' + b'I + \beta'I$ . Then

$$\begin{aligned} [XY] &= [(\alpha + bI + \beta I)(\alpha' + b'I + \beta'I)] \\ &= \alpha\alpha' - \alpha b'I - \alpha\beta'I - bI\alpha' + b * b' - b \wedge b' + b\beta' - \beta I\alpha' - \beta b' - \beta\beta' \end{aligned}$$

and

$$\begin{aligned} [Y][X] &= (\alpha' - b'I - \beta'I)(\alpha - bI - \beta I) \\ &= \alpha'\alpha - \alpha'bI - \alpha'\beta I - b'I\alpha + b' * b + b' \wedge b - b'\beta - \beta'I\alpha + \beta'b - \beta'\beta. \end{aligned}$$

Comparing these expressions we find that they are equal.  $\square$

We can now prove that  $N_4$  actually acts as a determinant on the STA.

**Theorem 6.24.** The norm function  $N_4$  satisfies the multiplicative property

$$N_4(xy) = N_4(x)N_4(y) \quad \forall x, y \in \mathcal{G}(\mathbb{R}^{1,3}).$$

*Proof.* Using that  $N_4(xy)$  is a scalar, and that  $N_2$  takes values in  $\mathcal{G}^0 \oplus \mathcal{G}^3 \oplus \mathcal{G}^4$ , we obtain

$$\begin{aligned} N_4(xy) &= \langle [(xy)^\square xy](xy)^\square xy \rangle_0 = \langle [y^\square x^\square xy]y^\square x^\square xy \rangle_0 \\ &= \langle x^\square xy[y^\square x^\square xy]y^\square \rangle_0 = \langle x^\square x[yy^\square x^\square x]yy^\square \rangle_0 \\ &= \langle N_2(x)[N_2(y^\square)N_2(x)]N_2(y^\square) \rangle_0 \\ &= \langle N_2(x)[N_2(x)][N_2(y^\square)]N_2(y^\square) \rangle_0 \\ &= \langle N_4(x)N_4(y^\square) \rangle_0 = N_4(x)N_4(y^\square), \end{aligned}$$

where we applied Lemma 6.21 and then Lemma 6.23. Finally, Lemma 6.22 gives the claimed identity.  $\square$

From (6.25) we find that the multiplicative group of the STA is given by

$$\mathcal{G}^\times(\mathbb{R}^{1,3}) = \{x \in \mathcal{G} : N_4(x) \neq 0\}$$

and that the inverse of  $x \in \mathcal{G}^\times$  is

$$x^{-1} = \frac{1}{N_4(x)}[x^\square x]x^\square.$$

Note that the above theorems regarding  $N_4$  only rely on the commutation properties of the different graded subspaces and not on the actual signature and field of the vector space.

Let us now turn our attention to the rotor group of the STA. The reverse equals the Clifford conjugate on the even subalgebra (it also corresponds to the Clifford conjugate defined on the relative space), so we find from (6.26) that the rotor group is

$$\begin{aligned} \text{Spin}^+(1, 3) &= \{x \in \mathcal{G}^+ : N_2(x) = x^\square x = 1\} \\ &= \{\alpha + \mathbf{a} + \mathbf{b}I + \beta I \in \mathcal{G}^+ : \alpha^2 - \mathbf{a}^2 + \mathbf{b}^2 - \beta^2 = 1 \text{ and } \alpha\beta = \mathbf{a} * \mathbf{b}\} \\ &= \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \cong \text{SL}(2, \mathbb{C}). \end{aligned}$$

The last isomorphism is related to the Dirac representation of the STA, while the exponentiation identity was obtained from Theorem 6.18 and gives a better picture of what the rotor group looks like. Namely, any rotor  $R$  can be written  $R = \pm e^{\mathbf{a} + \mathbf{b}I}$  for some relative vectors  $\mathbf{a}, \mathbf{b}$ , or by Theorem 6.19, either

$$R = e^{\alpha \mathbf{e}} e^{\varphi \mathbf{e}I},$$

where  $\alpha, \varphi \in \mathbb{R}$  and  $\mathbf{e} = f_0 \wedge f_1$  is a timelike unit blade, or

$$R = \pm e^{n \wedge f_2} = \pm e^{\alpha \mathbf{e}(1 + \mathbf{f})} = \pm 1 \pm \alpha \mathbf{e}(1 + \mathbf{f}),$$

with a null vector  $n$  and anticommuting timelike unit blades  $\mathbf{e}, \mathbf{f}$  (cp. Exercise 6.16). A simple rotor of the form  $e^{\mathbf{b}I}$  corresponds to a rotation in the spacelike plane  $\mathbf{b}^c$  with angle  $2|\mathbf{b}|$  (which is a rotation also in relative space), while  $e^{\mathbf{a}}$  corresponds to a hyperbolic rotation in the timelike plane  $\mathbf{a}$ , i.e. a boost in the relative space direction  $\mathbf{a}$  with hyperbolic angle  $2|\mathbf{a}|$ .

Picking a timelike and a spacelike unit reference vector, e.g.  $\gamma_0$  and  $\gamma_1$ , we obtain the Spin and Pin groups,

$$\text{Spin}(1, 3) = \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \sqcup \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \gamma_0 \gamma_1,$$

$$\text{Pin}(1, 3) = \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \sqcup \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \gamma_0 \sqcup \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \gamma_1 \sqcup \pm e^{\mathcal{G}^2(\mathbb{R}^{1,3})} \gamma_0 \gamma_1,$$

The Pin group forms a double-cover of the so called *Lorentz group*  $O(1,3)$ . Since the STA rotor group is connected, we find that  $O(1,3)$  has four connected components. Two of these are not in Spin and correspond to a single inversion of time resp. space. The Spin group covers the subgroup of *proper* Lorentz transformations preserving the total orientation, while the rotor group covers the connected, *proper orthochronous* Lorentz group  $SO^+(1, 3)$ , which also preserves the direction of time. The physical interpretations of the spacetime algebra will be further discussed in Section 11.

### 6.6.9 \*The Dirac algebra

Due to historic reasons, the *Dirac algebra*  $\mathcal{G}(\mathbb{C}^4) \cong \mathcal{G}(\mathbb{R}^{4,1})$  is actually the representation of the STA which is most commonly used in physical applications. The relation between these complex respectively real algebras is observed by noting that the pseudoscalar in  $\mathcal{G}(\mathbb{R}^{4,1})$  commutes with all elements and squares to minus the identity. By Proposition 5.3 we then find that the Dirac algebra is the complexification of the STA,

$$\mathcal{G}(\mathbb{R}^{4,1}) \cong \mathcal{G}(\mathbb{R}^{1,3}) \otimes \mathbb{C} \cong \mathcal{G}(\mathbb{C}^4) \cong \mathbb{C}^{4 \times 4}.$$

We construct this isomorphism explicitly by taking bases  $\{\gamma_0, \gamma_1, \gamma_2, \gamma_3\}$  of  $\mathbb{R}^{1,3}$  as above, and  $\{e_0, \dots, e_4\}$  of  $\mathbb{R}^{4,1}$  such that  $e_0^2 = -1$  and the other  $e_j^2 = 1$ . We write  $\mathcal{G}_5 := \mathcal{G}(\mathbb{R}^{4,1})$  and  $\mathcal{G}_4^{\mathbb{C}} := \mathcal{G}(\mathbb{R}^{1,3}) \otimes \mathbb{C}$ , and use the convention that Greek indices run from 0 to 3. The isomorphism  $F: \mathcal{G}_5 \rightarrow \mathcal{G}_4^{\mathbb{C}}$  is given by the following

1-to-1 correspondence of basis elements:

$\mathcal{G}_4^{\mathbb{C}}$	:	$1 \otimes 1$	$\gamma_\mu \otimes 1$	$\gamma_\mu \wedge \gamma_\nu \otimes 1$	$\gamma_\mu \wedge \gamma_\nu \wedge \gamma_\lambda \otimes 1$	$I_4 \otimes 1$	$1 \otimes i$
$\mathcal{G}_5$	:	$1$	$e_\mu e_4$	$-e_\mu \wedge e_\nu$	$-e_\mu \wedge e_\nu \wedge e_\lambda e_4$	$e_0 e_1 e_2 e_3$	$I_5$
$x^\square$ in $\mathcal{G}_4^{\mathbb{C}}$	:	+	-	-	+	+	+
$[x]$ in $\mathcal{G}_4^{\mathbb{C}}$	:	+	-	-	-	-	+
$\bar{x}$ in $\mathcal{G}_4^{\mathbb{C}}$	:	+	+	+	+	+	-

The respective pseudoscalars are  $I_4 := \gamma_0 \gamma_1 \gamma_2 \gamma_3$  and  $I_5 := e_0 e_1 e_2 e_3 e_4$ . We have also noted the correspondence between involutions in the different algebras. Clifford conjugate in  $\mathcal{G}_4^{\mathbb{C}}$  corresponds to reversion in  $\mathcal{G}_5$ , the  $[\cdot]$ -involution becomes the  $[\cdot]_{1,2,3,4}$ -involution, while complex conjugation in  $\mathcal{G}_4^{\mathbb{C}}$  corresponds to grade involution in  $\mathcal{G}_5$ . In other words,

$$F(x^\dagger) = F(x)^\square, \quad F([x]_{1,2,3,4}) = [F(x)], \quad F(x^*) = \overline{F(x)}.$$

We can use the correspondence above to find a norm function on  $\mathcal{G}_5$ . Since the properties of  $N_4 : \mathcal{G}(\mathbb{R}^{1,3}) \rightarrow \mathbb{R}$  are actually independent of the choice of field, we have that the complexification of  $N_4$  satisfies

$$N_4^{\mathbb{C}} : \mathcal{G}(\mathbb{C}^4) \rightarrow \mathbb{C},$$

$$x \mapsto [x^\square x] x^\square x.$$

Taking the modulus of this complex number, we arrive at a real-valued map  $N_5 : \mathcal{G}(\mathbb{R}^{4,1}) \rightarrow \mathbb{R}$  with

$$\begin{aligned} N_5(x) &:= \overline{N_4^{\mathbb{C}}(F(x))} N_4^{\mathbb{C}}(F(x)) \\ &= \overline{[F(x)^\square F(x)] F(x)^\square F(x)} [F(x)^\square F(x)] F(x)^\square F(x) \\ &= [[x^\dagger x]_{1,2,3,4} x^\dagger x]_5 [x^\dagger x]_{1,2,3,4} x^\dagger x \\ &= [[x^\dagger x]_{1,4} x^\dagger x] [x^\dagger x]_{1,4} x^\dagger x. \end{aligned}$$

In the final steps we noted that  $x^\dagger x \in \mathcal{G}^0 \oplus \mathcal{G}^1 \oplus \mathcal{G}^4 \oplus \mathcal{G}^5$  and that  $\mathbb{C} \subseteq \mathcal{G}_4^{\mathbb{C}}$  corresponds to  $\mathcal{G}^0 \oplus \mathcal{G}^5 \subseteq \mathcal{G}_5$ . Furthermore, since  $N_4^{\mathbb{C}}(xy) = N_4^{\mathbb{C}}(x) N_4^{\mathbb{C}}(y)$ , we have

$$\begin{aligned} N_5(xy) &= \overline{N_4^{\mathbb{C}}(F(x)F(y))} N_4^{\mathbb{C}}(F(x)F(y)) \\ &= \overline{N_4^{\mathbb{C}}(F(x))} \overline{N_4^{\mathbb{C}}(F(y))} N_4^{\mathbb{C}}(F(x)) N_4^{\mathbb{C}}(F(y)) \\ &= N_5(x) N_5(y) \end{aligned}$$

for all  $x, y \in \mathcal{G}$ . The invertible elements of the Dirac algebra are then as before

$$\mathcal{G}^\times(\mathbb{R}^{4,1}) = \{x \in \mathcal{G} : N_5(x) \neq 0\}$$

and the inverse of  $x \in \mathcal{G}^\times$  is

$$x^{-1} = \frac{1}{N_5(x)} [[x^\dagger x]_{1,4} x^\dagger x] [x^\dagger x]_{1,4} x^\dagger.$$

The above strategy could also have been used to obtain the expected result for  $N_3$  on  $\mathcal{G}(\mathbb{R}^{3,0}) \cong \mathcal{G}(\mathbb{C}^2)$  (with a corresponding isomorphism  $F$ ):

$$N_3(x) := \overline{N_2^{\mathbb{C}}(F(x))} N_2^{\mathbb{C}}(F(x)) = [x^\square x] x^\square x.$$

## 6.7 \*Norm functions and factorization identities

The norm functions

$$\begin{aligned}
N_0(x) &:= x, \\
N_1(x) &= x^\square x, \\
N_2(x) &= x^\square x, \\
N_3(x) &= [x^\square x]x^\square x, \\
N_4(x) &= [x^\square x]x^\square x, \\
N_5(x) &= [[x^\dagger x]_{1,4}x^\dagger x][x^\dagger x]_{1,4}x^\dagger x
\end{aligned}$$

constructed above (where we added  $N_0$  for completeness) all satisfy

$$N_k : \mathcal{G}(V) \rightarrow \mathbb{F},$$

where  $\dim V = k$ , and the multiplicative property

$$N_k(xy) = N_k(x)N_k(y) \quad (6.28)$$

for all  $x, y \in \mathcal{G}(V)$ . Furthermore, because these functions only involve products and grade involutions, and the proofs of the above identities only rely on commutation properties in the respective algebras, they even hold for arbitrary Clifford algebras  $Cl(X, R, r)$  over rings  $R$  with  $|X| = k = 0, 1, \dots, 5$ , respectively:

$$N_k : Cl(X, R, r) \rightarrow R.$$

The corresponding groups of invertible elements are thus

$$Cl^\times(X, R, r) = \{x \in Cl : N_k(x) \in R \text{ invertible}\}.$$

For matrix algebras, a similar multiplicative property is satisfied by the determinant. On the other hand, we have the following theorem concerning real matrices:

**Theorem 6.25.** *Assume that  $d : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$  is continuous and satisfies*

$$d(AB) = d(A)d(B) \quad (6.29)$$

for all  $A, B \in \mathbb{R}^{n \times n}$ . Then  $d$  must be either 0, 1,  $|\det|^\alpha$  or  $(\text{sign} \circ \det)|\det|^\alpha$  for some  $\alpha > 0$ .

In other words, we must have that  $d = d_1 \circ \det$ , where  $d_1 : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and satisfies  $d_1(\lambda\mu) = d_1(\lambda)d_1(\mu) \forall \mu, \nu \in \mathbb{R}$ . This function  $d_1$  is uniquely determined e.g. by whether  $d$  takes negative values, together with the value of  $d(\lambda \text{id})$  for any  $\lambda > 1$ . This means that the determinant is the *unique* real-valued function on real matrices with the product property (6.29). A proof of this theorem can be found in the appendix (Theorem A.6).

Now, looking at Table 5.1, we see that  $\mathcal{G}(\mathbb{R}^{k,k}) \cong \mathbb{R}^{2^k \times 2^k}$  for  $k = 0, 1, 2, \dots$ . From the above theorem we then know that there are *unique*<sup>20</sup> continuous functions  $N_{2k} : \mathcal{G}(\mathbb{R}^{k,k}) \rightarrow \mathbb{R}$  such that  $N_{2k}(xy) = N_{2k}(x)N_{2k}(y)$  and  $N_{2k}(\lambda) = \lambda^{2^k}$ . These are given by the usual determinant on the corresponding matrix algebra, and the corresponding multiplicative group  $\mathcal{G}^\times$  is the general linear group  $GL(2^k, \mathbb{R})$ .

<sup>20</sup>Actually, the functions are either  $\det$  or  $|\det|$ . The functions  $N_0, N_2$  and  $N_4$  constructed previously are smooth, however, so they must be equal to  $\det$ .



**Example 6.3.** The multiplicative property (6.28) of the norm functions leads to interesting factorization identities on rings. An example is  $N_1$  on  $\mathcal{G}(\mathbb{R}^{0,1}) \cong \mathbb{C}$ ,

$$(x_1^2 + x_2^2)(y_1^2 + y_2^2) = (x_1y_1 - x_2y_2)^2 + (x_1y_2 + x_2y_1)^2, \quad (6.30)$$

as well as the generalization given by  $N_2$  for anti-euclidean signatures (i.e. quaternions),

$$\begin{aligned} & (x_1^2 + x_2^2 + x_3^2 + x_4^2)(y_1^2 + y_2^2 + y_3^2 + y_4^2) \\ &= (x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4)^2 + (x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3)^2 \\ &+ (x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2)^2 + (x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1)^2, \end{aligned} \quad (6.31)$$

which hold for all  $x_j, y_k \in R$  and are called the *Gauss identity* and the *Lagrange identity*, respectively. These types of identities can be used to prove theorems in number theory. Using (6.31), one can for example prove that every positive integer can be written as a sum of four squares of integers. Or, in other words, that every positive integer is the norm (squared) of an integral quaternion. See e.g. [19] for a proof.

As far as we know, the general formulation and properties of the above norm functions were first given in [30] and independently discovered in [9, 40] (see also [32] for dimension  $k \leq 3$ ).

---

**Exercise 6.17.** Find explicit expressions for the norm functions  $N_4$  on  $\mathcal{G}(\mathbb{R}^4)$ , respectively  $N_5$  on  $\mathcal{G}(\mathbb{R}^5)$ , acting on a generic element  $x \in \mathcal{G}$ .

## 7 Euclidean and conformal geometry

### 7.1 Linearization of the Euclidean group

The *euclidean group*  $\mathbb{E}_n^+$  on  $\mathbb{R}^n$  consists of all orientation-preserving isometries, i.e. maps  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $|f(x) - f(y)| = |x - y|$  for all  $x, y \in \mathbb{R}^n$ . An element of  $\mathbb{E}_n^+$  can be shown to be a composition of rotations (orthogonal maps with determinant 1) and translations. As we have seen, rotations  $\underline{R} \in \text{SO}(n)$  can be represented by rotors  $R \in \text{Spin}(n)$  through

$$\mathbb{R}^n \ni x \mapsto \underline{R}(x) = R^* x R^{-1} = R x R^\dagger \in \mathbb{R}^n,$$

while translations are of the form

$$\mathbb{R}^n \ni x \mapsto T_a(x) = x + a \in \mathbb{R}^n, \quad a \in \mathbb{R}^n.$$

The euclidean group is sometimes slightly cumbersome to work with due to the fact that it not composed exclusively of linear transformations. We will therefore embed  $\mathbb{R}^n$  in a different geometric algebra where the euclidean isometries are transformed into linear maps.

Let  $\{e_1, \dots, e_n\}$  be an orthonormal basis of  $\mathbb{R}^n$  and introduce a new symbol  $e_0$  such that  $\{e_1, \dots, e_n, e_0\}$  is an orthonormal basis of  $\mathbb{R}^{n,0,1}$ , where  $e_0^2 = 0$ . Define a map

$$\begin{aligned} \mathbb{R}^n &\xrightarrow{\rho} \mathcal{G}(\mathbb{R}^{n,0,1}) \\ x &\mapsto 1 + e_0 x \end{aligned}$$

and let  $W$  denote the image of the map  $\rho$ . We extend a rotation  $\underline{R}$  to  $W$  as

$$\underline{R}(1 + e_0 x) = R(1 + e_0 x)R^\dagger = R R^\dagger + R e_0 x R^\dagger = 1 + e_0 R x R^\dagger = 1 + e_0 \underline{R}(x),$$

and hence have  $\underline{R} : W \rightarrow W$  and  $\underline{R} \circ \rho = \rho \circ \underline{R}$  on  $\mathbb{R}^n$ .

Let us now consider the translation map  $T_a(x) = x + a$ . For this we first introduce

$$A = e^{-\frac{1}{2}e_0 a} = 1 - \frac{1}{2}e_0 a,$$

and a slightly modified version of the grade involution  $x \rightarrow x^*$  by demanding that  $e_0^* = e_0$ , while as usual  $e_i^* = -e_i$ , for  $i = 1, \dots, n$ . We then obtain

$$A^* = 1 + \frac{1}{2}e_0 a \quad \text{and} \quad A^\dagger = 1 + \frac{1}{2}e_0 a = A^{-1}.$$

Hence,

$$\begin{aligned} \underline{A}(\rho(x)) &= A^*(1 + e_0 x)A^{-1} = (1 + \frac{1}{2}e_0 a)(1 + e_0 x)(1 + \frac{1}{2}e_0 a) \\ &= (1 + e_0 x + \frac{1}{2}e_0 a)(1 + \frac{1}{2}e_0 a) = 1 + \frac{1}{2}e_0 a + e_0 x + \frac{1}{2}e_0 a \\ &= 1 + e_0(x + a) = \rho(x + a) = \rho(T_a(x)), \end{aligned}$$

i.e.  $\underline{A} \circ \rho = \rho \circ T_a$  on  $\mathbb{R}^n$ .

This means that any isometry  $S = T_a \circ \underline{R} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  has the rotor representation  $\underline{A} \underline{R} = \underline{A} \circ \underline{R} : W \rightarrow W$ , with  $A \underline{R} = e^{-\frac{1}{2}e_0 a} e^{-\frac{1}{2}B} \in \mathcal{G}^\times(\mathbb{R}^{n,0,1})$ . Also note that an arbitrary point  $x = T_x(0) \in \mathbb{R}^n$  has the representation

$$\rho(x) = \underline{e^{-\frac{1}{2}e_0 x}}(\rho(0)) = (e^{-\frac{1}{2}e_0 x})^* \mathbf{1} (e^{-\frac{1}{2}e_0 x})^\dagger = e^{e_0 x}$$

in  $W \subseteq \mathcal{G}^\times(\mathbb{R}^{n,0,1})$ .

---

**Exercise 7.1.** Repeat the above construction, but replacing the null vector  $e$  by a null 2-blade in  $\mathbb{R}^{2,1}$ ,  $\mathbb{R}^{1,2}$ , or  $\mathbb{R}^{0,2}$ . Explain what simplifications can be made and show that the construction extends to the full euclidean group  $\mathbb{E}_n$  (not necessarily orientation-preserving).

## 7.2 Conformal algebra

For now, we refer to Chapter 10 in [12]. For an application of the conformal algebra to so-called *twistors*, see [3].

## 8 Representation theory

In this section we will use the classification of geometric algebras as matrix algebras, which was developed in Section 5, to work out the representation theory of these algebras. Since one can find representations of geometric algebras in many areas of mathematics and physics, this leads to a number of interesting applications. In this section we will consider two main examples in detail, namely normed division algebras and vector fields on higher-dimensional spheres. Another important application of the representation theory for geometric algebras is to *spinor spaces*, which will be treated in Section 9.

### 8.1 Real, complex and quaternionic representations

**Definition 8.1.** For  $\mathbb{K} = \mathbb{R}, \mathbb{C}$  or  $\mathbb{H}$ , we define a  $\mathbb{K}$ -*representation* of  $\mathcal{G}(V, q)$  as an  $\mathbb{R}$ -algebra homomorphism

$$\rho: \mathcal{G}(V, q) \rightarrow \text{End}_{\mathbb{K}}(W),$$

where  $W$  is a finite-dimensional vector space over  $\mathbb{K}$ .  $W$  is called a  $\mathcal{G}(V, q)$ -*module* over  $\mathbb{K}$ .

Note that a vector space over  $\mathbb{C}$  respectively<sup>21</sup>  $\mathbb{H}$  can be considered as a real vector space together with linear operators  $J$  respectively  $I, J, K$  in  $\text{End}_{\mathbb{R}}(W)$  that anticommute and square to minus the identity. In the definition above we assume that these operators commute with  $\rho(x)$  for all  $x \in \mathcal{G}$ , so that  $\rho$  can be said to respect the  $\mathbb{K}$ -structure of the space  $W$ . When talking about the dimension of the module  $W$  we will always refer to its dimension as a real vector space.

The standard strategy when studying representation theory is to look for irreducible representations.

**Definition 8.2.** A representation  $\rho$  is called *reducible* if  $W$  can be written as a direct sum of proper (not equal to 0 or  $W$ ) invariant subspaces, i.e.

$$W = W_1 \oplus W_2 \quad \text{and} \quad \rho(x)(W_j) \subseteq W_j \quad \forall x \in \mathcal{G}.$$

In this case we can write  $\rho = \rho_1 \oplus \rho_2$ , where  $\rho_j(x) := \rho(x)|_{W_j}$ . A representation is called *irreducible* (or simply an *irrep*) if it is not reducible.

*Remark.* The traditional definition of an irreducible representation is that it does not have any proper invariant subspaces (the above notion is then called *indecomposable*). However, because  $\mathcal{G}$  is the group algebra of a finite group (the Clifford group) one can verify that these two definitions are equivalent in this case (see Exercise 8.1).

**Proposition 8.1.** *Every  $\mathbb{K}$ -representation  $\rho$  of a geometric algebra  $\mathcal{G}(V, q)$  can be split up into a direct sum  $\rho = \rho_1 \oplus \dots \oplus \rho_m$  of irreducible representations.*

*Proof.* This follows directly from the definitions and the fact that  $W$  is finite-dimensional. □

<sup>21</sup>Also note that  $\mathbb{H}$  is not a field but a skew field.

**Definition 8.3.** Two  $\mathbb{K}$ -representations  $\rho_j : \mathcal{G}(V, q) \rightarrow \text{End}_{\mathbb{K}}(W_j)$ ,  $j = 1, 2$ , are said to be *equivalent* if there exists a  $\mathbb{K}$ -linear isomorphism  $F : W_1 \rightarrow W_2$  such that

$$F \circ \rho_1(x) \circ F^{-1} = \rho_2(x) \quad \forall x \in \mathcal{G}.$$

**Theorem 8.2.** *Up to equivalence, the only irreducible representations of the matrix algebras  $\mathbb{K}^{n \times n}$  and  $\mathbb{K}^{n \times n} \oplus \mathbb{K}^{n \times n}$  are*

$$\rho : \mathbb{K}^{n \times n} \rightarrow \text{End}_{\mathbb{K}}(\mathbb{K}^n)$$

and

$$\rho_{1,2} : \mathbb{K}^{n \times n} \oplus \mathbb{K}^{n \times n} \rightarrow \text{End}_{\mathbb{K}}(\mathbb{K}^n)$$

respectively, where  $\rho$  is the defining representation (ordinary matrix multiplication) and

$$\begin{aligned} \rho_1(x, y) &:= \rho(x), \\ \rho_2(x, y) &:= \rho(y). \end{aligned}$$

This follows from the classical fact that the algebras  $\mathbb{K}^{n \times n}$  are *simple* (i.e. have no proper two-sided ideals) and that simple algebras have only one irreducible representation up to equivalence. See e.g. [25] for details.

**Theorem 8.3** (Real representations). *From the above, together with the classification of real geometric algebras, follows the table of representations in Table 8.1, where  $\nu_{s,t}$  is the number of inequivalent irreducible representations and  $d_{s,t}$  is the (real) dimension of an irreducible representation for  $\mathcal{G}(\mathbb{R}^{s,t})$ . The cases for  $s, t > 8$  are obtained using the periodicity*

$$\begin{aligned} \nu_{s+8k,t} &= \nu_{s,t+8k} = \nu_{s,t}, \\ d_{s+8k,t} &= d_{s,t+8k} = 16^k d_{s,t}. \end{aligned} \tag{8.1}$$

$n$	$\mathcal{G}(\mathbb{R}^{n,0})$	$\nu_{n,0}$	$d_{n,0}$	$\mathcal{G}(\mathbb{R}^{0,n})$	$\nu_{0,n}$	$d_{0,n}$
0	$\mathbb{R}$	1	1	$\mathbb{R}$	1	1
1	$\mathbb{R} \oplus \mathbb{R}$	2	1	$\mathbb{C}$	1	2
2	$\mathbb{R}^{2 \times 2}$	1	2	$\mathbb{H}$	1	4
3	$\mathbb{C}^{2 \times 2}$	1	4	$\mathbb{H} \oplus \mathbb{H}$	2	4
4	$\mathbb{H}^{2 \times 2}$	1	8	$\mathbb{H}^{2 \times 2}$	1	8
5	$\mathbb{H}^{2 \times 2} \oplus \mathbb{H}^{2 \times 2}$	2	8	$\mathbb{C}^{4 \times 4}$	1	8
6	$\mathbb{H}^{4 \times 4}$	1	16	$\mathbb{R}^{8 \times 8}$	1	8
7	$\mathbb{C}^{8 \times 8}$	1	16	$\mathbb{R}^{8 \times 8} \oplus \mathbb{R}^{8 \times 8}$	2	8
8	$\mathbb{R}^{16 \times 16}$	1	16	$\mathbb{R}^{16 \times 16}$	1	16

Table 8.1: Number and dimension of irreducible representations of euclidean and anti-euclidean geometric algebras. The case of general signatures can be obtained analogously from Table 5.1.

Note that the cases when there are two inequivalent irreducible representations of  $\mathcal{G}(\mathbb{R}^{s,t})$  correspond exactly to the cases when the pseudoscalar  $I$  is central and squares to one. Furthermore, the two representations  $\rho_{\pm}$  are characterized by their value on the pseudoscalar,

$$\rho_{\pm}(I) = \pm \text{id}.$$

It is clear that these two representations are inequivalent, since if  $F: W_1 \rightarrow W_2$  is an isomorphism, and  $\rho(I) = \sigma \text{id}_{W_1}$ , then also  $F \circ \rho(I) \circ F^{-1} = \sigma \text{id}_{W_2}$ .

As one could expect, the corresponding representation theory over the complex field (where a representation is defined analogously as a  $\mathbb{C}$ -algebra homomorphism  $\rho: \mathcal{G}(\mathbb{C}^n) \rightarrow \text{End}_{\mathbb{C}} W$ ) is simpler:

**Theorem 8.4** (Complex representations). *Let  $\nu_n^{\mathbb{C}}$  denote the number of inequivalent irreducible complex representations of  $\mathcal{G}(\mathbb{C}^n)$ , and let  $d_n^{\mathbb{C}}$  denote the complex dimension of such a representation. Then, if  $n = 2k$  is even,*

$$\nu_n^{\mathbb{C}} = 1 \quad \text{and} \quad d_n^{\mathbb{C}} = 2^k,$$

while if  $n = 2k + 1$  is odd,

$$\nu_n^{\mathbb{C}} = 2 \quad \text{and} \quad d_n^{\mathbb{C}} = 2^k.$$

The inequivalent representations in the odd case are characterized by their value (sign) on the *complex volume element*  $I_{\mathbb{C}} := i^k I$ , which in that case is central, and always squares to one.

We will now consider the situation when the representation space  $W$  is endowed with an inner product. Note that if  $W$  is a real vector space over  $\mathbb{K}$  with an ( $\mathbb{R}$ -valued) inner product, then we can always find a  $\mathbb{K}$ -invariant (still  $\mathbb{R}$ -valued) inner product on  $W$ , i.e. such that the operators  $J$  or  $I, J, K$  are orthogonal. Namely, let  $\langle \cdot, \cdot \rangle_{\mathbb{R}}$  be an inner product on  $W$  and put

$$\langle x, y \rangle_{\mathbb{C}} := \sum_{\Gamma \in \{\text{id}, J\}} \langle \Gamma x, \Gamma y \rangle_{\mathbb{R}}, \quad \langle x, y \rangle_{\mathbb{H}} := \sum_{\Gamma \in \{\text{id}, I, J, K\}} \langle \Gamma x, \Gamma y \rangle_{\mathbb{R}}. \quad (8.2)$$

Then  $\langle Jx, Jy \rangle_{\mathbb{K}} = \langle x, y \rangle_{\mathbb{K}}$  and  $\langle Jx, y \rangle_{\mathbb{K}} = -\langle x, Jy \rangle_{\mathbb{K}}$ , etc.

In the same way, when  $V$  is euclidean or anti-euclidean, we can for a given representation  $\rho: \mathcal{G}(V) \rightarrow \text{End}_{\mathbb{K}}(W)$  find an inner product such that  $\rho$  acts orthogonally with unit vectors, i.e. such that  $\langle \rho(e)x, \rho(e)y \rangle = \langle x, y \rangle$  for all  $x, y \in W$  and  $e \in V$  with  $e^2 = \pm 1$ . We construct such an inner product by averaging a, possibly  $\mathbb{K}$ -invariant, inner product  $\langle \cdot, \cdot \rangle_{\mathbb{K}}$  on  $W$  over the Clifford group:

For general signatures, take an orthonormal basis  $E$  of  $V$  and put

$$\langle x, y \rangle := \sum_{\Gamma \in \mathcal{B}_E} \langle \rho(\Gamma)x, \rho(\Gamma)y \rangle_{\mathbb{K}}. \quad (8.3)$$

We then have that

$$\langle \rho(e_i)x, \rho(e_i)y \rangle = \langle x, y \rangle \quad (8.4)$$

for all  $e_i \in E$ , and if  $e_i \neq e_j$  in  $E$  have the same signature,

$$\begin{aligned} \langle \rho(e_i)x, \rho(e_j)y \rangle &= \langle \rho(e_i)\rho(e_i)x, \rho(e_i)\rho(e_j)y \rangle = \pm \langle x, \rho(e_i)\rho(e_j)y \rangle \\ &= \mp \langle x, \rho(e_j)\rho(e_i)y \rangle = \mp \langle \rho(e_j)x, \rho(e_j)\rho(e_i)y \rangle \\ &= -\langle \rho(e_j)x, \rho(e_i)y \rangle. \end{aligned}$$

Thus, in the (anti-)euclidean case, if  $e = \sum_i a_i e_i$  and  $\sum_i a_i^2 = 1$ , we obtain

$$\langle \rho(e)x, \rho(e)y \rangle = \sum_{i,j} a_i a_j \langle \rho(e_i)x, \rho(e_i)y \rangle = \langle x, y \rangle.$$

Hence, this inner product has the desired property. Also note that, for  $v \in V = \mathbb{R}^{n,0}$ , we have

$$\langle \rho(v)x, y \rangle = \langle x, \rho(v)y \rangle, \quad (8.5)$$

while for  $V = \mathbb{R}^{0,n}$ ,

$$\langle \rho(v)x, y \rangle = -\langle x, \rho(v)y \rangle, \quad (8.6)$$

i.e.  $\rho(v)$  is symmetric for euclidean spaces and antisymmetric for anti-euclidean spaces.

---

**Exercise 8.1.** Prove that a representation  $\rho$  of  $\mathcal{G}$  is irreducible (according to Definition 8.2) if and only if it does not have any proper invariant subspaces.

*Hint:* You may w.l.o.g. assume that the representation space has an invariant inner product.

## 8.2 Fermionic representations

We note that the real dimension  $d_{s,t}$  of an irreducible representation space always turns out to be a pure power of 2, and that the same holds concerning the complex dimension  $d_n^{\mathbb{C}}$ . Coincidentally, this is also the case for the dimension of the algebra  $\mathcal{G}$  itself, and for the dimension of the fermionic Fock space introduced in Section 4.3. These facts can be used to construct explicit irreducible representations of Clifford algebras of a form which often appears in the physics literature, and which in some cases can be regarded as the canonical choice.

The arguably most canonical or straightforward representation is the one where  $\mathcal{G}(V)$  just acts on itself by left multiplication (see Definition 6.2),

$$\rho = L: \mathcal{G} \rightarrow \text{End}(\mathcal{G}),$$

or, given an ON basis  $\{e_j\}_j$  of  $V$  and using the fermionic operators (4.2),

$$\rho(e_j) = L_{e_j} = e_j(\cdot) = e_j \lrcorner (\cdot) + e_j \wedge (\cdot) = c_j + c_j^\dagger, \quad j = 1, \dots, d = \dim V.$$

However, this representation is certainly reducible. Indeed, the dimension of  $\mathcal{G}(V)$  is  $2^d$  while according to Table 8.1 the real dimension of an irrep typically grows like  $2^{\lfloor d/2 \rfloor}$  when it is purely real, and  $2 \cdot 2^{\lfloor d/2 \rfloor}$  when it is complex or quaternionic. However, an agreement in dimensions can be reached if we can by some means identify pairs of basis elements in the representation, i.e. reduce to approximately half of the dimensions in  $V$ , say by fixing  $m = \lfloor d/2 \rfloor$  orthogonal planes in  $V$ , and by complexification.

Note first that in the case of fully mixed signature,  $V = \mathbb{R}^{m,m}$ ,  $d = 2m$ , there is a more or less<sup>22</sup> canonical split of the space into an euclidean and an anti-euclidean part, and that we may, given an orthonormal basis  $\{e_j, \epsilon_j\}_{j=1}^m$  of  $V$  with  $e_j^2 = 1$ ,  $\epsilon_j^2 = -1$ , define a basis of null-vectors (cf. Example 3.2)

$$n_j^\pm := \frac{1}{2}(e_j \pm \epsilon_j), \quad j = 1, \dots, m,$$

satisfying the anticommutation relations

$$\{n_j^\pm, n_k^\pm\} = 2n_j^\pm * n_k^\pm = 0, \quad \{n_j^+, n_k^-\} = 2n_j^+ * n_k^- = \delta_{jk}.$$

---

<sup>22</sup>it is still basis dependent

Comparing to the fermion relations (4.3), these are hence naturally represented on the (real or complexified) Fock space  $\mathcal{F} = \mathcal{G}(\mathbb{R}^m) = \mathcal{Cl}(\{e_j\}_{j=1}^m, \mathbb{R}, 1)$  via e.g.

$$\rho(n_j^+) := c_j^\dagger, \quad \rho(n_j^-) := c_j,$$

and the obtained representation  $\rho: \mathcal{G}(\mathbb{R}^{m,m}) \rightarrow \text{End } \mathcal{F}$  is found to be irreducible simply for dimensional reasons. We have here taken the  $m$  orthogonal timelike blades  $e_j \epsilon_j = 2n_j^- \wedge n_j^+$  to represent each fermionic dimension or ‘fermion’  $c_j^{(\dagger)}$ . Conversely, given  $m$  operators  $c_j, c_j^\dagger$  satisfying the anticommutation relations (4.3), we can form

$$\hat{e}_j := c_j^\dagger + c_j, \quad \hat{\epsilon}_j := c_j^\dagger - c_j,$$

with  $\hat{e}_j^2 = 1, \hat{\epsilon}_j^2 = -1$  etc., and therefore generate a representation of  $\mathcal{G}(\mathbb{R}^{m,m})$ .

In the complex case  $V = \mathbb{C}^n$  with  $n = 2m$  one may also make a (less canonical) division of the space into  $m$  orthogonal planes, exactly as above by taking  $\epsilon_j := ie_{m+j}$ , thereby producing a  $2^m$ -dimensional complex Fock space  $\mathcal{F} = \mathcal{G}(\mathbb{C}^m)$ . For odd  $n = 2m + 1$ , the requirement that  $\rho(e_1 \dots e_n) = (-i)^m \rho(I_{\mathbb{C}}) = \pm(-i)^m \text{id}_{\mathcal{F}}$  can be used to deduce the action of

$$\rho(e_n) = \pm(-i)^m \rho(e_1) \dots \rho(e_{2m}) = \pm \rho(e_1) \dots \rho(e_m) \rho(\epsilon_1) \dots \rho(\epsilon_m).$$

In the purely real euclidean case  $V = \mathbb{R}^{2m}$ , one may by means of complexification reduce to the previous case, with a non-canonical split and defining

$$n_j^\pm := \frac{1}{2}(e_j \pm ie_{m+j}), \quad \rho(n_j^+) := c_j^\dagger, \quad \rho(n_j^-) := c_j, \quad j = 1, \dots, m,$$

again with  $\mathcal{G}(\mathbb{C}^{2m})$  then acting irreducibly on a complex Fock space  $\mathcal{F} = \mathcal{G}(\mathbb{C}^m)$ .

However, sometimes the considered space comes equipped with a canonical pairing. For example, in the even-dimensional real case it could be that we also have a complex structure, i.e. a linear operator  $J \in \text{O}(V)$  such that  $J^2 = -\text{id}_V$ , as discussed above. Then, by selecting an ON basis for  $V$  of the form  $\{e_j, Je_j\}_{j=1}^m$ , we can form *canonical* complexified fermionic operators

$$n_j^\pm := \frac{1}{2}(e_j \pm iJe_j), \quad \rho(n_j^+) := c_j^\dagger, \quad \rho(n_j^-) := c_j, \quad j = 1, \dots, m.$$

A particular case one may keep in mind here is that of the abstract representation space or  $\mathcal{G}$ -module itself in the case that the irrep  $W \cong \mathbb{R}^{2m=d_{s,t}}$  of  $\mathcal{G}(V = \mathbb{R}^{s,t})$  turns out to be complex or quaternionic, i.e.

$$J^2 = -\text{id}_W, \quad \rho(e_j)J = J\rho(e_j) \quad \forall j = 1, \dots, s+t,$$

for some  $J \in \text{O}(W)$ . This complex structure may then be used to produce a canonical irreducible complex fermionic Fock space representation of the Clifford algebra  $\mathcal{G}(W)$  as above with  $\mathcal{F} = \mathcal{G}(\mathbb{C}^{d_{s,t}/2})$ . Examples of such two-level representations

$$\mathcal{G}(V) \rightarrow \text{End } W, \quad \mathcal{G}(W) \rightarrow \text{End } \mathcal{F}$$

appear abundantly within high-energy and particle physics by means of the framework of *second quantization*. See [31, Section 3.5] for a discussion of canonical pairings for representations in the context of a particular class of quantum mechanical models called *supersymmetric matrix models*. The above remarks also apply to the spinor representations which will be discussed in Section 9.



### 8.3 Examples

We are now ready for some examples which illustrate how representations of geometric algebras can appear naturally in various contexts in mathematics and how their representation theory can be used to prove important theorems.

#### 8.3.1 Normed division algebras

Our first example concerns the possible dimensions of normed division algebras. A *normed division algebra* is an algebra  $\mathcal{A}$  over  $\mathbb{R}$  (not necessarily associative) with unit and a norm  $|\cdot|$  such that

$$|xy| = |x||y| \tag{8.7}$$

for all  $x, y \in \mathcal{A}$  and such that every nonzero element is invertible. We will prove the following

**Theorem 8.5** (Hurwitz' Theorem). *If  $\mathcal{A}$  is a finite-dimensional normed division algebra over  $\mathbb{R}$ , then its dimension is either 1, 2, 4 or 8.*

*Remark.* This corresponds uniquely to  $\mathbb{R}$ ,  $\mathbb{C}$ ,  $\mathbb{H}$ , and the *octonions*  $\mathbb{O}$ , respectively. The proof of uniqueness requires some additional steps, see e.g. [5].

Let us first consider what restrictions that the requirement (8.7) puts on the norm. Assume that  $\mathcal{A}$  has dimension  $n$ . For every  $a \in \mathcal{A}$  we have a linear transformation

$$\begin{aligned} L_a: \mathcal{A} &\rightarrow \mathcal{A}, \\ x &\mapsto ax \end{aligned}$$

given by left multiplication by  $a$ . When  $|a| = 1$  we then have

$$|L_a x| = |ax| = |a||x| = |x|, \tag{8.8}$$

i.e.  $L_a$  preserves the norm. Hence, it maps the unit sphere  $S := \{x \in \mathcal{A} : |x| = 1\}$  in  $\mathcal{A}$  into itself. Furthermore, since every element in  $\mathcal{A}$  is invertible, we can for each pair  $x, y \in S$  find an  $a \in S$  such that  $L_a x = ax = y$ . Now, these facts imply a large amount of symmetry of  $S$ . In fact, we have the following

**Lemma 8.6.** *Assume that  $V$  is a finite-dimensional normed vector space. Let  $S_V$  denote the unit sphere in  $V$ . If, for every  $x, y \in S_V$ , there exists an operator  $L \in \text{End}(V)$  such that  $L(S_V) \subseteq S_V$  and  $L(x) = y$ , then  $V$  must be an inner product space.*

*\*Proof.* We will need the following fact: Every compact subgroup  $G$  of  $\text{GL}(n)$  preserves some inner product on  $\mathbb{R}^n$ . This can be shown by picking a Haar-measure  $\mu$  on  $G$  and averaging any inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{R}^n$  over  $G$  using this measure, (analogous to (8.2),(8.3) where the group is finite)

$$\langle x, y \rangle_G := \int_G \langle gx, gy \rangle d\mu(g). \tag{8.9}$$

Now, let  $G$  be the group of linear transformations on  $V \cong \mathbb{R}^n$  which preserve its norm  $|\cdot|$ .  $G$  is compact in the finite-dimensional operator norm topology, since  $G = \bigcap_{x \in V} \{L \in \text{End}(\mathbb{R}^n) : |Lx| = |x|\}$  is closed and bounded by 1.

Furthermore,  $L \in G$  is injective and therefore an isomorphism. The group structure is obvious. Hence,  $G$  is a compact subgroup of  $\mathrm{GL}(n)$ .

From the above we know that there exists an inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{R}^n$  which is preserved by  $G$ . Let  $|\cdot|_\circ$  denote the norm associated to this inner product, i.e.  $|x|_\circ^2 = \langle x, x \rangle$ . Take a point  $x \in \mathbb{R}^n$  with  $|x| = 1$  and rescale the inner product so that also  $|x|_\circ = 1$ . Let  $S$  and  $S_\circ$  denote the unit spheres associated to  $|\cdot|$  and  $|\cdot|_\circ$ , respectively. By the conditions in the lemma, there is for every  $y \in S$  an  $L \in G$  such that  $L(x) = y$ . But  $G$  also preserves the norm  $|\cdot|_\circ$ , so  $y$  must also lie in  $S_\circ$ . Hence,  $S$  is a subset of  $S_\circ$ . However, being unit spheres associated to norms,  $S$  and  $S_\circ$  are both homeomorphic to the standard sphere  $S^{n-1}$ , so we must have that they are equal. Therefore, the norms must be equal.  $\square$

We now know that our normed division algebra  $\mathcal{A}$  has some inner product  $\langle \cdot, \cdot \rangle$  such that  $\langle x, x \rangle = |x|^2$ . We call an element  $a \in \mathcal{A}$  *imaginary* if  $a$  is orthogonal to the unit element, i.e. if  $\langle a, 1_{\mathcal{A}} \rangle = 0$ . Let  $\mathrm{Im} \mathcal{A}$  denote the  $(n-1)$ -dimensional subspace of imaginary elements. We will observe that  $\mathrm{Im} \mathcal{A}$  acts on  $\mathcal{A}$  in a special way.

Take a curve  $\gamma: (-\epsilon, \epsilon) \rightarrow S$  on the unit sphere such that  $\gamma(0) = 1_{\mathcal{A}}$  and  $\gamma'(0) = a \in \mathrm{Im} \mathcal{A}$ . (Note that  $\mathrm{Im} \mathcal{A}$  is the tangent space to  $S$  at the unit element.) Then, because the product in  $\mathcal{A}$  is continuous,  $\forall x, y \in \mathcal{A}$

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} L_{\gamma(t)} x &= \lim_{h \rightarrow 0} \frac{1}{h} (L_{\gamma(h)} x - L_{\gamma(0)} x) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} (\gamma(h) - \gamma(0)) x = \gamma'(0) x = a x = L_a x \end{aligned}$$

and

$$\begin{aligned} 0 &= \frac{d}{dt} \Big|_{t=0} \langle x, y \rangle = \frac{d}{dt} \Big|_{t=0} \langle L_{\gamma(t)} x, L_{\gamma(t)} y \rangle \\ &= \langle \frac{d}{dt} \Big|_{t=0} L_{\gamma(t)} x, L_{\gamma(0)} y \rangle + \langle L_{\gamma(0)} x, \frac{d}{dt} \Big|_{t=0} L_{\gamma(t)} y \rangle \\ &= \langle L_a x, y \rangle + \langle x, L_a y \rangle. \end{aligned}$$

Hence,  $L_a^* = -L_a$  for  $a \in \mathrm{Im} \mathcal{A}$ . If, in addition,  $|a| = 1$  we have that  $L_a \in \mathrm{O}(\mathcal{A}, |\cdot|^2)$ , so  $L_a^2 = -L_a L_a^* = -\mathrm{id}$ . For an arbitrary imaginary element  $a$  we obtain by rescaling

$$L_a^2 = -|a|^2 \mathrm{id}. \quad (8.10)$$

This motivates us to consider the geometric algebra  $\mathcal{G}(\mathrm{Im} \mathcal{A}, q)$  with quadratic form  $q(a) := -|a|^2$ . By (8.10) and the universal property of geometric algebras (Theorem 2.1) we find that  $L$  extends to a representation of  $\mathcal{G}(\mathrm{Im} \mathcal{A}, q)$  on  $\mathcal{A}$ ,

$$\hat{L}: \mathcal{G}(\mathrm{Im} \mathcal{A}, q) \rightarrow \mathrm{End}(\mathcal{A}), \quad (8.11)$$

i.e. a representation of  $\mathcal{G}(\mathbb{R}^{0, n-1})$  on  $\mathbb{R}^n$ . The representation theory then demands that  $n$  is a multiple of  $d_{0, n-1}$ . By studying Table 8.1 and taking periodicity (8.1) into account we find that this is only possible for  $n = 1, 2, 4, 8$ .  $\square$

*Remark.* Not only  $\mathbb{R}$ ,  $\mathbb{C}$ , and  $\mathbb{H}$ , but also the octonion algebra  $\mathbb{O}$  can be constructed explicitly in terms of Clifford algebras. However, one cannot use simply the geometric product in this case since the octonion product is non-associative. With a choice of *octonionic structure*, e.g.

$$C := e_1 e_2 e_4 + e_2 e_3 e_5 + e_3 e_4 e_6 + e_4 e_5 e_7 + e_5 e_6 e_1 + e_6 e_7 e_2 + e_7 e_1 e_3 \in \mathcal{G}^3,$$

we can identify  $\mathbb{O} := \mathcal{G}^0 \oplus \mathcal{G}^1 \subseteq \mathcal{G}(\mathbb{R}^{0,7})$ , with a multiplication given by

$$a \diamond b := \langle ab(1 - C) \rangle_{0,1},$$

or alternatively,  $\mathbb{O} := \mathcal{G}^1(\mathbb{R}^8)$  with

$$a \diamond b := \langle ae_8b(1 - CI_7)(1 - I) \rangle_1,$$

where  $I_7 := e_1 \dots e_7$ , and  $e_8$  in this case represents the unit of  $\mathbb{O}$ . We refer to [28] for more on this construction, and to [5] for more on octonions in general.

### 8.3.2 Vector fields on spheres

In our next example we consider the  $N$ -dimensional unit spheres  $S^N$  and use representations of geometric algebras to construct vector fields on them. The number of such vector fields that can be found gives us information about the topological features of these spheres.

**Theorem 8.7** (Radon–Hurwitz). *On  $S^N$  there exist  $n_N$  pointwise linearly independent vector fields, where, if we write  $N$  uniquely as*

$$N + 1 = (2t + 1)2^{4a+b}, \quad t, a \in \mathbb{N}, \quad b \in \{0, 1, 2, 3\},$$

then

$$n_N = 8a + 2^b - 1.$$

For example,

$N$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$n_N$	0	1	0	3	0	1	0	7	0	1	0	3	0	1	0	8	0

**Corollary.**  $S^1$ ,  $S^3$  and  $S^7$  are parallelizable.

*Remark.* The number of vector fields constructed in this way is actually the maximum number of possible such fields on  $S^N$ . This is a much deeper result proven by Adams [2] using algebraic topology.

Our main observation is that if  $\mathbb{R}^{N+1}$  is a  $\mathcal{G}(\mathbb{R}^{0,n})$ -module then we can construct  $n$  pointwise linearly independent vector fields on the unit sphere

$$S^N = \{x \in \mathbb{R}^{N+1} : \langle x, x \rangle = 1\}.$$

Namely, suppose we have a representation  $\rho$  of  $\mathcal{G}(\mathbb{R}^{0,n})$  on  $\mathbb{R}^{N+1}$ . Take an inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{R}^{N+1}$  such that the action of  $\rho$  is orthogonal and pick any basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^{0,n}$ . We can now define a collection of smooth vector fields  $\{V_1, \dots, V_n\}$  on  $\mathbb{R}^{N+1}$  by

$$V_i(x) := \rho(e_i)x, \quad i = 1, \dots, n.$$

According to the observation (8.6) this action is antisymmetric, so that

$$\langle V_i(x), x \rangle = \langle \rho(e_i)x, x \rangle = -\langle x, \rho(e_i)x \rangle = 0.$$

Hence,  $V_i(x) \in T_x S^N$  for  $x \in S^N$ . By restricting to  $S^N$  we therefore have  $n$  tangent vector fields. It remains to show that these are pointwise linearly independent. Take  $x \in S^N$  and consider the linear map

$$\begin{aligned} i_x: \mathbb{R}^{0,n} &\rightarrow T_x S^N \\ v &\mapsto i_x(v) := \rho(v)x \end{aligned}$$

Since the image of  $i_x$  is  $\text{Span}_{\mathbb{R}}\{V_i(x)\}$  it is sufficient to prove that  $i_x$  is injective. But if  $i_x(v) = \rho(v)x = 0$  for some  $v \in \mathbb{R}^{0,n}$  then also  $v^2x = \rho(v)^2x = 0$ , so we must have  $v = 0$ .

Now, for a fixed  $N$  we want to find as many vector fields as possible, so we seek the highest  $n$  such that  $\mathbb{R}^{N+1}$  is a  $\mathcal{G}(\mathbb{R}^{0,n})$ -module. From the representation theory we know that this requires that  $N+1$  is a multiple of  $d_{0,n}$ . Furthermore, since  $d_{0,n}$  is a power of 2 we obtain the maximal such  $n$  when  $N+1 = p2^m$ , where  $p$  is odd and  $d_{0,n} = 2^m$ . Using Table 8.1 and the periodicity (8.1) we find that if we write  $N+1 = p2^{4a+b}$ , with  $0 \leq b \leq 3$ , then  $n = 8a + 2^b - 1$ . This proves the theorem.  $\square$

## 9 Spinors

This section is currently incomplete. We refer to the references below for more complete treatments.

In general, a *spinor*  $\Psi$  is an object upon which a rotor  $R$  acts by single-sided action like left multiplication  $\Psi \mapsto R\Psi$ , instead of two-sided action like e.g.  $\Psi \mapsto \underline{R}(\Psi) = R\Psi R^\dagger$ . More precisely, a *spinor space* is a representation space for the rotor group, but such that the action does not factor through the twisted adjoint action  $\widetilde{\text{Ad}}: \text{Spin}^+ \rightarrow \text{SO}^+$ . Due to historical reasons and varying applications, it is useful to have several different notions of spinor spaces available. The most commonly considered types of spinor spaces are abstract spinor modules and irreducible Spin representations, as well as ideal spinors, spinor operators and, more generally, mixed-action spinors.

### 9.1 Spin representations

Recall the representation theory of geometric algebras of Section 8. Given  $\mathcal{G}(\mathbb{R}^{s,t})$ , there is either one or two inequivalent irreducible  $\mathbb{K}$ -representations of real dimension  $d_{s,t}$ , where  $\mathbb{K}$  is either  $\mathbb{R}$ ,  $\mathbb{C}$ , or  $\mathbb{H}$ . This can be read off from Table 5.1 together with periodicity. We call such an irreducible representation space,

$$\mathcal{S}_{s,t} := \mathbb{K}^{d_{s,t}/\dim_{\mathbb{R}} \mathbb{K}} \cong \mathbb{R}^{d_{s,t}},$$

the corresponding *spinor module*<sup>23</sup> of  $\mathcal{G}(\mathbb{R}^{s,t})$ . Furthermore, because the Pin, Spin and rotor groups are contained in  $\mathcal{G}$ , this will also provide us with a canonical representation space for those groups.

**Proposition 9.1.** *The representation of the group  $\text{Pin}(s,t)$ , obtained by restricting an irreducible real representation  $\rho: \mathcal{G}(\mathbb{R}^{s,t}) \rightarrow \text{End}_{\mathbb{R}}(\mathcal{S}_{s,t})$  to  $\text{Pin} \subseteq \mathcal{G}$ , is irreducible. Furthermore, whenever there are two inequivalent such representations  $\rho$ , their corresponding restrictions to Pin are also inequivalent.*

*Proof.* The first statement follows from the fact that the standard basis of  $\mathcal{G}$  is contained in the Pin group. The second follows by recalling the role of the pseudoscalar  $I \in \text{Pin}$  in singling out inequivalent irreducible representations.  $\square$

**Definition 9.1.** The *real spinor representation* of the group  $\text{Spin}(s,t)$  is the homomorphism

$$\begin{aligned} \Delta_{s,t}: \text{Spin}(s,t) &\rightarrow \text{GL}(\mathcal{S}_{s,t}) \\ R &\mapsto \rho(R) \end{aligned}$$

given by restricting an irreducible real representation  $\rho: \mathcal{G}(\mathbb{R}^{s,t}) \rightarrow \text{End}_{\mathbb{R}}(\mathcal{S}_{s,t})$  to  $\text{Spin} \subseteq \mathcal{G}^+ \subseteq \mathcal{G}$ . Analogously, a real spinor representation of the rotor group is obtained by restricting to  $\text{Spin}^+(s,t)$ .

**Proposition 9.2.** *Depending on  $s$  and  $t$ , the representation  $\Delta_{s,t}$  decomposes as*

$$\begin{aligned} \Delta_{s,t} &= \Delta^+ \oplus \Delta^-, \\ \Delta_{s,t} &= \Delta^0 \oplus \Delta^0, \text{ or} \\ \Delta_{s,t} &= \Delta^0, \end{aligned}$$

<sup>23</sup>Sometimes called a *space of pinors*. Beware that the terminology regarding spinor spaces varies a lot in the literature, and can unfortunately be quite confusing.

where  $\Delta^+$ ,  $\Delta^-$ ,  $\Delta^0$  in the respective cases denote inequivalent irreducible representations of  $\text{Spin}(s, t)$ . This decomposition of  $\Delta_{s,t}$  is independent of which irreducible representation  $\rho$  is chosen in the definition (in the case that there are two inequivalent such representations). We write  $\mathcal{S}_{s,t} = \mathcal{S}_{\mathbb{K}}^+ \oplus \mathcal{S}_{\mathbb{K}}^-$ ,  $\mathcal{S}_{s,t} = \mathcal{S}_{\mathbb{K}}^0 \oplus \mathcal{S}_{\mathbb{K}}^0$ , and  $\mathcal{S}_{s,t} = \mathcal{S}_{\mathbb{K}}^0$ , for the respective representation spaces, where  $\mathbb{K}$  tells whether the representation is real, complex or quaternionic. The different cases of  $(s, t)$  modulo 8 are listed in Table 9.1.

*Proof.* This follows from Proposition 9.1 and the identification

$$\begin{array}{ccccc} \text{Spin}(s, t) & \subseteq & \mathcal{G}^+(\mathbb{R}^{s,t}) & \subseteq & \mathcal{G}(\mathbb{R}^{s,t}) & \rightarrow & \text{End } \mathcal{S}_{s,t} \\ \uparrow & & \uparrow & & & & \\ \text{Pin}(s, t-1) & \subseteq & \mathcal{G}(\mathbb{R}^{s,t-1}) & & & & \rightarrow \text{End } \mathcal{S}_{s,t-1} \end{array}$$

or

$$\begin{array}{ccccc} \text{Spin}(s, t) & \subseteq & \mathcal{G}^+(\mathbb{R}^{s,t}) & \subseteq & \mathcal{G}(\mathbb{R}^{s,t}) & \rightarrow & \text{End } \mathcal{S}_{s,t} \\ \uparrow & & \uparrow & & & & \\ \text{Pin}(t, s-1) & \subseteq & \mathcal{G}(\mathbb{R}^{t,s-1}) & & & & \rightarrow \text{End } \mathcal{S}_{t,s-1} \end{array}$$

induced by the isomorphism  $F$  in the proof of Proposition 5.1.  $\square$

**Example 9.1.** The spinor module of the space algebra  $\mathcal{G}(\mathbb{R}^3)$  is irreducible w.r.t. the spatial rotor group  $\text{Spin}(3, 0)$ ,

$$\mathcal{S}_{3,0} = \mathbb{C}^2 = \mathcal{S}_{\mathbb{H}}^0,$$

which corresponds to the standard action of  $\text{SU}(2)$  (the unit quaternions) on  $\mathbb{C}^2$  (see Section 6.6.7). On the other hand, the spinor module of the spacetime algebra  $\mathcal{G}(\mathbb{R}^{1,3})$  decomposes into a pair of equivalent four-dimensional irreducible representations of  $\text{Spin}(1, 3)$ ,

$$\mathcal{S}_{1,3} = \mathbb{H}^2 = \mathcal{S}_{\mathbb{C}}^0 \oplus \mathcal{S}_{\mathbb{C}}^0,$$

corresponding to two  $\mathbb{C}^2$ -representations for  $\text{SL}(2, \mathbb{C})$  (see Section 6.6.8). There is also an interesting decomposition in eight dimensions,

$$\mathcal{S}_{8,0} = \mathcal{S}_{0,8} = \mathbb{R}^{16} = \mathcal{S}_{\mathbb{R}}^+ \oplus \mathcal{S}_{\mathbb{R}}^-,$$

which is related to the notion of *triality* (see e.g. [28] and references therein).

**Definition 9.2.** The *complex spinor representation* of  $\text{Spin}(s, t)$  is the homomorphism (where  $n = s + t$ )

$$\begin{array}{ccc} \Delta_n^{\mathbb{C}}: \text{Spin}(s, t) & \rightarrow & \text{GL}_{\mathbb{C}}(\mathcal{S}) \\ R & \mapsto & \rho(R) \end{array}$$

obtained by restricting an irreducible complex representation  $\rho : \mathcal{G}(\mathbb{C}^{s+t}) \rightarrow \text{End}_{\mathbb{C}}(\mathcal{S})$  to  $\text{Spin}(s, t) \subseteq \mathcal{G}^+(\mathbb{R}^{s,t}) \subseteq \mathcal{G}(\mathbb{R}^{s,t}) \otimes \mathbb{C} \cong \mathcal{G}(\mathbb{C}^{s+t})$ .

**Proposition 9.3.** *When  $n$  is odd, the representation  $\Delta_n^{\mathbb{C}}$  is irreducible and independent of which irreducible representation  $\rho$  is chosen in the definition. When  $n$  is even, there is a decomposition*

$$\Delta_n^{\mathbb{C}} = \Delta_{\mathbb{C}}^+ \oplus \Delta_{\mathbb{C}}^-$$

*into a direct sum of two inequivalent irreducible complex representations of  $\text{Spin}(n)$ .*



In physics, spinors of the complex representation  $\Delta_n^{\mathbb{C}}$  are usually called *Dirac spinors*, and  $\Delta_{\mathbb{C}}^+$ ,  $\Delta_{\mathbb{C}}^-$  are *left-* and *right-handed Weyl (or chiral) spinor* representations. Elements of the real representations  $\mathcal{S}_{s,t}$  are called *Majorana spinors*, while those of  $\mathcal{S}_{\mathbb{K}}^+$ ,  $\mathcal{S}_{\mathbb{K}}^-$  are *left-* and *right-handed Majorana–Weyl spinors*.

Note that the spinor representations defined above do not descend to representations of  $\text{SO}(s,t)$  since  $\Delta_{s,t}^{(\mathbb{C})}(-1) = -\text{id}$ . Actually, together with the canonical tensor representations

$$\begin{aligned} \text{ad}|_{\mathcal{G}^1} : \quad \text{ad}_{\frac{1}{2}e_i \wedge e_j}(v) &= \frac{1}{2}[e_i \wedge e_j, v] = (e_i \wedge e_j) \lrcorner v = v_j e_i - v_i e_j, \\ \text{ad}|_{\mathcal{G}^k} : \quad \text{ad}_{\frac{1}{2}e_i \wedge e_j}(x) &= \sum_{r=1}^k \sum_{i_1, \dots, i_k} v_{i_1} \wedge \dots \wedge \text{ad}_{\frac{1}{2}e_i \wedge e_j}(v_{i_r}) \wedge \dots \wedge v_{i_k}, \end{aligned}$$

where  $v = \sum_i v_i e_i$ ,  $x = \sum_{i_1, \dots, i_k} v_{i_1} \wedge \dots \wedge v_{i_k} \in \mathcal{G}^k$ , and  $\{e_i\}_{i=1}^n$  is an orthonormal basis of  $\mathbb{R}^n$ , these additional representations provide us with the full set of so called *fundamental representations* of the Lie algebras  $\mathfrak{so}(n)$  and the corresponding Lie groups  $\text{Spin}(n)$ .

We refer to [27, 15] for more on this approach to spinor spaces, and proofs of the above propositions.

## 9.2 Spinors as elements of ideals

The above spaces of spinors can actually be considered as subspaces of the geometric algebra  $\mathcal{G}$  itself, namely as minimal left ideals of  $\mathcal{G}$ , with the action given by left multiplication on the algebra. For now, we refer to [28] for this approach.

## 9.3 Mixed-action spinors

A subspace of spinors  $\mathcal{S} \subseteq \mathcal{G}$  can of course be extended from a minimal left ideal of  $\mathcal{G}$  to some larger invariant subspace, like a sum of such ideals, or the whole algebra  $\mathcal{G}$ . However, it is possible to consider other, intermediate, subspaces  $\mathcal{S}$  by, instead of only left action, also taking advantage of right action. For now, we refer to [12, 29] for examples of this approach.



## 10 Some Clifford analysis on $\mathbb{R}^n$

This section is currently incomplete. We refer to e.g. [12, 14, 21].

## 11 The algebra of Minkowski spacetime

This section is under construction. For now, we refer to chapters 5,7 and 8 in [12]. See also e.g. [1, 20].

### 11.1 Relative space

For now, we refer to chapters 5,7 and 8 in [12].

### 11.2 Dirac-Hestenes spinors

The equation which describes the evolution of a Dirac spinor in spacetime is the Dirac equation, which in this representation is given by the *Dirac-Hestenes equation*,

$$\nabla\Psi\gamma_1\gamma_2 - A\Psi = m\Psi\gamma_0, \quad (11.1)$$

where  $\nabla := \gamma^\mu \frac{\partial}{\partial x^\mu}$  (we use the spacetime coordinate expansion  $x = x^\mu\gamma_\mu$ ) and  $A$  is the electromagnetic potential vector field. Here,  $\gamma_1\gamma_2$  again plays the role of the complex imaginary unit  $i$ .

### 11.3 Electromagnetic fields

Another interesting property of the spacetime algebra is that the electromagnetic field is most naturally represented as a bivector field in  $\mathcal{G}(\mathbb{R}^{1,3})$ . We can write any such bivector field as  $F = \mathbf{E} + I\mathbf{B}$ , where  $\mathbf{E}$  and  $\mathbf{B}$  are relative space vector fields. In the context of relativistic electrodynamics, these are naturally interpreted as the electric and magnetic fields, respectively. Maxwell's equations are compactly written as

$$\nabla F = J \quad (11.2)$$

in this formalism, where  $J$  is the source current. The physical quantity describing the energy and momentum present in an electromagnetic field is the *Maxwell stress-energy tensor* which in the STA formulation can be interpreted as a map  $T: \mathbb{R}^{1,3} \rightarrow \mathbb{R}^{1,3}$ , given by

$$T(x) := -\frac{1}{2}FxF = \frac{1}{2}FxF^\dagger. \quad (11.3)$$

For example, the energy of the field  $F$  relative to the  $\gamma_0$ -direction is  $\gamma_0 * T(\gamma_0) = \frac{1}{2}(\mathbf{E}^2 + \mathbf{B}^2)$ .

Rodrigues and Vaz [45, 46] have studied an interesting correspondence between the Dirac and Maxwell equations. With the help of the following theorem, they have proved that the electromagnetic field can be obtained from a spinor field satisfying an equation similar to the Dirac equation. This theorem also serves to illustrate how efficiently computations can be performed in the STA framework.

**Theorem 11.1.** *Any bivector  $F \in \mathcal{G}^2(\mathbb{R}^{1,3})$  can be written as*

$$F = \Psi\gamma_0\gamma_1\Psi^\dagger,$$

for some (nonzero) spinor  $\Psi \in \mathcal{G}^+(\mathbb{R}^{1,3})$ .

*Proof.* Take any bivector  $F = \mathbf{E} + I\mathbf{B} \in \mathcal{G}^2$ . Note that

$$F^2 = (\mathbf{E}^2 - \mathbf{B}^2) + 2\mathbf{E} * \mathbf{B}I = \rho e^{\phi I} \quad (11.4)$$

for some  $\rho \geq 0$  and  $0 \leq \phi < 2\pi$ . We consider the cases  $F^2 \neq 0$  and  $F^2 = 0$  separately.

If  $F^2 \neq 0$  then  $\mathbf{E}^2 - \mathbf{B}^2$  and  $\mathbf{E} * \mathbf{B}$  are not both zero and we can apply a *duality rotation* of  $F$  into

$$F' = \mathbf{E}' + I\mathbf{B}' := e^{-\phi I/4} F e^{-\phi I/4} \Rightarrow F'^2 = \rho, \quad (11.5)$$

i.e. such that  $\mathbf{E}'^2 - \mathbf{B}'^2 > 0$  and  $\mathbf{E}' * \mathbf{B}' = 0$ . Hence, we can select an orthonormal basis  $\{\mathbf{e}_i\}$  of the relative space, aligned so that  $\mathbf{E}' = E'\mathbf{e}_1$  and  $\mathbf{B}' = B'\mathbf{e}_2$ , where  $E' = |\mathbf{E}'|$  etc. Consider now a boost  $\mathbf{a} = \alpha\mathbf{e}_3$  of angle  $\alpha$  in the direction orthogonal to both  $\mathbf{E}'$  and  $\mathbf{B}'$ . Using that

$$e^{-\frac{\alpha}{2}\mathbf{e}_3} \mathbf{e}_1 e^{\frac{\alpha}{2}\mathbf{e}_3} = \mathbf{e}_1 e^{\alpha\mathbf{e}_3} = \mathbf{e}_1 (\cosh \alpha + \sinh \alpha \mathbf{e}_3) \quad (11.6)$$

and likewise for  $\mathbf{e}_2$ , we obtain

$$\begin{aligned} F'' &:= e^{-\alpha/2} F' e^{\alpha/2} = E' e^{-\frac{\alpha}{2}\mathbf{e}_3} \mathbf{e}_1 e^{\frac{\alpha}{2}\mathbf{e}_3} + I B' e^{-\frac{\alpha}{2}\mathbf{e}_3} \mathbf{e}_2 e^{\frac{\alpha}{2}\mathbf{e}_3} \\ &= E' \mathbf{e}_1 (\cosh \alpha + \sinh \alpha \mathbf{e}_3) + I B' \mathbf{e}_2 (\cosh \alpha + \sinh \alpha \mathbf{e}_3) \\ &= (E' \cosh \alpha - B' \sinh \alpha) \mathbf{e}_1 + I (B' \cosh \alpha - E' \sinh \alpha) \mathbf{e}_2 \\ &= \cosh \alpha ((E' - B' \tanh \alpha) \mathbf{e}_1 + I (B' - E' \tanh \alpha) \mathbf{e}_2), \end{aligned} \quad (11.7)$$

where we also noted that  $\mathbf{e}_1 \mathbf{e}_3 = -I\mathbf{e}_2$  and  $I\mathbf{e}_2 \mathbf{e}_3 = -\mathbf{e}_1$ . Since  $E'^2 - B'^2 > 0$  we can choose  $\alpha := \operatorname{arctanh}(\frac{B'}{E'})$  and obtain  $F'' = \sqrt{1 - (\frac{B'}{E'})^2} \mathbf{E}' = E'' \mathbf{e}_1$ , where  $E'' > 0$ . Finally, some relative space rotor  $e^{Ib/2}$  takes  $\mathbf{e}_1$  to our timelike target blade (relative space vector)  $\gamma_0 \gamma_1$ , i.e.

$$F'' = E'' e^{Ib/2} \gamma_0 \gamma_1 e^{-Ib/2}. \quad (11.8)$$

Summing up, we have that  $F = \Psi \gamma_0 \gamma_1 \Psi^\dagger$ , where

$$\Psi = \sqrt{E''} e^{\phi I/4} e^{\alpha/2} e^{Ib/2} \in \mathcal{G}^+. \quad (11.9)$$

When  $F^2 = 0$  we have that both  $\mathbf{E}^2 = \mathbf{B}^2$  and  $\mathbf{E} * \mathbf{B} = 0$ . Again, we select an orthonormal basis  $\{\mathbf{e}_i\}$  of the relative space so that  $\mathbf{E} = E\mathbf{e}_1$  and  $\mathbf{B} = B\mathbf{e}_2 = E\mathbf{e}_2$ . Note that

$$\begin{aligned} (1 - I\mathbf{e}_1 \mathbf{e}_2) \mathbf{e}_1 (1 + I\mathbf{e}_1 \mathbf{e}_2) &= \mathbf{e}_1 - I\mathbf{e}_1 \mathbf{e}_2 \mathbf{e}_1 + I\mathbf{e}_2 - I\mathbf{e}_1 \mathbf{e}_2 I\mathbf{e}_2 \\ &= 2(\mathbf{e}_1 + I\mathbf{e}_2). \end{aligned} \quad (11.10)$$

Thus,  $\frac{1}{\sqrt{2}}(1 - I\mathbf{e}_1 \mathbf{e}_2) \mathbf{E} \frac{1}{\sqrt{2}}(1 + I\mathbf{e}_1 \mathbf{e}_2) = \mathbf{E} + I\mathbf{B}$ . Using that  $\mathbf{e}_1$  can be obtained from  $\gamma_0 \gamma_1$  with some relative space rotor  $e^{Ib/2}$ , we have that  $F = \Psi \gamma_0 \gamma_1 \Psi^\dagger$ , where

$$\Psi = (\frac{E}{2})^{1/2} (1 - \frac{1}{E^2} I\mathbf{E}\mathbf{B}) e^{Ib/2} \in \mathcal{G}^+. \quad (11.11)$$

The case  $F = 0$  can be achieved not only using  $\Psi = 0$ , but also with e.g.  $\Psi = (1 + \gamma_0 \gamma_1)$ .  $\square$

Note that we can switch  $\gamma_0\gamma_1$  for any other non-null reference blade, e.g.  $\gamma_1\gamma_2$ .

*Remark.* In the setting of electrodynamics, where  $F = \mathbf{E} + I\mathbf{B}$  is an electromagnetic field, we obtain as a consequence of this theorem and proof the following result due to Rainich, Misner and Wheeler. If we define an *extremal field* as a field for which the magnetic (electric) field is zero and the electric (magnetic) field is parallel to one coordinate axis, the theorem of Rainich-Misner-Wheeler says that: “At any point of Minkowski spacetime any nonnull electromagnetic field can be reduced to an extremal field by a Lorentz transformation and a duality rotation.”

# A Appendix

## A.1 Notions in algebra

In order to make the presentation of this course easier to follow also without a wide mathematical knowledge, we give here a short summary of basic notions in algebra, most of which will be used in the course.

### A.1.1 Basic notions

A *binary composition*  $*$  on a set  $M$  is a map

$$\begin{aligned} M \times M &\rightarrow M \\ (x, y) &\mapsto x * y \end{aligned}$$

The composition is called

*associative* if  $(x * y) * z = x * (y * z) \forall x, y, z \in M$ ,

*commutative* if  $x * y = y * x \forall x, y \in M$ .

If there exists an element  $e \in M$  such that  $e * x = x * e = x \forall x \in M$  then  $e$  is called a *unit*. Units are unique because, if  $e$  and  $e'$  are units, then  $e = e * e' = e'$ .

An element  $x \in M$  has a *left inverse*  $y$  if  $y * x = e$ , *right inverse*  $z$  if  $x * z = e$ , and *inverse*  $y$  if  $y * x = x * y = e$ . If  $*$  is associative and if  $y$  and  $z$  are inverses to  $x$  then

$$z = z * e = z * (x * y) = (z * x) * y = e * y = y.$$

In other words, inverses are unique whenever  $*$  is associative.

Let  $*$  and  $\diamond$  be two binary compositions on  $M$ . We say that  $*$  is *distributive* over  $\diamond$  if

$$x * (y \diamond z) = (x * y) \diamond (x * z) \quad \forall x, y, z \in M$$

and

$$(y \diamond z) * x = (y * x) \diamond (z * x) \quad \forall x, y, z \in M.$$

From these notions one can define a variety of common mathematical structures as follows (see Example A.1 for examples).

**Monoid:** A set with an associative binary composition.

**Group:** Monoid with unit, where every element has an inverse.

**Abelian group:** Commutative group.

**Ring:** A set  $R$  with two binary compositions, called *addition*  $(+)$  and *multiplication*  $(\cdot)$ , such that  $(R, +)$  is an abelian group and  $(R, \cdot)$  is a monoid, and where multiplication is distributive over addition. Furthermore, it should hold that  $0 \cdot x = x \cdot 0 = 0$  for all  $x \in R$ , where  $0$  denotes the additive unit in  $R$  and is called *zero*.

**Ring with unit:** Ring where multiplication has a unit, often denoted  $1$  and called *one* or *the identity*.

**Characteristic:** The characteristic,  $\text{char } R$ , of a ring  $R$  with unit 1 is defined to be the smallest integer  $n > 0$  such that  $\underbrace{1 + \dots + 1}_{n \text{ terms}} = 0$ . If no such  $n$  exists, then we define  $\text{char } R = 0$ .

**Commutative ring:** Ring with commutative multiplication.

**(Skew) Field:** (Non-)Commutative ring with unit, where every nonzero element has a multiplicative inverse.

**Module:** An abelian group  $(M, \oplus)$  is called a *module* over the ring  $(R, +, \cdot)$  if we are given a map (often called *scalar multiplication*)

$$\begin{aligned} R \times M &\rightarrow M \\ (r, m) &\mapsto rm \end{aligned}$$

such that for all  $r, r' \in R$  and  $m, m' \in M$

- i)  $0_R m = 0_M$
- ii)  $1m = m$ , if  $R$  has the multiplicative unit 1
- iii)  $(r + r')m = (rm) \oplus (r'm)$
- iv)  $r(m \oplus m') = (rm) \oplus (rm')$
- v)  $(r \cdot r')m = r(r'm)$

*Remark.* Usually, both the zero  $0_R$  in  $R$  and the additive unit  $0_M$  in  $M$  are denoted 0. Also,  $\oplus$  is denoted  $+$  and is called addition as well.

**Submodule:** If  $A$  is a non-empty subset of an  $R$ -module  $B$  then  $A$  is called a *submodule* of  $B$  if

$$x, y \in A \Rightarrow x + y \in A \quad \forall x, y$$

and

$$r \in R, x \in A \Rightarrow rx \in A \quad \forall r, x,$$

i.e.  $A$  is closed under addition and multiplication by scalars. (In general, given some structure  $A$  (module, group, ...) a substructure is a non-empty subset of  $A$  that is closed under all the operations of the structure.)

**Ideal:** A non-empty subset  $J$  of a ring  $R$  is called a (*two-sided*) *ideal* if

$$x, y \in J \Rightarrow x + y \in J \quad \forall x, y$$

and

$$x \in J, r, q \in R \Rightarrow rxq \in J \quad \forall x, r, q.$$

$J$  is called a *left-sided* resp. *right-sided ideal* if the latter condition is replaced with

$$x \in J, r \in R \Rightarrow rx \in J \quad \forall x, r$$

resp.

$$x \in J, r \in R \Rightarrow xr \in J \quad \forall x, r.$$

**Span:** Given a module  $M$  over  $R$ ,  $\text{Span}_R X$  denotes the linear span over  $R$  of a subset  $X \subseteq M$ . It is a submodule of  $M$ .

**Vector space:** Module over a field.

**Hilbert space:** Vector space over  $\mathbb{R}$  or  $\mathbb{C}$  with a hermitian inner product (bilinear and symmetric for  $\mathbb{R}$  or sesquilinear for  $\mathbb{C}$ ) such that the induced topology of the corresponding norm is complete (i.e. Cauchy sequences converge).

**R-algebra:** An  $R$ -module  $A$  is called an  $R$ -algebra if there is defined an  $R$ -bilinear map  $A \times A \xrightarrow{*} A$ , usually called multiplication.

**Associative R-algebra:**  $R$ -algebra with associative multiplication (hence also a ring).

**Associative R-algebra with unit:** An associative  $R$ -algebra  $A$  with unit  $1_A$  (hence a ring with unit) and with an injective ring homomorphism  $R \hookrightarrow A$ , mapping the unit 1 in  $R$  to the unit  $1_A$  in  $A$ .

**Lie algebra:** (Typically non-associative)  $R$ -algebra  $A$  with multiplication commonly denoted  $[\cdot, \cdot]$  s.t.

$$\begin{aligned} [x, x] &= 0, & (\text{antisymmetry}) \\ [x, [y, z]] + [y, [z, x]] + [z, [x, y]] &= 0, & (\text{Jacobi identity}) \end{aligned}$$

for all  $x, y, z \in A$ .

**Lie group:** A group which is also a differentiable manifold (a topological space which is essentially composed of patches of  $\mathbb{R}^n$  that are smoothly glued together) and such that the group operations are smooth.

**Example A.1.**

- $\mathbb{N} = \{1, 2, 3, \dots\}$  with addition is a monoid,
- $U(1)$  (unit complex numbers) with multiplication is an abelian group,
- $(\mathbb{Z}, +, \cdot)$  is a commutative ring with characteristic 0,
- $\mathbb{Q}, \mathbb{R}$ , and  $\mathbb{C}$  are fields,
- $\mathbb{H}$  is a skew field,
- $\mathbb{R}$  is a module over  $\mathbb{Z}$  and a (infinite-dimensional) vector space over  $\mathbb{Q}$ ,
- $\mathbb{C}^n$  is a vector space over  $\mathbb{C}$ , and a Hilbert space with the standard sesquilinear inner product  $\bar{x}^T y = \sum_{j=1}^n \bar{x}_j y_j$ ,
- the set  $\mathbb{R}^{n \times n}$  of real  $n \times n$  matrices with matrix multiplication is a non-commutative associative  $\mathbb{R}$ -algebra with unit  $1_{n \times n} = \text{diag}(1, 1, \dots, 1)$ ,
- $\mathbb{R}^3$  with the cross product  $\times$  is a Lie algebra over  $\mathbb{R}$ ,

· all the following classical groups are Lie groups:

$$\begin{aligned}
\mathrm{GL}(n, \mathbb{F}) &= \{g \in \mathbb{F}^{n \times n} : \det_{\mathbb{F}} g \neq 0\}, \\
\mathrm{SL}(n, \mathbb{F}) &= \{g \in \mathbb{F}^{n \times n} : \det_{\mathbb{F}} g = 1\}, \\
\mathrm{O}(n) &= \{g \in \mathbb{R}^{n \times n} : \langle gx, gy \rangle_{\mathbb{R}} = \langle x, y \rangle_{\mathbb{R}} \ \forall x, y \in \mathbb{R}^n\}, \\
\mathrm{SO}(n) &= \{g \in \mathrm{O}(n) : \det_{\mathbb{R}} g = 1\}, \\
\mathrm{U}(n) &= \{g \in \mathbb{C}^{n \times n} : \langle gx, gy \rangle_{\mathbb{C}} = \langle x, y \rangle_{\mathbb{C}} \ \forall x, y \in \mathbb{C}^n\}, \\
\mathrm{SU}(n) &= \{g \in \mathrm{U}(n) : \det_{\mathbb{C}} g = 1\}, \\
\mathrm{Sp}(n) &= \{g \in \mathbb{H}^{n \times n} : \langle gx, gy \rangle_{\mathbb{H}} = \langle x, y \rangle_{\mathbb{H}} \ \forall x, y \in \mathbb{H}^n\},
\end{aligned}$$

where  $\langle x, y \rangle_{\mathbb{K}} = \bar{x}^T y$  denotes the standard hermitian inner product on the corresponding space  $\mathbb{K}^n$ .

**Group homomorphism:** A map  $\varphi : G \rightarrow G'$ , where  $(G, *)$  and  $(G', *)$  are groups, such that

$$\varphi(x * y) = \varphi(x) *' \varphi(y) \quad \forall x, y \in G$$

and  $\varphi(e) = e'$ .

**Module homomorphism:** A map  $\varphi : M \rightarrow M'$ , where  $M$  and  $M'$  are  $R$ -modules, such that

$$\varphi(rx + y) = r\varphi(x) + \varphi(y) \quad \forall r \in R, x, y \in M.$$

A map with this property is called  $R$ -linear, and the set of  $R$ -module homomorphisms  $\varphi : M \rightarrow M'$  (which is itself naturally an  $R$ -module) is denoted  $\mathrm{Hom}_R(M, M')$ .

**R-algebra homomorphism:** An  $R$ -linear map  $\varphi : A \rightarrow A'$ , where  $A$  and  $A'$  are  $R$ -algebras, such that  $\varphi(x *_A y) = \varphi(x) *_{A'} \varphi(y) \ \forall x, y \in A$ . If  $A$  and  $A'$  have units  $1_A$  resp.  $1_{A'}$  then we also require that  $\varphi(1_A) = 1_{A'}$ .

**Isomorphism:** A bijective homomorphism  $\varphi : A \rightarrow B$  (it follows that also  $\varphi^{-1}$  is a homomorphism). We say that  $A$  and  $B$  are isomorphic and write  $A \cong B$ .

**Endomorphism:** A homomorphism from an object to itself. For an  $R$ -module  $M$ , we denote  $\mathrm{End}_R M := \mathrm{Hom}_R(M, M)$ .

**Automorphism:** An endomorphism which is also an isomorphism.

**Relation:** A *relation* on a set  $X$  is a subset  $R \subseteq X \times X$ . If  $(x, y) \in R$  then we say that  $x$  is related to  $y$  and write  $xRy$ .

A relation  $R$  on  $X$  is called

- reflexive* if  $xRx \ \forall x \in X$ ,
- symmetric* if  $xRy \Rightarrow yRx \ \forall x, y \in X$ ,
- antisymmetric* if  $xRy \ \& \ yRx \Rightarrow x = y \ \forall x, y \in X$ ,
- transitive* if  $xRy \ \& \ yRz \Rightarrow xRz \ \forall x, y, z \in X$ .

**Partial order:** A reflexive, antisymmetric, and transitive relation.

**Total order:** A partial order such that either  $xRy$  or  $yRx$  holds  $\forall x, y \in X$ .

**Equivalence relation:** A reflexive, symmetric, and transitive relation.



### A.1.2 Direct products and sums

Suppose we are given an  $R$ -module  $M_i$  for each  $i$  in some index set  $I$ . A function  $f : I \rightarrow \bigcup_{i \in I} M_i$  is called a *section* if  $f(i) \in M_i$  for all  $i \in I$ . The *support* of a section  $f$ ,  $\text{supp } f$ , is the set  $\{i \in I : f(i) \neq 0\}$ .

**Definition A.1.** The (direct) product  $\prod_{i \in I} M_i$  of the modules  $M_i$  is the set of all sections. The (direct) sum  $\bigsqcup_{i \in I} M_i = \bigoplus_{i \in I} M_i$  is the set of all sections with finite support.

If  $I$  is finite,  $I = \{1, 2, \dots, n\}$ , then the product and sum coincide and is also denoted  $M_1 \oplus M_2 \oplus \dots \oplus M_n$ .

The  $R$ -module structure on the direct sum and product is given by “point-wise” summation and  $R$ -multiplication,

$$(f + g)(i) := f(i) + g(i), \quad \text{and} \quad (rf)(i) := r(f(i)),$$

for sections  $f, g$  and  $r \in R$ . If, moreover, the  $R$ -modules  $M_i$  also are  $R$ -algebras, then we can promote  $\prod_{i \in I} M_i$  and  $\bigoplus_{i \in I} M_i$  into  $R$ -algebras as well by defining “pointwise” multiplication

$$(fg)(i) := f(i)g(i).$$

Also note that we have canonical surjective  $R$ -module homomorphisms

$$\pi_j : \prod_{i \in I} M_i \ni f \mapsto f(j) \in M_j,$$

and canonical injective  $R$ -module homomorphisms

$$\sigma_j : M_j \ni m \mapsto \sigma_j(m) \in \bigoplus_{i \in I} M_i,$$

defined by  $\sigma_j(m)(i) := (i = j)m$ .

Let us consider the case  $M_i = R$ , for all  $i \in I$ , in a little more detail. Denote by  $X$  an arbitrary set and  $R$  a ring with unit. If  $f : X \rightarrow R$  then the support of  $f$  is given by

$$\text{supp } f = \{x \in X : f(x) \neq 0\}.$$

**Definition A.2.** The direct sum of  $R$  over  $X$ , or the free  $R$ -module generated by  $X$ , is the set of all functions from  $X$  to  $R$  with finite support, and is denoted  $\bigoplus_X R$  or  $\bigsqcup_X R$ .

For  $x \in X$ , we conveniently define  $\delta_x : X \rightarrow R$  through  $\delta_x(y) = (x = y)$ , i.e.  $\delta_x(y) = 0$  if  $y \neq x$ , and  $\delta_x(x) = 1$ . Then, for any  $f : X \rightarrow R$  with finite support, we have

$$f(y) = \sum_{x \in X} f(x)\delta_x(y) \quad \forall y \in X,$$

or, with the  $R$ -module structure taken into account,

$$f = \sum_{x \in X} f(x)\delta_x.$$

This sum is usually written simply  $f = \sum_{x \in X} f(x)x$  or  $\sum_{x \in X} f_x x$ , and is interpreted as a formal sum of the elements in  $X$  with coefficients in  $R$ .

---

**Exercise A.1.** Prove the following universality properties of the product and the sum:

a) Let  $N$  be an  $R$ -module and let  $\rho_j \in \text{Hom}_R(N, M_j)$  for all  $j \in I$ . Show that there exist unique  $\tau_j \in \text{Hom}_R(N, \prod_{i \in I} M_i)$ , such that the diagram below commutes:

$$\begin{array}{ccc} N & & \\ & \searrow^{\tau_j} & \\ \rho_j \downarrow & & \prod_{i \in I} M_i \\ & \swarrow_{\pi_j} & \\ M_j & & \end{array}$$

b) Let  $\rho_j : M_j \rightarrow N$  be in  $\text{Hom}_R(M_j, N)$ . Prove that there exist unique  $\tau_j \in \text{Hom}_R(\bigoplus_{i \in I} M_i, N)$ , making the diagram below commutative:

$$\begin{array}{ccc} M_j & & \\ & \searrow^{\sigma_j} & \\ \rho_j \downarrow & & \bigoplus_{i \in I} M_i \\ & \swarrow_{\tau_j} & \\ N & & \end{array}$$

### A.1.3 Lists

Let  $X$  be an arbitrary set and define the cartesian products of  $X$  inductively through

$$\begin{aligned} X^0 &= \{\emptyset\} \quad (\text{the empty list}) \\ X^1 &= X \\ X^{n+1} &= X \times X^n, \quad n > 0. \end{aligned}$$

Note that  $X$  is isomorphic to  $X \times X^0$ .

**Definition A.3.** List  $X := \bigcup_{n \geq 0} X^n$  is the set of all finite lists of elements in  $X$ .

We have a natural monoid structure on List  $X$  through concatenation of lists:

$$\begin{aligned} \text{List } X \times \text{List } X &\rightarrow \text{List } X \\ (x_1 x_2 \dots x_m, y_1 y_2 \dots y_n) &\mapsto x_1 \dots x_m y_1 \dots y_n \end{aligned}$$

This binary composition, which we shall call multiplication of lists, is obviously associative, and the empty list, which we can call 1, is the unit.

### A.1.4 The free associative $R$ -algebra generated by $X$

Let  $M = \text{List } X$  and define the free associative  $R$ -algebra generated by  $X$ , or the noncommutative polynomial ring over the ring  $R$  in the variables  $X$ , first as an  $R$ -module by

$$R\{X\} := \bigoplus_M R.$$

We shall now define a multiplication on  $R\{X\}$  which extends our multiplication on  $M$ , such that  $R\{X\}$  becomes a ring. We let

$$\begin{aligned} R\{X\} \times R\{X\} &\rightarrow R\{X\} \\ (f, g) &\mapsto fg, \end{aligned}$$

where

$$(fg)(m) := \sum_{m'm''=m} f(m')g(m'').$$

That is, we sum over all (finitely many) pairs  $(m', m'')$  such that  $m' \in \text{supp } f$ ,  $m'' \in \text{supp } g$  and  $m = m'm''$ .

If we interpret  $f$  and  $g$  as formal sums over  $M$ , i.e.

$$f = \sum_{m \in M} f_m m, \quad g = \sum_{m \in M} g_m m,$$

then we obtain

$$fg = \sum_{m', m'' \in M} f_{m'} g_{m''} m' m''.$$

*Remark.* We could of course also promote  $\bigoplus_M R$  to a ring in this way if we only required that  $M$  is a monoid and  $R$  a ring with unit.

**Exercise A.2.** Verify that  $R\{X\}$  is a ring with unit.

### A.1.5 Quotients

Let  $A$  be a submodule of the the  $R$ -module  $B$  and define, for  $x \in B$ ,

$$x + A := \{x + y : y \in A\}.$$

Sets of the form  $x + A$  are called *residue classes* and the set of these is denoted  $B/A$ , i.e.

$$B/A = \{x + A : x \in B\}.$$

The residue class  $x + A$  with representative  $x \in B$  is often denoted  $[x]$ .

We will now promote the set  $B/A$  to an  $R$ -module (called a *quotient module*). Addition is defined by

$$\begin{aligned} B/A \times B/A &\xrightarrow{+} B/A \\ (x + A, y + A) &\mapsto x + y + A. \end{aligned}$$

We have to verify that addition is well defined, so assume that  $x + A = x' + A$  and  $y + A = y' + A$ . Then we must have  $x - x' \in A$  and  $y - y' \in A$  so that

$$x' + y' + A = x + y + (x' - x) + (y' - y) + A.$$

Hence,  $x' + y' + A \subseteq x + y + A$  and similarly  $x + y + A \subseteq x' + y' + A$ . This shows that  $x' + y' + A = x + y + A$ .

Multiplication with ring elements is defined similarly by

$$\begin{aligned} R \times B/A &\rightarrow B/A \\ (r, x + A) &\mapsto rx + A. \end{aligned}$$

---

**Exercise A.3.** Verify that the  $R$ -multiplication above is well defined and that  $B/A$  becomes a module over  $R$ .

---

If instead  $J$  is an ideal in the ring  $R$ , we let

$$R/J = \{x + J : x \in R\}$$

and define addition through

$$\begin{aligned} R/J \times R/J &\xrightarrow{+} R/J \\ (x + J, y + J) &\mapsto x + y + J, \end{aligned}$$

and multiplication through

$$\begin{aligned} R/J \times R/J &\rightarrow R/J \\ (x + J, y + J) &\mapsto xy + J. \end{aligned}$$

We leave to the reader to verify that the addition and multiplication defined above are well defined and that they promote  $R/J$  to a ring.

**Example A.2.**  $n\mathbb{Z}$  with  $n$  a positive integer is an ideal in  $\mathbb{Z}$ , and the quotient  $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z}$  is a ring with characteristic  $n$  (and a field when  $n$  is prime).

### A.1.6 Tensor products of modules

Let  $A$  and  $B$  be  $R$ -modules and let  $D$  be the smallest submodule of  $\bigoplus_{A \times B} R$  containing the set

$$\left\{ (a + a', b) - (a, b) - (a', b), (a, b + b') - (a, b) - (a, b'), (ra, b) - r(a, b), \right. \\ \left. (a, rb) - r(a, b) : a, a' \in A, b, b' \in B, r \in R \right\}.$$

Then the tensor product  $A \otimes_R B$  (or simply  $A \otimes B$ ) of  $A$  and  $B$  is defined by

$$A \otimes_R B := \bigoplus_{A \times B} R / D.$$

The point with the above construction is the following

**Theorem A.1** (Universality). *To each bilinear  $A \times B \xrightarrow{\beta} C$ , where  $A, B$  and  $C$  are  $R$ -modules, there exists a unique linear  $\lambda : A \otimes B \rightarrow C$  such that the diagram below commutes*

$$\begin{array}{ccc} A \times B & \xrightarrow{\beta} & C \\ \pi \downarrow & \nearrow \lambda & \\ A \otimes B & & \end{array}$$

Here,  $\pi : A \times B \rightarrow A \otimes B$  is the canonical projection that satisfies  $\pi(a, b) = a \otimes b := (a, b) + D$ , where  $(a, b)$  on the right hand side is an element in  $\bigoplus_{A \times B} R$ .

---

**Exercise A.4.** Verify the above theorem.

### A.1.7 Sums and products of algebras

Let  $A$  and  $B$  be  $R$ -algebras with multiplication maps  $*_A$  resp.  $*_B$ . We have defined the direct sum  $A \oplus B$  as an  $R$ -algebra through pointwise multiplication,

$$\begin{aligned} (A \oplus B) \times (A \oplus B) &\rightarrow A \oplus B \\ ((a, b), (a', b')) &\mapsto (a *_A a', b *_B b'). \end{aligned}$$

We can also promote the tensor product  $A \otimes_R B$  into an  $R$ -algebra by introducing the product

$$\begin{aligned} (A \otimes_R B) \times (A \otimes_R B) &\rightarrow A \otimes_R B \\ (a \otimes b, a' \otimes b') &\mapsto (a *_A a') \otimes (b *_B b') \end{aligned}$$

and extending linearly. If  $1_A$  and  $1_B$  are units on  $A$  resp.  $B$ , then  $(1_A, 1_B)$  becomes a unit on  $A \oplus B$  and  $1_A \otimes 1_B$  becomes a unit on  $A \otimes_R B$ . Note that the above algebras may look similar, but because of multilinearity in the latter case they are very different since e.g.  $1_A \otimes 0 = 0 \otimes 1_B = 0$ , while  $(1_A, 1_B) = (1_A, 0) + (0, 1_B) \neq 0$ .

It is easy to see that if  $A$  and  $B$  are associative  $R$ -algebras, then also  $A \oplus B$  and  $A \otimes_R B$  are associative. Furthermore, it holds that

$$R^{n \times n} \otimes_R A^{m \times m} \cong A^{nm \times nm}, \quad (\text{A.1})$$

as  $R$ -algebras, where  $n, m$  are positive integers, and  $R$  is a ring with unit.

---

**Exercise A.5.** Prove the isomorphism (A.1) by considering the map

$$\begin{aligned} R^{n \times n} \times A^{m \times m} &\xrightarrow{F} A^{nm \times nm} \\ (r, a) &\mapsto F(r, a), \end{aligned}$$

with  $F(r, a)_{(i,j),(k,l)} := r_{i,k} a_{j,l}$ , and  $i, k \in \{1, \dots, n\}$ ,  $j, l \in \{1, \dots, m\}$ .

### A.1.8 The tensor algebra $\mathcal{T}(V)$

The tensor algebra  $\mathcal{T}(V)$  of a module  $V$  over a ring  $R$  can be defined as a direct sum of tensor products of  $V$  with itself,

$$\mathcal{T}(V) := \bigoplus_{k=0}^{\infty} \left( \bigotimes^k V \right),$$

where  $\bigotimes^0 V := R$ ,  $\bigotimes^1 V := V$ ,  $\bigotimes^{k+1} V := (\bigotimes^k V) \otimes_R V$ , and with the obvious associative multiplication (concatenation of tensors) extended linearly,

$$(v_1 \otimes \dots \otimes v_j) \otimes (w_1 \otimes \dots \otimes w_k) := v_1 \otimes \dots \otimes v_j \otimes w_1 \otimes \dots \otimes w_k.$$

Alternatively,  $\mathcal{T}(V)$  can be obtained as the free associative  $R$ -algebra generated by the set  $V$  itself,

$$\mathcal{T}(V) \cong R\{V\}/\hat{D},$$

where  $\hat{D}$  is the ideal generating linearity in each factor (similarly to  $D$  in the definition of the tensor product above). A third alternative definition, when  $R = \mathbb{F}$  is a field, is by simply choosing a basis  $E$  of  $V$ :

$$\mathcal{T}(V) \cong \mathbb{F}\{E\}.$$

---

**Exercise A.6.** Verify the equivalence of these three definitions of  $\mathcal{T}(V)$ .

## A.2 The concept of a category

A *category*  $\mathcal{C}$  consists in the following data:

- **Arrows of  $\mathcal{C}$ :** A class denoted  $\text{ar } \mathcal{C}$ .
- **Objects of  $\mathcal{C}$ :** A class denoted  $\text{ob } \mathcal{C}$ .
- **Domain:** A function  $\text{dom}: \text{ar } \mathcal{C} \rightarrow \text{ob } \mathcal{C}$ .
- **Codomain:** A function  $\text{cod}: \text{ar } \mathcal{C} \rightarrow \text{ob } \mathcal{C}$ . If  $a = \text{dom } f$  and  $b = \text{cod } f$  then we will write  $f: a \rightarrow b$  or  $a \xrightarrow{f} b$  (here objects will usually be denoted  $a, b, c, \dots$ , while arrows (or *morphisms* as they are called) usually are denoted  $f, g, h, \dots$ ).
- **Composition:** For each diagram  $a \xrightarrow{f} b \xrightarrow{g} c$  there is associated a unique arrow  $g \circ f: a \rightarrow c$ . Furthermore, this composition is associative in the following sense: If  $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} d$  is a diagram in  $\mathcal{C}$  then  $h \circ (g \circ f) = (h \circ g) \circ f$ .
- **Units:** For each object  $a$  in  $\text{ob } \mathcal{C}$  there is associated a unique arrow  $1_a$  such that  $\text{dom } 1_a = a = \text{cod } 1_a$ , and if  $a \xrightarrow{f} b$  then  $1_b \circ f = f = f \circ 1_a$ .

*Remark.* We can also consider objects as unit arrows. With this view a category consists only of arrows. Readers interested in foundational questions can use the word “class” informally, or as it is axiomatized in NBG (Neuman-Bernays-Gödel).

### A.2.1 Some examples of categories

Category	Objects	Arrows
Set	all sets	all functions between sets
FinSet	all finite sets	all functions between finite sets
Top	all topological spaces	all continuous functions between topological spaces
Vect	vector spaces	linear mappings
Grp	groups	group homomorphisms
Pos	partially ordered sets	monotone functions

All examples above are “subcategories” of Set, but there are other kinds of categories. The following examples of categories are relevant in this text.

**Example A.3.** Let  $A$  and  $B$  be  $R$ -modules, where  $R$  is a ring. We will denote a category  $\mathcal{C}_{A,B}$  whose objects consist of diagrams  $A \times B \xrightarrow{\beta} C$ , where  $\beta$  is bilinear and  $C$  an  $R$ -module. An arrow  $f$  from  $A \times B \xrightarrow{\beta} C$  to  $A \times B \xrightarrow{\beta'} C'$  is an  $R$ -linear map  $\varphi: C \rightarrow C'$  such that the diagram below commutes:

$$\begin{array}{ccc}
 & & C \\
 & \nearrow \beta & \\
 A \times B & & \\
 & \searrow \beta' & \\
 & & C'
 \end{array}
 \quad \downarrow \varphi$$

Note that the object  $A \times B \xrightarrow{\pi} A \otimes_R B$  is very special in  $\mathcal{C}_{A,B}$  since for every other object  $A \times B \xrightarrow{\beta} C$  there exists a unique arrow from  $A \times B \xrightarrow{\pi} A \otimes_R B$  to  $A \times B \xrightarrow{\beta} C$  (see Theorem A.1). Such an object in a category is called *initial*.

### A.2.2 Universal objects in a category

An object  $i$  in a category  $\mathcal{C}$  is called *initial* if for every object  $a$  in  $\mathcal{C}$  there is a unique arrow  $f_a: i \rightarrow a$  in  $\mathcal{A}$ . An object  $j$  in a category  $\mathcal{C}$  is called *final* if for every object  $b$  in  $\mathcal{C}$  there is a unique arrow  $g_b: b \rightarrow j$  in  $\mathcal{A}$ .

Note that if  $p$  and  $q$  are both initial objects in  $\mathcal{A}$  then there are unique arrows  $p \xrightarrow{f} q$  and  $q \xrightarrow{g} p$  such that  $g \circ f: p \rightarrow p$ . Furthermore, this arrow  $g \circ f$  is also unique since there also is a unique arrow from  $p$  to  $p$ . Since identity on  $p$  is such an arrow, we conclude that  $g \circ f = \text{id}_p$ . Similarly, we obtain  $f \circ g = \text{id}_q$ .

We formulate this by saying that universal objects (initial or final) are unique up to isomorphisms.

**Example A.4.** Let  $V$  be a vector space over the field  $\mathbb{F}$ , and define a category  $\mathcal{C}_V$  in the following way: The objects consist of all diagrams like  $V \xrightarrow{\lambda} \mathcal{A}$ , where  $\lambda$  is linear and  $\mathcal{A}$  an associative  $\mathbb{F}$ -algebra. An arrow from  $V \xrightarrow{\lambda} \mathcal{A}$  to  $V \xrightarrow{\lambda'} \mathcal{A}'$  is an algebra homomorphism  $f: \mathcal{A} \rightarrow \mathcal{A}'$  such that the diagram below commutes:

$$\begin{array}{ccc} V & \xrightarrow{\lambda} & \mathcal{A} \\ & \searrow \lambda' & \downarrow \varphi \\ & & \mathcal{A}' \end{array}$$

Note that  $V \xrightarrow{\tau} \mathcal{T}(V)$  is an initial object in  $\mathcal{C}_V$ , where  $\tau$  denotes the canonical embedding into the tensor algebra (see A.1.8).

**Example A.5.** In our last example, we let  $V \xrightarrow{q} \mathbb{F}$  be a fixed quadratic form on the  $\mathbb{F}$ -vector space  $V$ . Now introduce a category  $\mathcal{C}_{V,q}$  whose objects are diagrams like  $V \xrightarrow{\lambda} \mathcal{A}$ , where  $\mathcal{A}$  is an associative  $\mathbb{F}$ -algebra with unit denoted  $1_{\mathcal{A}}$ ,  $\lambda$  is linear, and  $\lambda(v)^2 = q(v)1_{\mathcal{A}}$  for all  $v \in V$ . An arrow from  $V \xrightarrow{\lambda} \mathcal{A}$  to  $V \xrightarrow{\lambda'} \mathcal{A}'$  is an  $\mathbb{F}$ -algebra homomorphism  $f: \mathcal{A} \rightarrow \mathcal{A}'$  such that the diagram below commutes:

$$\begin{array}{ccc} V & \xrightarrow{\lambda} & \mathcal{A} \\ & \searrow \lambda' & \downarrow \varphi \\ & & \mathcal{A}' \end{array}$$

Here we note that  $V \xrightarrow{j} \mathcal{G}(V,q)$  is an initial object in  $\mathcal{C}_{V,q}$ , where  $j$  denotes the canonical (quotient) map (see Theorem 2.1). Furthermore, if  $V$  has an orthogonal basis  $E$ , then  $V \xrightarrow{i} \mathcal{C}l(E, \mathbb{F}, q|_E)$  is also an initial object in  $\mathcal{C}_{V,q}$ , where  $i$  denotes the canonical embedding map (see Theorem 2.8). Hence these objects must be the same up to isomorphisms.

## A.3 Expansion of the inner product

Let  $\Lambda(n, m)$  denote the set of ordered  $m$ -subsets  $\lambda = (\lambda_1 < \lambda_2 < \dots < \lambda_m)$  of the set  $\{1, 2, \dots, n\}$ , with  $m \leq n$ . The ordered complement of  $\lambda$  is denoted  $\lambda^c = (\lambda_1^c < \lambda_2^c < \dots < \lambda_{n-m}^c)$ . Furthermore, we denote by  $\text{sgn } \lambda$  the sign of the

permutation  $(\lambda_1, \lambda_2, \dots, \lambda_m, \lambda_1^c, \lambda_2^c, \dots, \lambda_{n-m}^c)$ . For an  $n$ -blade  $A = a_1 \wedge \dots \wedge a_n$  we use the corresponding notation  $A_\lambda = a_{\lambda_1} \wedge a_{\lambda_2} \wedge \dots \wedge a_{\lambda_m}$ , etc.

**Theorem A.2.** *For any  $m$ -vector  $x \in \mathcal{G}^m$  and  $n$ -blade  $A \in \mathcal{B}_n$ ,  $m \leq n$ , we have*

$$x \lrcorner A = \sum_{\lambda \in \Lambda(n, m)} (\operatorname{sgn} \lambda) (x * A_\lambda) A_{\lambda^c}. \quad (\text{A.2})$$

*Proof.* First note that both sides of (A.2) are linear in  $x$  and that the l.h.s. is linear and alternating in the  $a_k$ 's. To prove that a multilinear mapping  $F : V \times V \times \dots \times V = V^n \rightarrow W$  is alternating it is enough to show that  $F(x_1, \dots, x_n) = 0$  whenever  $x_s = x_{s+1}$  for  $s = 1, \dots, n-1$ . So for the right hand side, suppose that  $a_s = a_{s+1}$  and consider a fixed  $\lambda \in \Lambda(n, m)$ . If  $\{s, s+1\} \subseteq \lambda$  or  $\{s, s+1\} \subseteq \lambda^c$ , then  $A_\lambda$  or  $A_{\lambda^c}$  vanishes. So let us assume e.g. that  $s \in \lambda$  and  $s+1 \in \lambda^c$ , say  $s = \lambda_i$ ,  $s+1 = \lambda_j^c$ . We define  $\mu \in \Lambda(n, m)$  such that

$$\begin{aligned} (\lambda, \lambda^c) &= (\lambda_1, \lambda_2, \dots, \lambda_i = s, \dots, \lambda_m, \lambda_1^c, \lambda_2^c, \dots, \lambda_j^c = s+1, \dots, \lambda_{n-m}^c) \\ (\mu, \mu^c) &= (\mu_1, \mu_2, \dots, \mu_i = s+1, \dots, \mu_m, \mu_1^c, \mu_2^c, \dots, \mu_j^c = s, \dots, \mu_{n-m}^c) \end{aligned}$$

Then  $\mu \in \Lambda(n, m)$ ,  $\operatorname{sgn} \mu = -\operatorname{sgn} \lambda$ , and since  $a_s = a_{s+1}$ , we have  $A_\lambda = A_\mu$  and  $A_{\lambda^c} = A_{\mu^c}$ . Thus,  $(\operatorname{sgn} \lambda)(x * A_\lambda)A_{\lambda^c} + (\operatorname{sgn} \mu)(x * A_\mu)A_{\mu^c} = 0$ .

We conclude that both sides of (A.2) are alternating and multilinear in  $a_1, \dots, a_n$ . Hence, we may without loss of generality assume that  $x = e_1 \dots e_m$  and  $A = e_1 \dots e_m \dots e_n$  (otherwise both sides are zero), where  $\{e_1, \dots, e_d\}$ ,  $m \leq n \leq d$ , denotes an orthogonal basis for  $V = \mathcal{G}^1$ . But then the theorem is obvious, since

$$\begin{aligned} &\sum_{\lambda \in \Lambda(n, m)} \operatorname{sgn} \lambda ((e_1 \dots e_m) * (e_{\lambda_1} \dots e_{\lambda_m})) e_{\lambda_1^c} \dots e_{\lambda_{n-m}^c} \\ &= ((e_1 \dots e_m) * (e_1 \dots e_m)) e_{m+1} \dots e_d. \end{aligned}$$

□

## A.4 Extension of morphisms

Let  $A$  be an abelian group such that  $a + a = 0$  for all  $a \in A$  (such as  $(\mathbb{Z}_2, +)$  or  $(\mathcal{P}(X), \Delta)$ ) and let  $M$  be a monoid with unit 0. A map  $\varphi : A \rightarrow M$  is then called a homomorphism if  $\varphi(0) = 0$  and  $\varphi(a + b) = \varphi(a) + \varphi(b)$  for all  $a, b \in A$ . We have the following

**Theorem A.3.** *If  $H$  is a subgroup of  $A$  and  $\varphi : H \rightarrow M$  is a homomorphism then there exists an extension of  $\varphi$  to a homomorphism  $\psi : A \rightarrow M$  such that  $\psi|_H = \varphi$ .*

*Proof.* Let  $a \in A \setminus H$  and form  $H_a := H \cup (a + H)$ . Define  $\varphi_a : H_a \rightarrow M$  by  $\varphi_a(h) := \varphi(h)$  if  $h \in H$  and  $\varphi_a(a + h) := \varphi(h)$ . It is easy to see that  $H_a$  is a subgroup of  $A$  and that  $\varphi_a$  is a homomorphism that extends  $\varphi$ .

Let us introduce a partial order  $\preceq$  on pairs  $(H', \varphi')$  where  $H'$  is a subgroup of  $A$  and  $\varphi'$  is a homomorphism  $H' \rightarrow M$  which extends  $\varphi$ . We define  $(H', \varphi') \preceq (H'', \varphi'')$  if  $H' \subseteq H''$  and  $\varphi''|_{H'} = \varphi'$ . By the Hausdorff maximality theorem there exists a maximal chain

$$\mathcal{K} = \{(H^i, \varphi^i) : i \in I\},$$



where  $I$  is a total ordered index set w.r.t  $\leq$ , and

$$i \leq j \Leftrightarrow (H^i, \varphi^i) \preceq (H^j, \varphi^j).$$

Now, let  $H^* := \bigcup_{i \in I} H^i$ . Then  $H^*$  is a subgroup of  $A$  such that  $H^i \subseteq H^* \forall i \in I$ , since if  $a, b \in H^*$  then there exists some  $i \in I$  such that both  $a, b \in H^i$ , i.e.  $a + b \in H^i \subseteq H^*$ . We also define  $\varphi^* : H^* \rightarrow M$  by  $\varphi^*(a) = \varphi^i(a)$  if  $a \in H^i$ . This is well-defined.

Now, if  $H^* \neq A$  then we could find  $a \in A \setminus H^*$  and extend  $\varphi^*$  to a homomorphism  $\varphi_a^* : H_a^* \rightarrow M$  as above. But then  $\mathcal{K} \cup \{(H_a^*, \varphi_a^*)\}$  would be an even greater chain than  $\mathcal{K}$ , which is a contradiction. This proves the theorem.  $\square$

Now, consider the case when  $A = (\mathcal{P}(X), \Delta)$ ,  $H = \mathcal{F}(X)$ ,  $M = \{\pm 1\}$  with multiplication  $\cong \mathbb{Z}_2$ , and  $\varphi(A) = (-1)^{|A|}$  for  $A \in \mathcal{F}(X)$ . We then obtain

**Theorem A.4.** *There exists a homomorphism*

$$\varphi : (\mathcal{P}(X), \Delta) \rightarrow (\{-1, 1\}, \cdot)$$

such that  $\varphi(A) = (-1)^{|A|}$  for finite subsets  $A \subseteq X$ , and  $\varphi(A \Delta B) = \varphi(A)\varphi(B)$  for all  $A, B \subseteq X$ .

In particular, if  $A, B \subseteq X$  are disjoint then  $\varphi(A \cup B) = \varphi(A \Delta B) = \varphi(A)\varphi(B)$ . This extends the even/odd property of finite sets to arbitrary infinite sets; cf. Theorem 4.9.

## A.5 Matrix theorems

In the following theorem we assume that  $R$  is an arbitrary commutative ring and

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1m} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nm} \end{bmatrix} \in R^{n \times m}, \quad a_j = \begin{bmatrix} a_{1j} \\ \vdots \\ a_{nj} \end{bmatrix},$$

i.e.  $a_j$  denotes the  $j$ :th column in  $A$ . If  $I \subseteq \{1, \dots, n\}$  and  $J \subseteq \{1, \dots, m\}$  we let  $A_{I,J}$  denote the  $|I| \times |J|$ -matrix minor obtained from  $A$  by deleting the rows and columns not in  $I$  and  $J$ . Further, let  $k$  denote the rank of  $A$ , i.e. the highest integer  $k$  such that there exists  $I, J$  as above with  $|I| = |J| = k$  and  $\det A_{I,J} \neq 0$ . By renumbering the  $a_{ij}$ :s we can without loss of generality assume that  $I = J = \{1, 2, \dots, k\}$ .

**Theorem A.5** (Basis minor). *If the rank of  $A$  is  $k$ , and*

$$d := \det \begin{bmatrix} a_{11} & \cdots & a_{1k} \\ \vdots & & \vdots \\ a_{k1} & \cdots & a_{kk} \end{bmatrix} \neq 0,$$

then every  $d \cdot a_j$  is a linear combination of  $a_1, \dots, a_k$ .

*Proof.* Pick  $i \in \{1, \dots, n\}$  and  $j \in \{1, \dots, m\}$  and consider the  $(k+1) \times (k+1)$ -matrix

$$B_{i,j} := \begin{bmatrix} a_{11} & \cdots & a_{1k} & a_{1j} \\ \vdots & & \vdots & \vdots \\ a_{k1} & \cdots & a_{kk} & a_{kj} \\ a_{i1} & \cdots & a_{ik} & a_{ij} \end{bmatrix}.$$

Then  $\det B_{i,j} = 0$ . Expanding  $\det B_{i,j}$  along the bottom row for fixed  $i$  we obtain

$$a_{i1}C_1 + \dots + a_{ik}C_k + a_{ij}d = 0, \quad (\text{A.3})$$

where the  $C_l$  are independent of the choice of  $i$  (but of course dependent on  $j$ ). Hence,

$$C_1a_1 + \dots + C_ka_k + da_j = 0, \quad (\text{A.4})$$

and similarly for all  $j$ .  $\square$

The following shows that the basic factorization  $\det(AB) = \det(A)\det(B)$  is a unique property of the determinant.

**Theorem A.6** (Uniqueness of determinant). *Assume that  $d: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$  is continuous and satisfies*

$$d(AB) = d(A)d(B) \quad (\text{A.5})$$

for all  $A, B \in \mathbb{R}^{n \times n}$ . Then  $d$  must be either 0, 1,  $|\det|^\alpha$  or  $(\text{sign} \circ \det)|\det|^\alpha$  for some  $\alpha > 0$ .

*Proof.* First, we have that ( $I$  here denotes the unit matrix)

$$\begin{aligned} d(0) &= d(0^2) = d(0)^2, \\ d(I) &= d(I^2) = d(I)^2, \end{aligned}$$

so  $d(0)$  and  $d(I)$  must be either 0 or 1. Furthermore,

$$\begin{aligned} d(0) &= d(0A) = d(0)d(A), \\ d(A) &= d(IA) = d(I)d(A), \end{aligned}$$

for all  $A \in \mathbb{R}^{n \times n}$ , which implies that  $d = 1$  if  $d(0) = 1$  and  $d = 0$  if  $d(I) = 0$ . We can therefore assume that  $d(0) = 0$  and  $d(I) = 1$ .

Now, an arbitrary matrix  $A$  can be written as

$$A = E_1E_2 \dots E_kR,$$

where  $R$  is on reduced row-echelon form (reduced as much as possible by Gaussian elimination) and  $E_i$  are elementary row operations of the form

$$\begin{aligned} R_{ij} &:= (\text{swap rows } i \text{ and } j), \\ E_i(\lambda) &:= (\text{scale row } i \text{ by } \lambda), \text{ or} \\ E_{ij}(c) &:= (\text{add } c \text{ times row } j \text{ to row } i). \end{aligned}$$

Because  $R_{ij}^2 = I$ , we must have  $d(R_{ij}) = \pm 1$ . This gives, since

$$E_i(\lambda) = R_{1i}E_1(\lambda)R_{1i},$$

that  $d(E_i(\lambda)) = d(E_1(\lambda))$  and

$$d(\lambda I) = d(E_1(\lambda) \dots E_n(\lambda)) = d(E_1(\lambda)) \dots d(E_n(\lambda)) = d(E_1(\lambda))^n.$$

In particular, we have  $d(E_1(0)) = 0$  and of course  $d(E_1(1)) = d(I) = 1$ .

If  $A$  is invertible, then  $R = I$ . Otherwise,  $R$  must contain a row of zeros so that  $R = E_i(0)R$  for some  $i$ . But then  $d(R) = 0$  and  $d(A) = 0$ . When  $A$  is invertible we have  $I = AA^{-1}$  and  $1 = d(I) = d(A)d(A^{-1})$ , i.e.  $d(A) \neq 0$  and  $d(A^{-1}) = d(A)^{-1}$ . Hence,

$$A \in \text{GL}(n) \iff d(A) \neq 0.$$

We thus have that  $d$  is completely determined by its values on  $R_{ij}$ ,  $E_1(\lambda)$  and  $E_{ij}(c)$ . Note that we have not yet used the continuity of  $d$ , but let us do so now. We can split  $\mathbb{R}^{n \times n}$  into three connected components, namely  $\text{GL}^-(n)$ ,  $\det^{-1}(0)$  and  $\text{GL}^+(n)$ , where the determinant is less than, equal to, and greater than zero, respectively. Since  $E_1(1), E_{ij}(c) \in \text{GL}^+(n)$  and  $E_1(-1), R_{ij} \in \text{GL}^-(n)$ , we have by continuity of  $d$  that

$$\begin{aligned} d(R_{ij}) = +1 &\implies d \text{ is } > 0, = 0, \text{ resp. } > 0 \\ d(R_{ij}) = -1 &\implies d \text{ is } < 0, = 0, \text{ resp. } > 0 \end{aligned} \tag{A.6}$$

on the respective component. Using that  $d(E_1(-1))^2 = d(E_1(-1)^2) = d(I) = 1$ , we have  $d(E_1(-1)) = \pm 1$  and  $d(E_1(-\lambda)) = d(E_1(-1))d(E_1(\lambda)) = \pm d(E_1(\lambda))$ , where the sign depends on (A.6). Note that on  $\mathbb{R}_{>0} := \{\lambda \in \mathbb{R} : \lambda > 0\}$  we have a continuous map  $d \circ E_1 : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$  such that

$$d \circ E_1(\lambda\mu) = d \circ E_1(\lambda) \cdot d \circ E_1(\mu) \quad \forall \lambda, \mu \in \mathbb{R}_{>0}.$$

Forming  $f := \ln \circ d \circ E_1 \circ \exp$ , we then have a continuous map  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(\lambda + \mu) = f(\lambda) + f(\mu).$$

By extending linearity from  $\mathbb{Z}$  to  $\mathbb{Q}$  and then  $\mathbb{R}$  by continuity, we must have that  $f(\lambda) = \alpha\lambda$  for some  $\alpha \in \mathbb{R}$ . Hence,  $d \circ E_1(\lambda) = \lambda^\alpha$  for  $\lambda > 0$ . Continuity also demands that  $\alpha > 0$ .

It only remains to consider  $d \circ E_{ij} : \mathbb{R} \rightarrow \mathbb{R}_{>0}$ . We have  $d \circ E_{ij}(0) = d(I) = 1$  and  $E_{ij}(c)E_{ij}(\gamma) = E_{ij}(c + \gamma)$ , i.e.

$$d \circ E_{ij}(c + \gamma) = d \circ E_{ij}(c) \cdot d \circ E_{ij}(\gamma) \quad \forall c, \gamma \in \mathbb{R}. \tag{A.7}$$

Proceeding as above,  $g := \ln \circ d \circ E_{ij} : \mathbb{R} \rightarrow \mathbb{R}$  is linear and continuous, so that  $g(c) = \alpha_{ij}c$  for some  $\alpha_{ij} \in \mathbb{R}$ , hence  $d \circ E_{ij}(c) = e^{\alpha_{ij}c}$ . One can verify that the following identity holds for all  $i \neq j$ :

$$E_{ji}(-1) = E_j(-1)R_{ij}E_{ji}(1)E_{ij}(-1).$$

This gives  $d(E_{ji}(-1)) = (\pm 1)(\pm 1)d(E_{ji}(1))d(E_{ij}(-1))$  and, using (A.7),

$$\begin{aligned} d(E_{ij}(1)) &= d(E_{ji}(2)) = d(E_{ji}(1+1)) = d(E_{ji}(1))d(E_{ji}(1)) \\ &= d(E_{ij}(2))d(E_{ij}(2)) = d(E_{ij}(4)), \end{aligned}$$

which requires  $\alpha_{ij} = 0$ .

We conclude that  $d$  is completely determined by  $\alpha > 0$ , where  $d \circ E_1(\lambda) = \lambda^\alpha$  and  $\lambda \geq 0$ , plus whether  $d$  takes negative values or not. This proves the theorem.  $\square$

## References

- [1] R. Ablamowicz, G. Sobczyk. *Lectures on Clifford (Geometric) Algebras and Applications*. Birkhäuser Boston, Inc., Boston, MA, 2004.
- [2] J. F. Adams. *Vector fields on spheres*. Ann. of Math. 75 (1962), 603–632.
- [3] E. Arcaute, A. Lasenby, C. J. L. Doran. *Twistors in Geometric Algebra*. Adv. Appl. Clifford Algebr. 18 (2008), 373–394.
- [4] E. Artin. *Geometric algebra*. Interscience, New York, 1957.
- [5] J. C. Baez. *The octonions*. Bull. Amer. Math. Soc. (N.S.) 39 (2002), 145–205.
- [6] I. M. Benn, R. W. Tucker. *An introduction to spinors and geometry with applications in physics*. Adam Hilger, Ltd., Bristol, 1987.
- [7] J. S. R. Chrisolm, A. K. Common (eds.). *Clifford Algebras and their Applications in Mathematical Physics*. D. Reidel Publishing Co., Dordrecht, 1986.
- [8] W. K. Clifford. *Mathematical papers*. Edited by Robert Tucker, with an introduction by H. J. Steven Smith, Chelsea Publishing Co., New York, 1968.
- [9] P. Dadbeh. *Inverse and Determinant in 0 to 5 Dimensional Clifford Algebra*. [arXiv:1104.0067](https://arxiv.org/abs/1104.0067) [math-ph]
- [10] P. de la Harpe. *The Clifford algebra and the spinor group of a Hilbert space*. Compositio Math. 25 (1972), 245–261.
- [11] C. J. L. Doran, D. Hestenes, F. Sommen, N. Van Acker. *Lie groups as spin groups*. J. Math. Phys. 34 (1993), 3642–3669.
- [12] C. J. L. Doran, A. N. Lasenby. *Geometric Algebra for Physicists*. Cambridge University Press, Cambridge, 2003.
- [13] R. Geroch. *Spinor structure of space-times in general relativity. I*. J. Math. Phys. 9 (1968), 1739–1744.
- [14] J. E. Gilbert, M. A. M. Murray. *Clifford algebras and Dirac operators in harmonic analysis*. Cambridge University Press, Cambridge, 1991.
- [15] R. Goodman, N. R. Wallach. *Representations and Invariants of the Classical Groups*. Cambridge University Press, Cambridge, 1998.
- [16] H. G. Grassmann. *A new branch of mathematics*. The Ausdehnungslehre of 1844 and other works. Translated from the German and with a note by Lloyd C. Kannenberg. With a foreword by Albert C. Lewis. Open Court Publishing Co., Chicago, IL, 1995.
- [17] L. C. Grove. *Classical groups and geometric algebra*. American Mathematical Society, Providence, RI, 2002.

- [18] S. F. Gull, A. N. Lasenby, C. J. L. Doran. *Imaginary Numbers are not Real - the Geometric Algebra of Spacetime*. Found. Phys. 23 (1993), 1175–1201.
- [19] I. N. Herstein. *Topics in Algebra, 2nd ed.* John Wiley and Sons, 1975.
- [20] D. Hestenes. *New foundations for classical mechanics, 2nd ed.* Kluwer Academic Publishers Group, Dordrecht, 1999.
- [21] D. Hestenes, G. Sobczyk. *Clifford Algebra to Geometric Calculus*. D. Reidel Publishing Co., Dordrecht, 1984.
- [22] D. Hestenes. *Space-Time Algebra*. Gordon and Breach, New York, 1966.
- [23] L. Hörmander. *Riemannian geometry : lectures given during the fall of 1990*. Matematiska institutionen, Lunds universitet, 1990.
- [24] N. Jacobson. *Lie Algebras*. Dover Publications, Inc., New York, 1979.
- [25] S. Lang. *Algebra, 3rd ed.* Addison-Wesley, 1993.
- [26] A. N. Lasenby, C. J. L. Doran and S. F. Gull. *Gravity, gauge theories and geometric algebra*. R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci. 356 (1998), 487–582.
- [27] B. H. Lawson, Jr., M.-L. Michelsohn. *Spin Geometry*. Princeton University Press, Princeton, NJ, 1989.
- [28] P. Lounesto. *Clifford Algebras and Spinors*. Cambridge University Press, Cambridge, 1997.
- [29] D. Lundholm. *On the Geometry of Supersymmetric Quantum Mechanical Systems*. J. Math. Phys. 49, 062101 (2008).
- [30] D. Lundholm. *Geometric (Clifford) algebra and its applications*. M.Sc. thesis, KTH, Trita-MAT. MA, ISSN 1401-2278; 2006:01, [arXiv:math/0605280](https://arxiv.org/abs/math/0605280)
- [31] D. Lundholm, *Zero-energy states in supersymmetric matrix models*, Ph.D. thesis, KTH, 2010, <http://urn.kb.se/resolve?urn=urn:nbn:se:kth:diva-12846>
- [32] N. G. Marchuk, S. E. Martynova. *Notions of determinant, spectrum, and Hermitian conjugation of Clifford algebra elements*. [arXiv:math-ph/0307043](https://arxiv.org/abs/math-ph/0307043)
- [33] A. Naeve, L. Svensson. *Discrete Integration and Derivation*. Presented at the 5:th International Conference on Clifford Algebras and their Applications in Mathematical Physics, Ixtapa-Zihuatanejo, Mexico, June 27-July 4, 1999. (Online version at <http://kmr.nada.kth.se>)
- [34] M. Nakahara. *Geometry, Topology and Physics, 2nd ed.* Institute of Physics, Bristol, 2003.
- [35] A. Petersson. *Enumeration of spanning trees in simplicial complexes*. Licentiate Thesis, Uppsala Universitet, 2009, U.U.D.M. Report 2009:13.

- [36] R. J. Plymen. *The Laplacian and the Dirac operator in infinitely many variables*. *Compositio Math.* 41 (1980), 137–152.
- [37] I. R. Porteous. *Clifford algebras and the classical groups*. Cambridge University Press, Cambridge, 1995.
- [38] M. Riesz. *Clifford Numbers and Spinors*. University of Maryland, 1958, Kluwer Academic Publishers, 1993.
- [39] G. Schubring (Editor). *Hermann Günther Grassmann (1809-1877) : visionary mathematician, scientist and neohumanist scholar*. Kluwer Acad. Publ., Dordrecht, 1996.
- [40] D. Shirokov. *Concepts of trace, determinant and inverse of Clifford algebra elements*. [arXiv:1108.5447](https://arxiv.org/abs/1108.5447) [math-ph]
- [41] B. Simon. *Representations of finite and compact groups*. American Mathematical Society, Providence, RI, 1996.
- [42] G. Sommer (Editor). *Geometric Computing with Clifford Algebras: Theor. Found. and Appl. in Computer Vision and Robotics*. Springer, Berlin, 2001.
- [43] L. Svensson. *Lecture notes from a course in Clifford Algebra*. KTH, 2000.
- [44] L. Svensson, A. Naeve. *Combinatorial Aspects of Clifford Algebra*. Presented at the International Workshop on Applications of Geometric Algebra, Cambridge, 5-6 Sept. 2002. (Online version at <http://kmr.nada.kth.se>)
- [45] J. Vaz, Jr. and W.A. Rodrigues, Jr. *Equivalence of the Dirac and Maxwell equations and quantum mechanics*. *Int. J. Theor. Phys.* **32**:945-958, 1993.
- [46] J. Vaz, Jr. and W.A. Rodrigues, Jr. *Maxwell and Dirac theories as an already unified theory*. [arXiv:hep-th/9511181](https://arxiv.org/abs/hep-th/9511181) v1, 1995.
- [47] G. P. Wene. *The Clifford algebra of an infinite-dimensional space*. *J. Math. Phys.* 30 (1989), 249–251.

## Index

- $I_{\mathbb{C}}$ , 97
- $V^{\times}$ , 63
- $\Gamma$ , 63
- $\mathcal{B}$ , 26
- $\mathcal{B}^{\bullet}$ , 26
- $\mathcal{B}^{\times}$ , 26
- $Cl(X, R, r)$ , 14
- $Cl_{\mathcal{F}}$ , 50
- $\tilde{\Gamma}$ , 63
- $\mathcal{G}^{\times}$ , 63
- $\mathcal{G}(V, q)$ , 7
- $\mathcal{T}(V)$ , 120
- $k$ -vector, 17
- 2-blade, 1
- 2-vector, 1
- 3-blade, 2
  
- adjoint action, 64
- algebra, 114
  - associative, 114
  - unital, 114
- annihilation operator, 48
- anti-derivation, 25
- anti-euclidean space, 11
- anti-lorentzian space, 11
- anticommutation relations, 6, 48, 98
- anticommutator, 49
- antisymmetry, 114, 115
- arrow, 121
- associative (composition), 112
- automorphism, 115
  
- Betti number, 46
- bilinear form, 6
- binary composition, 112
- bivector, 18
- blade, 26
- boson, 49
- Bott periodicity, 60
- boundary (simplicial complex), 45
- boundary map, 42
  
- canonical basis, 9
- Cartan–Dieudonné’s theorem, 66, 69
- category, 121
- center (of an algebra), 13
- chain, 42
  
- chain group, 42
- chain map, 43
- characteristic, 113
- classification, 54
  - complex, 59
  - real, 57
- Clifford algebra, 14
  - infinite-dimensional, 50
- Clifford conjugate, 20
- Clifford group, 63
- Clifford product, 7, 15
- Clifford, William Kingdon, 1
- codomain, 121
- commutative (composition), 112
- complete triangulation, 44
- complex morphism, 43
- complex number, 12, 13
- complex structure, 99
- composition, 121
- content map, 44
- creation operator, 48
- cross product, 24
- cycle (simplicial complex), 45
  
- degenerate geometric algebra, 12
- degenerate space, 10
- determinant, 32
  - uniqueness, 91, 125
- Dirac algebra, 89
- direct product, 116
- direct sum, 116
- distributive (composition), 112
- domain, 121
- double Cayley algebra, 25
- dual, 24
  - outer product, 24
  - outermorphism, 31
  
- endomorphism, 115
- equivalence relation, 115
- euclidean group, 93
- euclidean space, 11
- even subspace/subalgebra, 17, 55
- exterior algebra, 11
  
- fermion, 49, 98
- fermionic operator, 48, 98

field, 113  
     skew, 113  
 filtered algebra, 18  
 final object, 122  
 Fock space, 49, 98  
 free associative algebra, 117  
  
 geometric algebra, 7  
 geometric product, 7  
 grade involution, 19, 80  
 graded tensor product, 61  
 grading (of the Clifford algebra), 18  
 Gram–Schmidt orthogonalization, 30  
 Grassmann algebra, 11  
 Grassmann, Hermann Günther, 1  
 group, 112  
     abelian, 112  
  
 Hilbert space, 114  
 Hodge  $*$ , 24  
 homology, 45  
 homomorphism  
     group, 115  
     module, 115  
     R-algebra, 115  
 Hurwitz’ theorem, 100  
  
 ideal, 113  
 index map, 44  
 initial object, 122  
 inner product, 20  
     left, 19  
     right, 19  
 inverse, 112  
 invertible elements, 63  
 invertible vectors, 63  
 involution, 19, 20, 80  
 irrep, 95  
 isomorphism, 115  
  
 Jacobi identity, 114  
 Jordan–von Neumann theorem, 6  
  
 Killing form, 80  
 Kirchoff’s matrix tree theorem, 47  
  
 Lagrange identity, 92  
 left action, 64, 98  
 Lie algebra, 114  
 Lie group, 114  
 light-cone, 28  
  
 lightlike vector/blade, 83  
 line segment, 1  
 linear dependence, 22  
 Lipschitz group, 63  
 list, 117  
 lorentzian space, 11  
  
 main involution  
     first, 19  
     second, 20  
 meet, 24  
 Minkowski spacetime, 86  
 module, 113  
 monoid, 112  
 morphism, 121  
 mother algebra, 62  
 multivector, 18  
     simple, 26  
  
 noncommutative polynomial ring, 117  
 nondegenerate geometric algebra, 12,  
     18  
 nondegenerate space, 10, 18  
 norm function, 72, 91  
 normed division algebra, 100  
 null vector/blade, 83  
 null-cone, 28  
 number operator, 49  
  
 object, 121  
 octonion, 100, 101  
 octonionic structure, 101  
 odd subspace, 17  
 order  
     partial, 115  
     total, 115  
 orientation, 12, 15  
 oriented simplex, 42  
 orthogonal basis, 9  
 orthogonal group, 33, 65  
 orthonormal basis, 9  
 outer product, 1, 19  
     dual, 24  
 outermorphism, 30  
     dual, 31  
  
 parallelizable sphere, 102  
 parallelogram identity, 6  
 plane algebra, 11, 12, 82  
 polarization (of quadratic form), 6



- projection, 34
- projection on grade, 19
- projector, 54
- pseudoscalar, 12
  
- quadratic form, 6
- quadratic space, 6
- quaternion, 13, 100
- quotient module, 118
  
- R-algebra, 114
- Radon–Hurwitz’ theorem, 102
- rank (homology), 46
- reciprocal basis, 28
- reflection, 65
- reflexivity, 115
- rejection, 35
- relation, 115
- representation, 95
  - complex, 97
  - fermionic, 98
  - indecomposable, 95
  - irreducible, 95
  - real, 96
  - reducible, 95
- residue class, 118
- reversion, 19
- right action, 64
- ring, 112
  - commutative, 113
  - with unit, 112
- rotor, 5, 63
  
- scalar, 14, 18, 113
- scalar product, 19
- second quantization, 99
- section (function), 116
- signature, 10, 14
- simplex dimension, 42
- simplicial complex, 42
- space algebra, 11, 12, 18, 84
- spacelike vector/blade, 83
- spacetime algebra, 86
- span, 114
- spanning tree, 46
- Sperner’s lemma, 44
- spin field, 11
- spinor, 104
  - Dirac, 107
  - Majorana, 107
  - Majorana–Weyl, 107
  - Weyl, 107
- spinor module, 104
- spinor representation
  - complex, 105
  - real, 104
- spinor space, 104
- state (Fock space), 49
- submodule, 113
- Sylvester’s Law of Inertia, 10
- symmetric difference, 14
- symmetry, 115
  
- tensor algebra, 7, 120
- timelike vector/blade, 83
- torsion (homology), 46
- transitivity, 115
- triviality, 105
- trivector, 2
- twisted adjoint action, 64
  
- unit, 112, 121
- universality, 8, 16, 55, 101, 119, 122
  
- vacuum state, 49
- vector, 1, 18
- vector space, 114
- versor, 63
  - group, 63
  - unit, 63
- volume element, 12
  - complex, 97