



LUDWIG-
MAXIMILIANS-
UNIVERSITÄT
MÜNCHEN

MATHEMATISCHES INSTITUT



Summer term 2020

Prof. D. Kotschick
G. Placini

Mathematical Gauge Theory II

Sheet 2

Exercise 1. (Anti-linear automorphism II) Let $J, J': V \rightarrow V'$ be complex anti-linear isomorphisms between standard Clifford modules. Show that there exists a number $\lambda \in S^1$ so that $J'(\lambda\phi) = \lambda J(\phi)$ for all $\phi \in V$.

Exercise 2. (The group $\text{Spin}(n)$) Let $J_0: \mathbb{C}^N \rightarrow \mathbb{C}^N$ be a complex anti-linear automorphism of the standard Clifford module γ_0 . The group $\text{Spin}(n)$ is defined as the set of pairs $(\tau, \sigma) \in \text{Spin}^c(n)$ so that σ commutes with J_0 . Prove that the homomorphism

$$\begin{aligned} q: \text{Spin}(n) &\longrightarrow \text{SO}(n) \\ (\tau, \sigma) &\longmapsto \tau \end{aligned}$$

is surjective with kernel $\{(I_n, \pm I_N)\} \cong \mathbb{Z}_2$.

Exercise 3. ($\text{Spin}^c(n)$ reconstructed from $\text{Spin}(n)$) We consider the quotient

$$(\text{Spin}(n) \times S^1)/\mathbb{Z}_2,$$

where (τ, σ, λ) gets identified with $(\tau, -\sigma, -\lambda)$. Prove that the homomorphism

$$\begin{aligned} (\text{Spin}(n) \times S^1)/\mathbb{Z}_2 &\longrightarrow \text{Spin}^c(n) \\ [\tau, \sigma, \lambda] &\longmapsto (\tau, \lambda\sigma) \end{aligned}$$

is an isomorphism.

Exercise 4. (Fundamental representation of $\text{SU}(2)$) Show that there exists a fixed matrix $M \in \text{SU}(2)$ such that

$$MAM^\dagger = \bar{A} \quad \forall A \in \text{SU}(2).$$

You can email the solutions until Tuesday, May 5th at noon.