

A theory of computable functionals

Helmut Schwichtenberg

Mathematisches Institut, LMU, München

Gothenburg, 2. September 2022


A theory of computable functionals (TCF)

Similar to HA^ω , but

- add inductively and coinductively defined predicates,
- distinguish computationally relevant (c.r.) and non-computational (n.c.) predicates,
- add realizability predicates (internal “meta”-step),
- allow partial functionals, defined by equations (possibly non-terminating, like corecursion),
- minimal logic: only \rightarrow , \forall primitive. \vee , \exists , \wedge inductively defined.

Minlog implements TCF.

- TCF has an intended model: **partial continuous functionals**.
- Defined via **information systems** (Scott). Has function spaces.
- It consists of **ideals** (infinite) approximated by **tokens** (finite).
- Ideals are **consistent** and **deductively closed** sets of tokens.
- Tokens are constructor trees with possibly $*$ at some leaves.
- Examples: natural numbers \mathbb{N} , binary trees \mathbb{Y} .



- $\{S_0, S(S^*)\}$ is inconsistent.
- $\{S^*, S(S^*)\}$ is an ideal.
- $\{S^*, S(S^*), S(S_0)\}$ is an ideal ("total").
- $\{S^*, S(S^*), S(S(S^*)), \dots\}$ is an infinite ideal ("cototal").


An ideal x in a closed base type

- is **cototal** if for each of its tokens $t(*)$ with a distinguished occurrence of $*$ there is another token of the form $t(C\vec{*})$ in x ,
- **total** if it is cototal and finite.


The base type \mathbb{Y} (binary trees) is given by the constructors

$$\begin{aligned} - &: \mathbb{Y} && \text{(leaf),} \\ C &: \mathbb{Y} \rightarrow \mathbb{Y} \rightarrow \mathbb{Y} && \text{(branch).} \end{aligned}$$


Example of a cototal ideal in \mathbb{Y}



Example of a total ideal in \mathbb{Y} : deductive closure of



Example of a neither total nor cotal ideal: deductive closure of



Totality $T_{\mathbb{N}}$ is inductively defined as the least fixed point (lfp) of the clauses

$$0 \in T_{\mathbb{N}}, \quad n \in T_{\mathbb{N}} \rightarrow Sn \in T_{\mathbb{N}}.$$

Cototality ${}^{\text{co}}T_{\mathbb{N}}$ is coinductively defined as the greatest fixed point (gfp) of its closure axiom

$$n \in {}^{\text{co}}T_{\mathbb{N}} \rightarrow n \equiv 0 \vee \exists_{n'}(n' \in {}^{\text{co}}T_{\mathbb{N}} \wedge n \equiv Sn').$$

Similarity \sim_Y is a binary variant of totality. It is inductively defined as the least fixed point (lfp) of the clauses

$$\begin{aligned} & - \sim_Y -, \\ & t_1 \sim_Y t'_1 \rightarrow t_2 \sim_Y t'_2 \rightarrow Ct_1 t_2 \sim_Y Ct'_1 t'_2. \end{aligned}$$

Bisimilarity \approx_Y is a binary variant of cototality. It is coinductively defined as the greatest fixed point (gfp) of its closure axiom

$$\begin{aligned} t \approx_Y t' & \rightarrow ((t \equiv -) \wedge (t' \equiv -)) \vee \\ & \exists_{t_1, t_2, t'_1, t'_2} (t_1 \approx_Y t'_1 \wedge t_2 \approx_Y t'_2 \wedge t \equiv Ct_1 t_2 \wedge t' \equiv Ct'_1 t'_2) \end{aligned}$$

Lemma

For every closed base type bisimilarity implies Leibniz equality.

- Example: Υ . Let a range over tokens, t over ideals.
- By induction on the height of extended tokens a^* we prove

$$\forall_{a^*, t, t'} (t \approx_{\Upsilon} t' \rightarrow a^* \in t \rightarrow a^* \in t').$$

- It suffices to consider the case $Ca_1^*a_2^*$.
- From $t \approx t'$ by closure we have ideals t_1, t_2, t'_1, t'_2 with

$$t_1 \approx t'_1 \wedge t_2 \approx t'_2 \wedge t \equiv Ct_1t_2 \wedge t' \equiv Ct'_1t'_2.$$

- Then $a_i^* \in t_i$, and by IH $a_i^* \in t'_i$. Thus $Ca_1^*a_2^* \in t'$.

Axioms for (co)inductive predicates I^\pm , $\text{co}I^\pm$. Examples:

- Even. The introduction axioms (or clauses) are $\text{Even}_{0,1}^+$:

$$0 \in \text{Even}, \quad n \in \text{Even} \rightarrow S(Sn) \in \text{Even}$$

and the elimination axiom is Even^- :

$$0 \in X \rightarrow \forall_n (n \in \text{Even} \rightarrow n \in X \rightarrow S(Sn) \in X) \rightarrow \text{Even} \subseteq X.$$

“Every competitor X satisfying the clauses is above X .”

- Similar: T_l^\pm , $\text{co}T_l^\pm$, \sim_l^\pm and \approx_l^\pm
- The n.c. **Leibniz equality** \equiv is defined by

$$\equiv^+ : x^T \equiv x^T \quad \equiv^- : x \equiv y \rightarrow \forall_x Xxx \rightarrow Xxy$$

We can deduce the property Leibniz used as a definition.

Lemma (Compatibility of EqD)

$$x \equiv y \rightarrow A(x) \rightarrow A(y).$$

Proof: By the elimination axiom with

$$X := \{ x, y \mid A(x) \rightarrow A(y) \}.$$

Using compatibility of \equiv one proves symmetry and transitivity.

Define **falsity** by $\mathbf{F} := (\text{ff} \equiv \text{tt})$.

Theorem (Ex-falso-quodlibet)

We can derive $\mathbf{F} \rightarrow A$ from assumptions $\text{Ef}_Y : \forall \vec{x} (\mathbf{F} \rightarrow Y \vec{x})$ for predicate variables Y strictly positive in A , and $\text{Ef}_I : \forall \vec{x} (\mathbf{F} \rightarrow I \vec{x})$ for inductive predicates I without a nullary clause.

Bisimilarity axioms:

For every closed base type bisimilarity implies Leibniz equality.

Justification: holds in the intended model.

For closed base types ι it follows that

$$\begin{aligned}t \sim_{\iota} t' &\leftrightarrow t, t' \in T_{\iota} \wedge t \equiv t', \\t \approx_{\iota} t' &\leftrightarrow t, t' \in {}^{\text{co}}T_{\iota} \wedge t \equiv t' .\end{aligned}$$

This is helpful because it gives us a tool (induction, coinduction) to prove equalities $t \equiv t'$, which otherwise would be difficult.

Corollary

$$t \sim_\iota t \leftrightarrow t \in T_\iota,$$

$$t \approx_\iota t \leftrightarrow t \in {}^{\text{co}}T_\iota,$$

\sim_ι is an equivalence relation on T_ι ,

\approx_ι is an equivalence relation on ${}^{\text{co}}T_\iota$.

Definition (Pointwise equality¹)

$$(x \dot{=}_{\iota} y) := (x \approx_{\iota} y)$$
$$(f \dot{=}_{\tau \rightarrow \sigma} g) := \forall_{x,y} (x \dot{=}_{\tau} y \rightarrow fx \dot{=}_{\sigma} gy).$$

Definition (Extensionality)

$$(x \in \text{Ext}_{\tau}) := (x \dot{=}_{\tau} x).$$

¹Robin Gandy, On the axiom of extensionality – Part I, JSL 1956 and Gaisi Takeuti, On a generalized logic calculus, Jap. J. Math. 1953

Example of a non-extensional functional:

- Define f, g of type $\mathbb{N} \rightarrow \mathbb{N}$ by the computation rules $fn = 0$ and $g0 = 0, g(Sn) = gn$.
- Then $f \perp_{\mathbb{N}} = 0$ by the computation rules for f .
- For $g \perp_{\mathbb{N}}$ no computation rule fits, but by the definition of $[[\lambda_{\bar{x}} M]]$ we have that $[[g \perp_{\mathbb{N}}]]$ is the empty ideal $[[\perp_{\mathbb{N}}]]$.
- Hence $f \dot{=} g$, i.e., $\forall_{n,m}(n \dot{=}_{\mathbb{N}} m \rightarrow fn \dot{=}_{\mathbb{N}} gm)$, since $n \dot{=}_{\mathbb{N}} m$ implies $n \in T_{\mathbb{N}}$ and $n \equiv m$.
- Therefore the functional F defined by $Fh = h \perp_{\mathbb{N}}$ maps the pointwise equal f, g to different values.

Lemma

Ext_τ and ${}^{\text{co}}T_\tau$ are equivalent for closed types of level ≤ 1 .

Proof.

For closed base types this has been proved above. In case of level 1 we use induction on the height of the type. Let $\tau \rightarrow \sigma$ be a closed type of level 1. The following are equivalent.

$$f \in \text{Ext}_{\tau \rightarrow \sigma}$$

$$f \dot{=}_{\tau \rightarrow \sigma} f$$

$$\forall_{x,y} (x \dot{=}_\tau y \rightarrow fx \dot{=}_\sigma fy)$$

$$\forall_{x \in {}^{\text{co}}T_\tau} (fx \dot{=}_\sigma fx) \quad \text{by the Corollary, since } \text{lev}(\tau) = 0$$

$$\forall_{x \in {}^{\text{co}}T_\tau} (fx \in \text{Ext}_\sigma).$$

By IH the final formula is equivalent to $f \in {}^{\text{co}}T_{\tau \rightarrow \sigma}$. □

For arbitrary closed types the relation $\dot{=}_{\tau}$ is a “partial equivalence relation”, which means the following.

Lemma

For every closed type τ the relation $\dot{=}_{\tau}$ is an equivalence relation on Ext_{τ} .

Lemma (Compatibility of terms)

For every term $t(\vec{x})$ with extensional constants and free variables among \vec{x} we have

$$\vec{x} \dot{=}_{\vec{\rho}} \vec{y} \rightarrow t(\vec{x}) \dot{=}_{\tau} t(\vec{y}).$$

Lemma (Extensionality of terms)

For every term $t(\vec{x})$ with extensional constants and free variables among \vec{x} we have

$$\vec{x} \in \text{Ext}_{\vec{\rho}} \rightarrow t(\vec{x}) \in \text{Ext}_{\tau}.$$

Need “realizability extensions” of c.r. predicates and formulas:

- Assume that we have a global assignment giving for every c.r. predicate variable X of arity $\vec{\rho}$ an n.c. predicate variable X^r of arity $(\vec{\rho}, \xi)$ where ξ is the type variable associated with X .
- We introduce $I^r / {}^{co}I^r$ for c.r. (co)inductive predicates $I / {}^{co}I$, e.g.,

$$\text{Even}^r 00 \quad \text{Even}^r nm \rightarrow \text{Even}^r (S(Sn))(Sm).$$

- A predicate or formula C is **r-free** if it does not contain any of these X^r , I^r or ${}^{co}I^r$.
- A derivation M is **r-free** if it contains **r-free** formulas only.

Definition (C^r for r -free c.r. formulas C)

Let $z \mathbf{r} C$ mean $C^r z$.

$$z \mathbf{r} P \vec{t} := P^r \vec{t} z,$$

$$z \mathbf{r} (A \rightarrow B) := \begin{cases} \forall_w (w \mathbf{r} A \rightarrow z w \mathbf{r} B) & \text{if } A \text{ is c.r.} \\ A \rightarrow z \mathbf{r} B & \text{if } A \text{ is n.c.} \end{cases}$$

$$z \mathbf{r} \forall_x A := \forall_x (z \mathbf{r} A).$$

Definition (Extracted term for an \mathbf{r} -free proof M of a c.r. A)

$$\text{et}(u^A) \quad := z_u^{\tau(A)} \quad (z_u^{\tau(A)} \text{ uniquely associated to } u^A),$$

$$\text{et}((\lambda_{u^A} M^B)^{A \rightarrow B}) := \begin{cases} \lambda_{z_u} \text{et}(M) & \text{if } A \text{ is c.r.} \\ \text{et}(M) & \text{if } A \text{ is n.c.} \end{cases}$$

$$\text{et}((M^{A \rightarrow B} N^A)^B) := \begin{cases} \text{et}(M)\text{et}(N) & \text{if } A \text{ is c.r.} \\ \text{et}(M) & \text{if } A \text{ is n.c.} \end{cases}$$

$$\text{et}((\lambda_x M^A)^{\forall_x A}) := \text{et}(M),$$

$$\text{et}((M^{\forall_x A(x)} t)^{A(t)}) := \text{et}(M).$$

It remains to define extracted terms for the **axioms**. Consider a (c.r.) inductively defined predicate I .

- $\text{et}(I_i^+) := C_i$ and $\text{et}(I^-) := \mathcal{R}$, where the **constructor** C_i and the **recursion** operator \mathcal{R} refer to ι_I associated with I .
- $\text{et}({}^{\text{co}}I^-) := D$ and $\text{et}({}^{\text{co}}I_i^+) := {}^{\text{co}}\mathcal{R}$, where the **destructor** D and the **corecursion** operator ${}^{\text{co}}\mathcal{R}$ refer to the ι_I .

Let I be an inductive predicate and ι_I its associated algebra. One can show that

- every constructor of ι_I is extensional w.r.t. its clause I_i^+ ,
- $\mathcal{R}_{\iota_I}^\alpha$ is extensional w.r.t. the least-fixed-point axiom I^- ,
- the destructor of ι_I is extensional w.r.t. the closure axiom $\text{co}I^-$, and
- $\text{co}\mathcal{R}_{\iota_I}^\alpha$ is extensional w.r.t. the greatest-fixed-point axiom $\text{co}I^+$.

Since the term $\text{et}(M)$ extracted from a closed proof M of a c.r. formula A is built from these constants by abstraction and application, by the lemma on extensionality of terms we can conclude that $\text{et}(M)$ is extensional w.r.t. A .

Theorem (Soundness)

Let M be an \mathbf{r} -free derivation of a formula A from assumptions $u_i: C_i$ ($i < n$). Then we can derive

$$\begin{cases} \text{et}(M) \mathbf{r} A & \text{if } A \text{ is c.r.} \\ A & \text{if } A \text{ is n.c.} \end{cases}$$

from assumptions

$$\begin{cases} z_{u_i} \mathbf{r} C_i & \text{if } C_i \text{ is c.r.} \\ C_i & \text{if } C_i \text{ is n.c.} \end{cases}$$

We express

- Kolmogorov's view of "formulas as problems"²
- Feferman's dictum "to assert is to realize"³

by **invariance axioms**:

For **r**-free c.r. formulas A we require as axioms

$$\text{InvAll}_A: \forall_z (z \mathbf{r} A \rightarrow A).$$

$$\text{InvEx}_A: A \rightarrow \exists_z (z \mathbf{r} A).$$

²Zur Deutung der intuitionistischen Logik, Math. Zeitschr., 1932

³Constructive theories of functions and classes, Logic Colloquium 78, p.208

Invariance axioms used in the proof of soundness (1):

Case $(\lambda_{u^A} M^B)^{A \rightarrow B}$ with B n.c. We need a derivation of $A \rightarrow B$.

Subcase A c.r. By IH we have a derivation of B from $z \mathbf{r} A$. Using the invariance axiom $A \rightarrow \exists_z(z \mathbf{r} A)$ we get the required derivation of B from A :

$$\frac{\frac{A \rightarrow \exists_z(z \mathbf{r} A) \quad A}{\exists_z(z \mathbf{r} A)}}{\frac{B}{\exists^-}} \quad \begin{array}{l} [z \mathbf{r} A] \\ | \text{IH} \\ B \end{array}$$

Invariance axioms used in the proof of soundness (2):

Case $(M^{A \rightarrow B} N^A)^B$ with B n.c. Goal: find a derivation of B .

Subcase A c.r. By IH we have derivations of $A \rightarrow B$ and of $\text{et}(N) \mathbf{r} A$. From the invariance axiom $\forall_z (z \mathbf{r} A \rightarrow A)$ we obtain the required derivation of B by \rightarrow^- from the derivation of $A \rightarrow B$ and

$$\frac{\frac{\forall_z (z \mathbf{r} A \rightarrow A) \quad \text{et}(N)}{\text{et}(N) \mathbf{r} A \rightarrow A} \quad \text{IH}}{\text{et}(N) \mathbf{r} A} \quad | \text{IH}}{A}$$

Conclusion

- In TCF the computational content of a proof M is represented by an extracted term $\text{et}(M)$ **in the language of TCF**.
- The Soundness theorem provides a formal verification in TCF that the extracted term realizes the formula (“specification”). This is automated in Minlog.
- Since extraction ignores n.c. parts of the proof, $\text{et}(M)$ is much shorter than M .
- For efficiency, in a second step one can translate the extracted term to a functional programming language. Minlog does this for Scheme and Haskell.

Question: TCF has an internal “meta”-theory, via realizability.
Any relation to Truth theories?