



Prof. T. Ø. SØRENSEN PhD  
T. König

Winter term 2016/17  
December 22, 2016

## FUNCTIONAL ANALYSIS II

### ASSIGNMENT 10

**Problem 37.** (*Modulus, positive and negative part of an operator*)

Let  $\mathcal{H}$  be a Hilbert space and let  $A \in \mathcal{B}(\mathcal{H})$  be self-adjoint. Prove:

- (a) The operator  $|A| = \sqrt{A^*A}$  constructed with Hilbert space techniques (see Problem 18) coincides with  $|A|$  defined via the functional calculus.
- (b)  $A \leq |A|$ .
- (c) There exists a unique pair  $A_+, A_- \in \mathcal{B}(\mathcal{H})$  of self-adjoint operators such that

$$A_+, A_- \geq 0, \quad A_+ A_- = 0, \quad A = A_+ - A_-.$$

**Problem 38.** (*A class of operator monotone functions*)

A continuous real-valued function  $f$  on an interval  $I \subset \mathbb{R}$  is said to be *operator monotone* (on the interval  $I$ ), if  $A \leq B$  implies that  $f(A) \leq f(B)$  for all self-adjoint operators  $A, B \in \mathcal{B}(\mathcal{H})$  such that  $\sigma(A) \subset I$  and  $\sigma(B) \subset I$ .

- (a) Prove that  $f_\alpha$  given by  $f_\alpha(t) := \frac{t}{1+\alpha t}$  is operator monotone on  $[0, \infty)$  if  $\alpha \geq 0$ .
- (b) Let  $\alpha \in [0, 1]$ , and  $A, B \in \mathcal{B}(\mathcal{H})$  be such that  $0 \leq A \leq B$ . Prove that  $0 \leq A^\alpha \leq B^\alpha$ , i.e. that  $x \mapsto x^\alpha$  is operator monotone on  $\mathbb{R}_+$ . [Hint: You may use without proof that for all  $x \geq 0, \alpha \in (0, 1)$  we have  $x^\alpha = \frac{\sin(\alpha\pi)}{\pi} \int_0^\infty \frac{x}{x+\lambda} \frac{d\lambda}{\lambda^{1-\alpha}}.$ ]
- (c) Find a counterexample for (b) when  $\alpha > 1$ .

**Problem 39.** (*Functional calculus on a multiplication operator*)

Let  $A : L^2([0, 1]) \rightarrow L^2([0, 1])$  be given by  $Af(x) := xf(x)$  for a.e.  $x \in [0, 1]$ .

- (a) Prove that  $A = A^*$ ,  $\|A\| = 1$  and  $\sigma(A) = [0, 1]$ .
- (b) Give the explicit action of  $f(A) \in \mathcal{B}(L^2([0, 1]))$  for any bounded measurable function  $f : [0, 1] \rightarrow \mathbb{C}$ .
- (c) For any  $\psi \in L^2([0, 1])$  express  $(\psi, f(A)\psi)$  as an integral with respect to the measure  $\Omega \mapsto (\psi, E_\Omega \psi)$ , where  $E$  denotes the projection-valued measure given by  $A$ .

**Problem 40.** (*Spectral projections*)

Let  $\mathcal{H}$  be a Hilbert space, let  $A \in \mathcal{B}(\mathcal{H})$  be self-adjoint, and let  $E$  denote the projection-valued measure given by  $A$ . Prove:

- (a) For any Borel set  $\Omega \subset \sigma(A)$ , the subspace  $R(E_\Omega)$  is invariant under  $A$ .
- (b) If  $\Omega \subset \sigma(A)$  is closed, then  $\sigma(A|_{R(E_\Omega)}) \subset \Omega$ .

*This sheet is to be discussed in the exercise class on Thursday, January 12.  
For more details please visit <http://www.math.lmu.de/~tkoenig/16FA2exercises.php>*