

CHAPTER 8

Toolbox

8. Representable Functors

Definition 8.8.1. Let $\mathcal{F} : \mathcal{C} \rightarrow \mathbf{Set}$ be a covariant functor. A pair (A, x) with $A \in \mathcal{C}, x \in \mathcal{F}(A)$ is called a *representing (generic, universal) object* for $\mathcal{F}$ and $\mathcal{F}$ is called a *representable functor*, if for each $B \in \mathcal{C}$ and $y \in \mathcal{F}(B)$ there exists a unique $f \in \text{Mor}_{\mathcal{C}}(A, B)$ such that $\mathcal{F}(f)(x) = y$:

$$\begin{array}{ccc} A & & \mathcal{F}(A) \ni x \\ \downarrow f & & \downarrow \mathcal{F}(f) \quad \downarrow \\ B & & \mathcal{F}(B) \ni y \end{array}$$

Proposition 8.8.2. Let (A, x) and (B, y) be representing objects for $\mathcal{F}$. Then there exists a unique isomorphism $f : A \rightarrow B$ such that $\mathcal{F}(f)(x) = y$.

$$\begin{array}{ccccc} & A & & \mathcal{F}(A) & & x \\ & \swarrow & \downarrow h & \swarrow & \downarrow \mathcal{F}(h) & \downarrow \\ 1_A & & B & & \mathcal{F}(B) & \downarrow 1_{\mathcal{F}(A)} & y \\ & \swarrow & \downarrow k & \swarrow & \downarrow \mathcal{F}(k) & \downarrow & \\ 1_B & & A & & \mathcal{F}(A) & \downarrow 1_{\mathcal{F}(B)} & x \\ & \swarrow & \downarrow h & \swarrow & \downarrow \mathcal{F}(h) & \downarrow & y \\ & & B & & \mathcal{F}(B) & & \end{array}$$

Examples 8.8.3. 1. Let $X \in \mathbf{Set}$ and let R be a ring. $\mathcal{F} : R\text{-Mod} \rightarrow \mathbf{Set}$, $\mathcal{F}(M) := \text{Map}(X, M)$ is a covariant functor. A representing object for $\mathcal{F}$ is given by $(RX, x : X \rightarrow RX)$ with the property, that for all $(M, y : X \rightarrow M)$ there exists a unique $f \in \text{Hom}_R(RX, M)$ such that $\mathcal{F}(f)(x) = \text{Map}(X, f)(x) = fx = y$

$$\begin{array}{ccc} X & \xrightarrow{x} & RX \\ & \searrow y & \downarrow f \\ & & M. \end{array}$$

2. Given modules M_R and ${}_R N$. Define $\mathcal{F} : \mathbf{Ab} \rightarrow \mathbf{Set}$ by $\mathcal{F}(A) := \text{Bil}_R(M, N; A)$. Then $\mathcal{F}$ is a covariant functor. A representing object for $\mathcal{F}$ is given by $(M \otimes_R N, \otimes : M \times N \rightarrow M \otimes_R N)$ with the property that for all $(A, f : M \times N \rightarrow A)$ there exists

a unique $g \in \text{Hom}(M \otimes_R N, A)$ such that $\mathcal{F}(g)(\otimes) = \text{Bil}_R(M, N; g)(\otimes) = g \otimes = f$

$$\begin{array}{ccc} M \times N & \xrightarrow{\otimes} & M \otimes_R N \\ & \searrow f & \downarrow g \\ & & A. \end{array}$$

3. Given a $\mathbb{K}$ -module V . Define $\mathcal{F} : \mathbf{Alg} \rightarrow \mathbf{Set}$ by $\mathcal{F}(A) := \text{Hom}(V, A)$. Then $\mathcal{F}$ is a covariant functor. A representing object for $\mathcal{F}$ is given by $(T(V), \iota : V \rightarrow T(V))$ with the property that for all $(A, f : V \rightarrow A)$ there exists a unique $g \in \text{Mor}_{\mathbf{Alg}}(T(V), A)$ such that $\mathcal{F}(g)(\iota) = \text{Hom}(V, g)(\iota) = g\iota = f$

$$\begin{array}{ccc} V & \xrightarrow{\iota} & T(V) \\ & \searrow f & \downarrow g \\ & & A. \end{array}$$

4. Given a $\mathbb{K}$ -module V . Define $\mathcal{F} : \mathbf{cAlg} \rightarrow \mathbf{Set}$ by $\mathcal{F}(A) := \text{Hom}(V, A)$. Then $\mathcal{F}$ is a covariant functor. A representing object for $\mathcal{F}$ is given by $(S(V), \iota : V \rightarrow S(V))$ with the property that for all $(A, f : V \rightarrow A)$ there exists a unique $g \in \text{Mor}_{\mathbf{Alg}}(S(V), A)$ such that $\mathcal{F}(g)(\iota) = \text{Hom}(V, g)(\iota) = g\iota = f$

$$\begin{array}{ccc} V & \xrightarrow{\iota} & S(V) \\ & \searrow f & \downarrow g \\ & & A. \end{array}$$

Proposition 8.8.4. $\mathcal{F}$ has a representing object (A, a) if and only if there is a natural isomorphism $\varphi : \mathcal{F} \cong \text{Mor}_{\mathcal{C}}(A, -)$ (with $a = \varphi(A)^{-1}(1_A)$).

PROOF. $\Rightarrow$: The map

$$\varphi(B) : \mathcal{F}(B) \ni y \mapsto f \in \text{Mor}_{\mathcal{C}}(A, B) \text{ with } \mathcal{F}(f)(a) = y$$

is bijective with the inverse map

$$\psi(B) : \text{Mor}_{\mathcal{C}}(A, B) \ni f \mapsto \mathcal{F}(f)(a) \in \mathcal{F}(B).$$

In fact we have $y \mapsto f \mapsto \mathcal{F}(f)(a) = y$ and $f \mapsto y := \mathcal{F}(f)(a) \mapsto g : \mathcal{F}(g)(a) = y = \mathcal{F}(f)(a)$. By uniqueness we get $f = g$. Hence all $\varphi(B)$ are bijective with inverse map $\psi(B)$. It is sufficient to show that ψ is a natural transformation.

Given $g : B \rightarrow C$. Then the following diagram commutes

$$\begin{array}{ccc} \text{Mor}_{\mathcal{C}}(A, B) & \xrightarrow{\psi(B)} & \mathcal{F}(B) \\ \text{Mor}_{\mathcal{C}}(A, g) \downarrow & & \downarrow \mathcal{F}(g) \\ \text{Mor}_{\mathcal{C}}(A, C) & \xrightarrow{\psi(C)} & \mathcal{F}(C) \end{array}$$

since $\psi(C)\text{Mor}_{\mathcal{C}}(A, g)(f) = \psi(C)(gf) = \mathcal{F}(gf)(a) = \mathcal{F}(g)\mathcal{F}(f)(a) = \mathcal{F}(g)\psi(B)(f)$.

$\Leftarrow$: Let A be given. Let $a := \varphi(A)^{-1}(1_A)$. For $y \in \mathcal{F}(B)$ we get $y = \varphi(B)^{-1}(f) = \varphi(B)^{-1}(f1_A) = \varphi(B)^{-1}\text{Mor}_{\mathcal{C}}(A, f)(1_A) = \mathcal{F}(f)\varphi(A)^{-1}(1_A) = \mathcal{F}(f)(a)$ for a uniquely determined $f \in \text{Mor}_{\mathcal{C}}(A, B)$. $\square$

Proposition 8.8.5. *Given a representable functor $\mathcal{F}_X : \mathcal{C} \rightarrow \mathbf{Set}$ for each $X \in \mathcal{D}$. Given a natural transformation $\mathcal{F}_g : \mathcal{F}_Y \rightarrow \mathcal{F}_X$ for each $g : X \rightarrow Y$ (contravariant!) such that $\mathcal{F}$ depends functorially on X , i.e. $\mathcal{F}_{1_X} = 1_{\mathcal{F}_X}$, $\mathcal{F}_{hg} = \mathcal{F}_g\mathcal{F}_h$. Then the representing objects (A_X, a_X) for $\mathcal{F}_X$ depend functorially on X , i.e. for each $g : X \rightarrow Y$ there is a unique homomorphism $A_g : A_X \rightarrow A_Y$ (with $\mathcal{F}_X(A_g)(a_X) = \mathcal{F}_g(A_Y)(a_Y)$) and the following identities hold $A_{1_X} = 1_{A_X}$, $A_{hg} = A_h A_g$.*

PROOF. Choose a representing object (A_X, a_X) for $\mathcal{F}_X$ for each $X \in \mathcal{C}$ (by the axiom of choice). Then there is a unique homomorphism $A_g : A_X \rightarrow A_Y$ with

$$\mathcal{F}_X(A_g)(a_X) = \mathcal{F}_g(A_Y)(a_Y) \in \mathcal{F}_X(A_Y),$$

for each $g : X \rightarrow Y$ because $\mathcal{F}_g(A_Y) : \mathcal{F}_Y(A_Y) \rightarrow \mathcal{F}_X(A_Y)$ is given. We have $\mathcal{F}_X(A_1)(a_X) = \mathcal{F}_1(A_X)(a_X) = a_X = \mathcal{F}_X(1)(a_X)$ hence $A_1 = 1$, and $\mathcal{F}_X(A_{hg})(a_X) = \mathcal{F}_{hg}(A_Z)(a_Z) = \mathcal{F}_g(A_Z)\mathcal{F}_h(A_Z)(a_Z) = \mathcal{F}_g(A_Z)\mathcal{F}_Y(A_h)(a_Y) = \mathcal{F}_X(A_h)\mathcal{F}_g(A_Y)(a_Y) = \mathcal{F}_X(A_h)\mathcal{F}_X(A_g)(a_X) = \mathcal{F}_X(A_h A_g)(a_X)$ hence $A_h A_g = A_{hg}$ for $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ in $\mathcal{D}$. $\square$

Corollary 8.8.6. 1. $\text{Map}(X, M) \cong \text{Hom}_R(RX, M)$ is a natural transformation in M (and in X !). In particular $\mathbf{Set} \ni X \mapsto RX \in R\text{-Mod}$ is a functor.

2. $\text{Bil}_R(M, N; A) \cong \text{Hom}(M \otimes_R N, A)$ is a natural transformation in A (and in $(M, N) \in \mathbf{Mod}\text{-}R \times R\text{-Mod}$). In particular $\mathbf{Mod}\text{-}R \times R\text{-Mod} \ni M, N \mapsto M \otimes_r N \in \mathbf{Ab}$ is a functor.

3. $R\text{-Mod}\text{-}S \times S\text{-Mod}\text{-}T \ni (M, N) \mapsto M \otimes_S N \in R\text{-Mod}\text{-}T$ is a functor.