



Prof. T. Ø. SØRENSEN PhD  
S. Gottwald

Winter term 2014/15  
October 30, 2014

## FUNCTIONAL ANALYSIS II

### ASSIGNMENT 4

**Problem 13** (Spectrum of the product). Let  $X$  be a Banach space and  $S, T \in \mathcal{B}(X)$ .

- (i) Prove that  $\sigma(TS) \cup \{0\} = \sigma(ST) \cup \{0\}$ .
- (ii) Show that  $\sigma(TS) = \sigma(ST)$  is not true in general.

**Problem 14** (Spectrum of self-adjoint operators). Let  $A$  be a bounded self-adjoint operator on a Hilbert space  $\mathcal{H}$ , i.e.  $A^* = A$ . Prove the following:

- (i)  $\sigma(A) \subset \left[ \inf_{x \in \mathcal{H}, \|x\|=1} \langle x, Ax \rangle, \sup_{x \in \mathcal{H}, \|x\|=1} \langle x, Ax \rangle \right] \subset \mathbb{R}$ .
- (ii)  $\sigma_r(A) = \emptyset$ .
- (iii) If  $x, y \in \mathcal{H}$  and  $\lambda \neq \mu$  are such that  $Ax = \lambda x$  and  $Ay = \mu y$  then  $\langle x, y \rangle = 0$ .
- (iv) If  $\sigma(A) = \{0\}$  then  $A = \mathbb{O}$ .

**Problem 15** (Weyl sequences). Let  $X$  be a Banach space and  $T \in \mathcal{B}(X)$ . A sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  is called a *Weyl sequence* of  $T$  at  $\lambda \in \mathbb{C}$ , if  $\|x_n\| = 1$  for all  $n \in \mathbb{N}$  and  $\|Tx_n - \lambda x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . Prove:

- (i) If  $T$  has a Weyl sequence at  $\lambda \in \mathbb{C}$  then  $\lambda \in \sigma(T)$ .
- (ii) If  $\lambda \in \partial\sigma(T)$  then  $T$  has a Weyl sequence at  $\lambda \in \mathbb{C}$ .

Now, let  $\mathcal{H}$  be a Hilbert space and let  $T \in \mathcal{B}(\mathcal{H})$  be self-adjoint.

- (iii) Prove that  $T$  has a Weyl sequence at  $\lambda$  iff  $\lambda \in \sigma(T)$ .

**Problem 16** (Multiplication operators II). Let  $(X, \mu)$  be a  $\sigma$ -finite measure space, let  $1 \leq p < \infty$ , and for a measurable function  $h : X \rightarrow \mathbb{C}$  let

$$\Omega_h := \{f \in L^p(X, \mu) : hf \in L^p(X, \mu)\}.$$

Let  $M_h : \Omega_h \rightarrow L^p(X, \mu)$ ,  $f \mapsto hf$ .

- (i) Prove that  $M_h \in \mathcal{B}(L^p(X, \mu))$  iff  $h \in L^\infty(X, \mu)$ .

Assuming  $h \in L^\infty(X, \mu)$  prove the following:

- (ii)  $\sigma_p(M_h) = \{\lambda \in \mathbb{C} : \mu(\{x \in X : h(x) = \lambda\}) > 0\}$ .
- (iii)  $\rho(M_h) = \{\lambda \in \mathbb{C} : \exists c > 0 \text{ such that } |\lambda - h(x)| \geq c \text{ a.e.}\}$ .

For more details please visit <http://www.math.lmu.de/~gottwald/14FA2/>