



Prof. T. Ø. SØRENSEN PhD
S. Gottwald

Winter term 2014/15
Dec 11, 2014

FUNCTIONAL ANALYSIS II

ASSIGNMENT 10

Problem 37. Let \mathcal{H} be a Hilbert space and let $A \in \mathcal{B}(\mathcal{H})$ be self-adjoint.

- (i) Show that the operator $|A| = \sqrt{A^*A}$ constructed with Hilbert space techniques (see Problem 18) coincides with $|A|$ defined via the functional calculus.
- (ii) Prove that the operator norm limit of a convergent sequence of positive definite operators on \mathcal{H} is positive definite.
- (iii) Prove that $A_n := 2(\frac{4}{n}\mathbb{I} + (|A| - A)^2)^{-1}(|A| - A)^2|A| \in \mathcal{B}(\mathcal{H})$, $A_n \geq \mathbb{O}$ for all $n \in \mathbb{N}$, and that $A_n \xrightarrow{n \rightarrow \infty} |A| - A$ in $\mathcal{B}(\mathcal{H})$ (hence concluding that $A \leq |A|$). [Hint: No functional calculus argument is needed here.]
- (iv) Reprove that $A \leq |A|$ by using the functional calculus.
- (v) Prove that there exists a unique pair $A_+, A_- \in \mathcal{B}(\mathcal{H})$ of self-adjoint operators such that

$$A_+, A_- \geq \mathbb{O}, \quad A_+ A_- = \mathbb{O}, \quad A = A_+ - A_-.$$

Problem 38 (Operator monotone functions). A continuous real-valued function f on an interval I is said to be *operator monotone* (on the interval I), if $A \leq B$ implies that $f(A) \leq f(B)$ for all self-adjoint operators $A, B \in \mathcal{B}(\mathcal{H})$ such that $\sigma(A) \subset I$ and $\sigma(B) \subset I$.

- (i) Prove that f_α given by $f_\alpha(t) := \frac{t}{1+\alpha t}$ is operator monotone on \mathbb{R}_+ if $\alpha \geq 0$.
- (ii) Let $\alpha \in [0, 1]$, and $A, B \in \mathcal{B}(\mathcal{H})$ be such that $0 \leq A \leq B$. Prove that $0 \leq A^\alpha \leq B^\alpha$, i.e. that $x \mapsto x^\alpha$ is operator monotone on \mathbb{R}_+ . [Hint: You may want to use that for all $x \geq 0$ we have $x^\alpha = \frac{\sin(\alpha\pi)}{\pi} \int_0^\infty \frac{x}{x+\lambda} \frac{d\lambda}{\lambda^{1-\alpha}}$ for all $\alpha \in (0, 1)$.]
- (iii) Find a counterexample for (ii) when $\alpha > 1$.

Problem 39. Let $A : L^2([0, 1]) \rightarrow L^2([0, 1])$ be given by $Af(x) := xf(x)$ for a.e. $x \in [0, 1]$.

- (i) Prove that $A = A^*$, $\|A\| = 1$ and $\sigma(A) = [0, 1]$.
- (ii) Give the explicit action of $f(A) \in \mathcal{B}(L^2([0, 1]))$ for any bounded measurable function $f : [0, 1] \rightarrow \mathbb{C}$.
- (iii) For any $\psi \in L^2([0, 1])$ express $(\psi, f(A)\psi)$ as an integral with respect to the measure $\Omega \mapsto (\psi, E_\Omega\psi)$, where E denotes the projection valued measure given by A .

Problem 40. Let \mathcal{H} be a Hilbert space, let $A \in \mathcal{B}(\mathcal{H})$ be self-adjoint, and let E denote the projection valued measure given by A . Prove:

- (i) For any Borel set $\Omega \subset \sigma(A)$, the subspace $R(E_\Omega)$ is invariant under A .
- (ii) If $\Omega \subset \sigma(A)$ is closed, then $\sigma(A|_{R(E_\Omega)}) \subset \Omega$.