

## TOPOLOGIE IV – EXERCISE SHEET 2

**Universality of colimits in  $\mathbf{Gpd}$ .** Given a map  $f: X \rightarrow Y$  between groupoids, the pullback functor sits inside a commutative diagram

$$\begin{array}{ccc} \mathbf{Gpd}_{/Y} & \xrightarrow{f^*} & \mathbf{Gpd}_{/X} \\ \parallel & & \parallel \\ \mathbf{Fun}(Y, \mathbf{Gpd}) & \xrightarrow{f^*} & \mathbf{Fun}(X, \mathbf{Gpd}) \end{array}$$

Since colimits in those functor categories are formed pointwise, they are preserved by precomposition. In particular, the base change along  $f$  functor

$$f^*: \mathbf{Gpd}_{/Y} \rightarrow \mathbf{Gpd}_{/X}$$

preserves colimits. We say that colimits are *universal* in  $\mathbf{Gpd}$ . As an exercise, show that colimits in  $\mathbf{Cat}$  are not universal.

**Truncated maps.** Let  $\mathcal{C}$  be a category with finite limits. For  $k \geq -2$ , a map  $f: x \rightarrow y$  of  $\mathcal{C}$  is  $(k+1)$ -truncated if and only if the diagonal  $\Delta_f: x \rightarrow x \times_y x$  is  $k$ -truncated.

Using this characterisation, one can show that any left exact functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between categories with finite limits preserves truncatedness of objects and morphisms. If  $F$  is conservative, then it furthermore reflects truncatedness. For instance, since the forgetful functor  $\mathbf{Gpd}_* \rightarrow \mathbf{Gpd}$  is conservative and preserves limits, it preserves and reflects truncatedness.

**Lifting problems and finding sections.** Let  $\mathcal{C}$  be a category. For any cospan in  $\mathcal{C}$

$$x \xrightarrow{f} b \xleftarrow{p} e$$

whose limit exists, consider the following diagram

$$\begin{array}{ccccc} \mathrm{Hom}_{\mathcal{C}_{/x}}(x, f^*p) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(x, f^*p) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(x, e) \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow f_* \\ * & \xrightarrow{\mathrm{id}_x} & \mathrm{Hom}_{\mathcal{C}}(x, x) & \xrightarrow{u_*} & \mathrm{Hom}_{\mathcal{C}}(x, b) \end{array}$$

In particular, the two following lifting problems are equivalent

$$\begin{array}{ccc} & e & \\ & \nearrow & \downarrow p \\ x & \xrightarrow{f} & b \end{array} \quad \rightsquigarrow \quad \begin{array}{ccc} & x \times_b e & \\ & \nearrow & \downarrow f^*p \\ x & \xrightarrow{\quad} & x \end{array}$$

As a slogan, every lifting problem is equivalent to the problem of constructing a section.

**A criterion for connectivity.** For  $X$  a pointed groupoid and  $n \geq 0$ , the following are equivalent:

- (i)  $X$  is  $n$ -connected, or in other words  $\tau_{\leq n} X \simeq *$
- (ii)  $\mathrm{Hom}_*(X, Y) \simeq *$  for every pointed and  $n$ -truncated groupoid  $Y$
- (iii) for any  $m$ -truncated morphism  $f: Y \rightarrow Z$  between pointed groupoids, the map

$$f_*: \mathrm{Hom}_*(X, Y) \rightarrow \mathrm{Hom}_*(X, Z)$$

is  $(m - n - 1)$ -truncated.

*Proof.* Clearly (iii) implies (ii).

We now show that (ii) implies (i). For any groupoid  $Y$ , the evaluation map  $\text{ev}_*: \text{Hom}(X, Y) \rightarrow Y$  is an equivalence, since all of its fibers are contractible by assumption. Yoneda lemma then implies that  $\tau_{\leq n} X \simeq *$ .

Finally, we turn to the implication (i) implies (iii). Since the functor  $\text{Hom}_*(X, -)$  preserves limits, the map

$$f_*: \text{Hom}_*(X, Y) \rightarrow \text{Hom}_*(X, Z)$$

induced by composition with some  $f: Y \rightarrow Z$  is  $(k+1)$ -truncated if and only if the map

$$(\Delta_f)_*: \text{Hom}_*(X, Y) \rightarrow \text{Hom}_*(X, Z)$$

induced by the diagonal  $\Delta_f: Y \rightarrow Y \times_Z Y$  is  $k$ -truncated. By induction, it thus suffices to show that  $f_*$  is an equivalence when  $f$  is  $(n-1)$ -truncated. In this case, the fiber above  $u: X \rightarrow Z$  of  $f_*$  is the groupoid of pointed sections of  $u^*f: W \rightarrow X$

$$\begin{array}{ccc} \text{Hom}_{*/X}(X, u^*f) & \longrightarrow & \text{Hom}_*(X, Y) \\ \downarrow & \lrcorner & \downarrow f_* \\ * & \xrightarrow{u} & \text{Hom}_*(X, Z) \end{array}$$

By assumption  $X$  is connected, and thus  $u^*f$  has typical fiber  $F$  for some  $(n-1)$ -truncated groupoid  $F$ . Since  $\text{Aut}(F)$  is a reunion of connected components of the  $(n-1)$ -truncated groupoid  $\text{End}(F) \simeq \text{Hom}(F, F)$ , it is itself  $(n-1)$ -truncated and  $\text{BAut}(F)$  is  $n$ -truncated. Since  $X$  is  $n$ -connected, the classifying map  $X \rightarrow \text{BAut}(F)$  is constant, and pasting cartesian squares

$$\begin{array}{ccccc} X \times F & \longrightarrow & F & \longrightarrow & \text{BAut}_*(F) \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & * & \longrightarrow & \text{BAut}(F) \end{array}$$

yields an identification  $W \simeq X \times F$  over  $X$ . Using the base point of  $W$  to turn  $F$  into a pointed groupoid:

$$\begin{aligned} \text{Hom}_{*/X}(X, u^*f) &\simeq \text{Hom}_{*/X}(X, X \times F) \\ &\simeq \text{Hom}_*(X, F) \\ &\simeq * \end{aligned}$$

where the last step uses again that  $X$  is  $n$ -connected. Finally the map

$$f_*: \text{Hom}_*(X, Y) \rightarrow \text{Hom}_*(X, Z)$$

has contractible fibers, and therefore is an equivalence. This concludes the proof.  $\square$

**Exercise 1.** Observe that  $* \wedge (-)$  and  $S^0 \wedge (-)$  are left adjoint to the functors  $*: \text{Gpd}_* \rightarrow \text{Gpd}_*$  and  $\text{id}_{\text{Gpd}_*}$  respectively, and thus

$$* \wedge (-) \simeq * \quad \text{et} \quad S^0 \wedge (-) \simeq \text{id}_{\text{Gpd}_*}$$

Since  $(-) \wedge (-)$  preserves colimits in each variable, we have a pushout square

$$\begin{array}{ccc} \text{id}_{\text{Gpd}_*} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & S^1 \wedge (-) \end{array}$$

and a canonical natural isomorphism  $S^1 \wedge (-) \simeq \Sigma$ . In particular

$$\begin{aligned} \Sigma(- \wedge -) &\simeq \Sigma(-) \wedge (-) \\ &\simeq (-) \wedge \Sigma(-) \end{aligned}$$

since the smash product is commutative.

**Exercise 2.** Let  $X$  and  $Y$  two pointed groupoids being respectively  $m$ - and  $n$ -connected with  $m$  and  $n$  non-negative. For  $Z$  pointed and  $(m + n + 1)$ -truncated, the criterion above shows that  $\text{Hom}_*(Y, Z)$  is  $m$ -truncated and thus

$$\begin{aligned} \text{Hom}_*(X \wedge Y, Z) &\simeq \text{Hom}_*(X, \text{Hom}_*(Y, Z)) \\ &\simeq * \end{aligned}$$

Since this holds uniformly in  $Z$ , the smash product  $X \wedge Y$  must be  $(m + n + 1)$ -connected.

The result is false when  $m$  and  $n$  are allowed to be negative. For instance, smashing with the  $(-1)$ -connected space  $S^0 \vee S^0$  does not preserve connectedness.

**Exercise 3** ([DH21, lemma 2.17]). Mather's second cube lemma follows immediately from the universality of pushouts in  $\text{Gpd}$ . Let now  $\mathcal{C}$  be a category with universal pushouts. Recall that the endofunctor  $\Sigma: \mathcal{C}_* \rightarrow \mathcal{C}_*$  is defined by the following cocartesian square

$$\begin{array}{ccc} \text{id} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \Sigma \end{array}$$

in the category of endomorphisms of pointed objects  $\mathcal{C}_*$ .

Consider now the following cube

$$\begin{array}{ccccc} & & \Omega\Sigma & \longrightarrow & * \\ & \nearrow \xi_1 & \downarrow \xi_2 & \nearrow & \downarrow \\ \text{fib}(\text{id} \rightarrow \Sigma) & \longrightarrow & \Omega\Sigma & \longrightarrow & \Sigma \\ \downarrow & \nearrow & \downarrow & \nearrow & \downarrow \\ \text{id} & \longrightarrow & * & \longrightarrow & \Sigma \end{array}$$

where the top face is obtained by pulling back the bottom face along the base point  $* \rightarrow \Sigma$ . Since the bottom face is a pushout, the one must be as well by assumption.

By pasting cartesian squares all four other squares appearing in the cube are cartesian. Looking at the front face yields an identification

$$\begin{array}{ccc} \text{fib}(\text{id} \rightarrow \Sigma) & \xlongequal{\quad} & \text{id} \times \Omega\Sigma \\ \searrow \xi_2 & & \swarrow \text{pr}_2 \\ & \Omega\Sigma & \end{array}$$

above  $\Omega\Sigma$ . Finally, we obtain a pushout square

$$\begin{array}{ccc} \text{id} \times \Omega\Sigma & \xrightarrow{\text{pr}_2} & \Omega\Sigma \\ \downarrow a & & \downarrow \\ \Omega\Sigma & \longrightarrow & * \end{array}$$

where  $a$  is the composite

$$\text{id} \times \Omega\Sigma \xlongequal{\quad} \text{fib}(\text{id} \rightarrow \Sigma) \xrightarrow{\xi_1} \Omega\Sigma$$

Observe that  $a$  is in general only conjugated to  $\text{pr}_2$  by an automorphism of  $\text{id} \times \Omega\Sigma$  but is not homotopic to it. Indeed, base changing the defining pushout square of  $\Sigma$  along  $\text{pr}_1: \Sigma \times \Omega\Sigma \rightarrow \Sigma$  yields

$$\begin{array}{ccc}
\text{id} \times \Omega\Sigma & \xrightarrow{\text{pr}_2} & \Omega\Sigma \\
\downarrow \text{pr}_2 & & \downarrow \\
\Omega\Sigma & \xrightarrow{\quad \sqcup \quad} & \Sigma \times \Omega\Sigma
\end{array}$$

but  $\Sigma \times \Omega\Sigma$  is not terminal in general.

**Exercise 4** ([DH21, theorem 1.4]). Let  $\mathcal{C}$  be a category with finite products and pushouts. Remember that for two pointed objects  $x$  and  $y$ , the natural identification  $x \star y \simeq \Sigma(x \wedge y)$  is obtained by computing both sides as the colimit of the following diagram

$$\begin{array}{ccccc}
& * & \longleftarrow & * & \longrightarrow & * \\
\uparrow & & & \uparrow & & \uparrow \\
x & \longleftarrow & x \vee y & \longrightarrow & y \\
\downarrow & & \downarrow & & \downarrow \\
x & \longleftarrow & x \times y & \longrightarrow & y
\end{array}$$

In particular, the structure maps  $x \rightarrow x \star y$  and  $y \rightarrow x \star y$  both naturally factor through the point, which is not obvious from the definition.

The following diagram

$$\begin{array}{ccccc}
x \times y & \xrightarrow{\text{pr}_2} & y \\
\downarrow \text{pr}_1 & & \downarrow \\
x & \longrightarrow & * & \longrightarrow & x \star y \\
\downarrow & & \downarrow & & \downarrow \\
* & \longrightarrow & \Sigma x & \longrightarrow & \Sigma x \vee (x \star y)
\end{array}$$

gives canonical identifications

$$\begin{aligned}
\text{cofib}(\text{pr}_2: x \times y \rightarrow y) &\simeq \Sigma x \vee (x \star y) \\
&\simeq \Sigma x \vee \Sigma(x \wedge y)
\end{aligned}$$

When  $\mathcal{C}$  has universal pushouts, then combining this with the result from previous exercise we obtain natural isomorphisms

$$\begin{aligned}
\Sigma\Omega\Sigma &\simeq \text{cofib}(\text{pr}_2: \text{id} \times \Omega\Sigma \rightarrow \Omega\Sigma) \\
&\simeq \Sigma \vee \Sigma(\text{id} \wedge \Omega\Sigma) \\
&\simeq \Sigma \vee (\text{id} \wedge \Sigma\Omega\Sigma)
\end{aligned}$$

of endofunctors of  $\mathcal{C}_*$ . Plugging in the formula for  $\Sigma\Omega\Sigma$  then yields

$$\begin{aligned}
\Sigma\Omega\Sigma &\simeq \Sigma \vee (\text{id} \wedge (\Sigma \vee \Sigma(\text{id} \wedge \Omega\Sigma))) \\
&\simeq \Sigma \vee \Sigma(\text{id}^{\wedge 2}) \vee (\text{id}^{\wedge 2} \wedge \Sigma\Omega\Sigma)
\end{aligned}$$

and by induction

$$\Sigma\Omega\Sigma \simeq \bigvee_{i=1}^n \Sigma(\text{id}^{\wedge i}) \vee (\text{id}^{\wedge n} \wedge \Sigma\Omega\Sigma)$$

for all  $n \geq 1$ . In particular there is a well defined comparison morphism

$$\bigvee_{i \geq 1} \Sigma(\text{id}^{\wedge i}) \rightarrow \Sigma\Omega\Sigma$$

between endofunctors of  $\mathcal{C}_*$ .

Fix now  $X$  a pointed and connected groupoid. For  $n \geq 1$ , both the left map and the composite in the following diagram

$$\bigvee_{i=1}^n \Sigma(X^{\wedge i}) \longrightarrow \bigvee_{i \geq 1} \Sigma(X^{\wedge i}) \longrightarrow \Sigma\Omega\Sigma X$$

are the canonical inclusions, and therefore are both at least  $n$ -connected by the second exercise. By the cancellation property for connected morphisms, the right map is also  $n$ -connected. Since this holds for all  $n$ , we get the James splitting

$$\Sigma\Omega\Sigma X \simeq \bigvee_{i \geq 1} \Sigma(X^{\wedge i})$$

**Exercise 5.** Let  $\mathbb{K}$  be  $\mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ , and  $d := [\mathbb{K} : \mathbb{R}]$ . For  $n \geq 1$ , recall that  $\mathbb{P}^{n+1}(\mathbb{K})$  is obtained from  $\mathbb{P}^n(\mathbb{K})$  via the following cell attachment in  $\mathbf{Top}$

$$\begin{array}{ccc} S^{d(n+1)-1} & \xrightarrow{\gamma_n^{\mathbb{K}}} & \mathbb{P}^n(\mathbb{K}) \\ \downarrow & \lrcorner & \downarrow \\ D^{d(n+1)} & \longrightarrow & \mathbb{P}^{n+1}(\mathbb{K}) \end{array}$$

along the tautological spherical fibration  $S^{d(n+1)-1} \rightarrow \mathbb{P}^n(\mathbb{K})$ . But all objects at play are cofibrant and the left vertical map is a cofibration, so this pushout is furthermore an homotopy pushout. The cocartesian square in  $\mathbf{Gpd}$

$$\begin{array}{ccc} S^{d(n+1)-1} & \xrightarrow{\gamma_n^{\mathbb{K}}} & \mathbb{P}^n(\mathbb{K}) \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \mathbb{P}^{n+1}(\mathbb{K}) \end{array}$$

thus yields an identification

$$\mathrm{Th}(\gamma_n^{\mathbb{K}}) \simeq \mathbb{P}^{n+1}(\mathbb{K})$$

This implies

$$\mathrm{Th}(\gamma_{\infty}^{\mathbb{K}}) \simeq \mathbb{P}^{\infty}(\mathbb{K})$$

which is evident from the description of  $\gamma_{\infty}^{\mathbb{K}}$  as the universal principal  $(\mathbb{K}^{\times})$ -bundle.

#### REFERENCES

- [DH21] Sanath Devalapurkar and Peter Haine, *On the James and Hilton–Milnor splittings, and the metastable EHP sequence*, Documenta Mathematica **26** (2021), 1423–1464.