Mathematisches Institut
der Universität München
Prof. Dr. B. Pareigis

SS 2002
Set 10

Problem set for
Quantum Groups and Noncommutative Geometry

(37) We have seen that in representation theory and in corepresentation theory of quantum groups H such as $\mathbb{K}G$, $U(\mathfrak{g})$, $SL_q(2)$, $U_q(sl(2))$ the ordinary tensor product (in $\mathbb{K}$-$\mathcal{M}od$) of two (co-)reprensentations is in a canonical way again a (co-)reprensentation. For two H-modules M and N describe the module structure on $M \otimes N$ if

(a) $H = \mathbb{K}G$: $g(m \otimes n) = \ldots$ for $g \in G$;
(b) $H = U(\mathfrak{g})$: $g(m \otimes n) = \ldots$ for $g \in \mathfrak{g}$;
(c) $H = U_q(sl(2))$:
  (i) $E(m \otimes n) = \ldots$,
  (ii) $F(m \otimes n) = \ldots$,
  (iii) $K(m \otimes n) = \ldots$
  for the elements $E, F, K \in U_q(sl(2))$.

(38) Let G be a monoid.

(a) The *category of G-families of vector spaces* $\mathcal{M}^G = \prod_{g \in G} \mathcal{V}ec$ has families of vector spaces $(V_g | g \in G)$ as objects and families of linear maps $(f_g : V_g \to W_g | g \in G)$ as morphisms. The composition is $(f_g | g \in G) \circ (h_g | g \in G) = (f_g \circ h_g | g \in G)$. Show that $\mathcal{M}^G$ is a monoidal category with the tensor product

$$(V_g | g \in G) \otimes (W_g | g \in G) := (\oplus_{h,k \in G, hk=g} V_h \otimes W_k | g \in G).$$

Where do unit and associativity laws of G enter the proof?

(b) A vector space V together with a family of subspaces $(V_g \subseteq V | g \in G)$ is called *G-graded*, if $V = \oplus_{g \in G} V_g$ holds. Let $(V, (V_g | g \in G))$ and $(W, (W_g | g \in G))$ be G-graded vector spaces. A linear map $f : V \to W$ is called *G-graded*, if $f(V_g) \subseteq W_g$ for all $g \in G$. The G-graded vector spaces and G-graded linear maps form the category $\mathcal{M}^{[G]}$ of *G-graded vector spaces*. Show that $\mathcal{M}^{[G]}$ is a monoidal category with the tensor product $V \otimes W$, where the subspaces $(V \otimes W)_g$ are defined by

$$(V \otimes W)_g := \sum_{h,k \in G, hk=g} V_h \otimes W_k.$$

(c) Show that the monoidal category $\mathcal{M}^G$ of G-families of vector spaces is monoidally equivalent to the monoidal category $\mathcal{M}^{[G]}$ of G-graded vector spaces.

(d) Show that the monoidal category $\mathcal{M}^{[G]}$ of G-graded vector spaces is monoidally equivalent to the monoidal category of $\mathbb{K}G$-comodules $\mathcal{M}^{\mathbb{K}G}$. (Hint: Use the following constructions. For a G-family $(V_g|g \in G)$ construct a G-graded vector space $\hat{V} := \oplus_{g\in G} V_g$ (exterior direct sum) with the subspaces $\hat{V}_g := \mathrm{Im}(V_g)$ in the direct sum. Conversely if $(V, (V_g|g \in G))$ is a G-graded vector space then $(V_g|g \in G)$ is a G-family of vector spaces. For a G-graded vector space $(V, (V_g|g \in G))$ construct the $\mathbb{K}G$-comodule V with the structure map $\delta : V \longrightarrow V \otimes \mathbb{K}G$, $\delta(v) := v \otimes g$ for all (homogeneous elements) $v \in V_g$ and for all $g \in G$. Conversely let $(V, \delta : V \longrightarrow V \otimes \mathbb{K}G)$ be a $\mathbb{K}G$-comodule. Then construct the vector space V with den graded (homogenous) components $V_g := \{v \in V|\delta(v) = v \otimes g\})$.

(39) Let $(\mathcal{D}, \omega)$ be a diagram in $\mathcal{V}ec$. Let $\mathcal{D}$ be a monoidal category and ω be a monoidal functor. Then $(\mathcal{D}, \omega)$ is called a *monoidal diagram.*

Let $(\mathcal{D}, \omega)$ be a monoidal diagram $\mathcal{V}ec$. Let $A \in \mathcal{V}ec$ be an algebra. A natural transformation $\varphi : \omega \longrightarrow \omega \otimes B$ is called monoidal *monoidal* if the diagrams

$$\begin{tikzcd}
\omega(X) \otimes \omega(Y) \arrow[r, "\varphi(X)\otimes\varphi(Y)"] \arrow[d, "\rho"'] & \omega(X) \otimes \omega(Y) \otimes B \otimes B \arrow[d, "\rho\otimes m"] \\
\omega(X \otimes Y) \arrow[r, "\varphi(X\otimes Y)"] & \omega(X \otimes Y) \otimes B
\end{tikzcd}$$

and

$$\begin{tikzcd}
\mathbb{K} \arrow[r, "\cong"] \arrow[d] & \mathbb{K} \otimes \mathbb{K} \arrow[d] \\
\omega(I) \arrow[r, "\varphi(I)"] & \omega(I) \otimes B
\end{tikzcd}$$

commute.

We denote the set of monoidal natural transformations by $\mathrm{Nat}^{\otimes}(\omega, \omega \otimes B)$. Show that $\mathrm{Nat}^{\otimes}(\omega, \omega \otimes B)$ is a functor in B.

(40) Show that the *adjoint action* $H \otimes H \ni h \otimes a \mapsto \sum h_{(1)} a S(h_{(2)}) \in H$ makes H an H-module algebra.

Due date: Tuesday, 02.07.2002, 16:15 in Lecture Hall E41